Status, Trends, and Potential of Biological Communities of the Grand Calumet River Basin

submitted by

Philip B. Moy, USACE Project Manager
Richard L. Whitman, USGS Project Manager

Paul J. Gerovac, Meredith E. Becker, and Richard L. Whitman

Lake Michigan Ecological Research Station
Biological Resources Division
U.S. Geological Survey
1100 N. Mineral Springs Rd.
Porter, Indiana 46304

Once the downstream section of the Calumet River system that dominated most of Northwest Indiana, the now highly modified Grand Calumet River now constitutes one of the smallest major watersheds of the Calumet region. Though only 13 miles long and with almost no natural surface drainage area, it now flows through one of the most industrialized areas in the United States. Despite intensive .ID isd(2 eredith1e4(ielopum)9(nta, the Grand)]TJ0.0079 Tc -0

Industrial expansion experienced a lull with the onset of the Great Depression that extended through World War II (Cook and Jackson 1978). This lack of activity lasted into the 1950's, but the effects of industries already present were still apparent. In 1930, the Grand Calumet River was described by Peattie (1930) as a stagnant lagoon, an “open sewer” devoid of plant life, though bordering marshes still offered “favorable localities for plant growth.” Air pollution in 1966 was comprised of 41% fuel combustion (458,000 tons per year), 35% industry (392,000 tons per year), 22% transportation (241,000 tons per year), and 0.02% refuse disposal (Lerner 1977).

The establishment of industry created the landscape apparent today. The Grand Calumet River experiences the great force of pollution inherent in industry and urbanization. Years of un-managed pollution from rapid industrial growth is now buried in the sediments, and any ecological integrity of the system has been severely degraded.

POLLUTANTS

The 1972 Federal Water Pollution Control Act required the Indiana Stream Pollution Control Board (ISPCB) to issue permits to stream dischargers through the National Pollution Discharge Elimination System, and a chance at reviving the Grand Calumet River was granted. The steel mills removed solid and acid wastes from their effluent by installing catch basins and by using aeration and filtration techniques. Though there have been numerous reports of non-compliance, the NPDES system is recognized as a contributor to improved water quality.

Recognized pollutant sources include urban runoff, landfills, dumpsites, industrial effluent, and sewage treatment plants. The Grand Calumet River has a history of high levels of bacteria, nutrients, cyanides, lead, arsenic, cadmium, PCBs, phenols, oils, grease, chlorides, and other contaminants in the water and sediments. Combined sewer overflows from Gary, East Chicago, and Hammond sewage treatment plants flush raw sewage and fecal contamination into the Grand Calumet River and also Lake Michigan via the Indiana Harbor Canal.

The effects of these pollutants on humans and on the River ecology are great. Ammonia is released from stock yards and in the coking operation of steel production, and both ammonia and phosphorus are found in sewage, fertilizer, meat packing and industrial waste, and detergents. These two nutrients contribute to toxic algal blooms, increased aquatic plant and algal growth, and lower oxygen levels when they are flushed into Lake Michigan. Problems with Lake ecology and fisheries can occur when nutrient levels are high.

Other contaminants produced by industry also cause adverse health effects in humans. Hydrocarbons from refineries have an acute lethal toxicity and will decrease the dissolved oxygen concentration in water. Lead can cause convulsions, anemia, and kidney and brain damage. PCBs were once used in transformers and diodes, but they were banned in the 1970's. They resist degradation and remain in the environment and can cause vomiting, rashes, abdominal pain, temporary blindness, liver damage, cancer, and birth defects in humans.

Despite successful attempts to improve the River's water quality, the sediments will not be cleaned simply by changing current pollution practices. For over 100 years these contaminants have accumulated ee 4fe sediyspppply beteby l

wetlands have the potential to impact not only the Grand Calumet River watershed but also the southern Lake Michigan lakeplain, and ultimately the entire Great Lakes basin

The purpose of this chapter is to describe the special habitats along the Grand Calumet River corridor that support diverse biotic communities. The ecological significance of these sites will be outlined in the contexts of the River's watershed, the southern Lake Michigan lakeplain, and the Great Lakes basin. Potential impacts of dredging and of associated habitat restoration projects will also be discussed.

PRE-SETTLEMENT CONDITIONS

The southern Lake Michigan lakeplain

The post-glacial landscape of the southern Lake Michigan lakeplain is the product of constant change. For most of its history, regional physical processes such as climatic conditions, glacial mechanics, and fluctuating lake levels drove this dynamic system. The region's biotic communities have been influenced by three major biomes; eastern deciduous forest, tallgrass prairie, and boreal. The physically changing terrain together with the availability of diverse genetic material created an ecological rhythm that marked time with constantly evolving biotic communities.

The Great Lakes are relatively young in geologic terms; their history encompasses only the last several thousand years. During the Pleistocene Epoch climatic changes caused glaciers to advance southward across North America, extending as far south as Ohio and the Missouri River. As the glaciers flowed across the landscape, they carved out the Great Lakes basin. With the end of the Wisconsin Glaciation, some 11,000 years ago, the ice fields retreated from the region leaving behind the predecessors of the modern Great Lakes (Brown 1997).

The following geologic history of the lakeplain and formation of the Grand Calumet River are summarized from *Geologic History of the Little Calumet and Grand Calumet Rivers* *w*

Boreal and tundra flora established along the receding edge of the glacier. As the ice

While one segment of modern culture was beginning to understand the biological significance of the lakeplain ecosystem another was systematically destroying the ecological processes and patterns that had produced it. Pepoon laments the destruction of habitat that had already occurred by 1927: "... Very much of this rare flora has disappeared forever, owing to drainage, railroad building, sand hauling and filling, extension of urban communities, and the tremendous influx of factories. Much of the pond water remaining is practically sewage. The choice plants of these sloughs and marshes, as far east as Dune Park, have disappeared or are vanishing rapidly" (Pepoon 1927).

CURRENT CONDITIONS

The landscape of the Calumet region has changed dramatically during the past 100 years. The Grand Calumet River typifies these changes. It was once described as being more like a bayou than a river (Moore 1959), but it has since been channelized and redirected to flow into the Mississippi River basin. Today roughly ninety percent of its water comes from industrial and municipal discharges. The sandy soils of the river bed have been replaced by sediments contaminated with the residue of urban industrial activities. Despite these changes, aspects of the natural systems are still evident along the river corridor and throughout the watershed. Identifying and understanding the significance of the native species and community types is an integral part of assessing the ecological state of the river corridor.

The Indiana portion of the strandplain covers over 30,000 acres, and within that roughly 2,000 acres of ridge and swale still exist. Aerial photographs taken over the last sixty years document the physical transformation of the region from a natural system to an urban industrial complex. The first complete set of aerial photographs of the Calumet region date back to 1938. Undisturbed sections of ridge and swale topography are easily recognized by their distinctive linear pattern. The 1938 photographs show that the strandplain had already been divided into three distinct units. The city of Gary separated the Miller Woods area in the east from the central ridge and swale section in west Gary, Hammond, and East Chicago. The central ridge and swale section was divided from the Wolf Lake - George Lake area by the cities of Hammond, East Chicago, and Whiting. Shoreline armoring and fill, that would ultimately isolate the strandplain from Lake Michigan, had also begun. Although these areas were isolated from each other, there were still fairly large blocks of natural terrain in the Miller Woods and central ridge and swale area.

The Miller Woods area currently contains roughly one thousand acres of fairly contiguous native landscape including the Miller Woods Unit of the Indiana Dunes National Lakeshore, the City of Gary's Marquette Park, and private property. Over 430 species of native plants have been documented in the Miller Woods unit of the Indiana Dunes National Lakeshore. The dune complex north of the Lagoons supports panne and foredune communities. The ridge and swale complex to the south of the lagoons supports some of the highest quality black oak savanna in the Chicago Region (Wilhelm 1990). The Miller area

There are approximately one thousand acres of natural terrain left in the central ridge and swale area. The habitat has been fragmented into isolated pockets ranging in size from 170 acres to as little as five acres.

basin or have their best examples here. Thirty elements are ranked imperiled, of which thirteen occur exclusively or predominately within the basin, or have their best example there. Seventy-nine elements are ranked as rare, and 33 of these occur exclusively or are best represented within the basin.

Seven natural systems have been identified that support biodiversity in the Great Lakes basin. They are: open lake, coastal shore, coastal marsh, lakeplain, tributary and connecting channel, inland terrestrial upland, and inland wetland. Open lakes, coastal marsh, coastal shore and lakeplain are unique to the basin. Of these, coastal shore and lakeplain support a disproportionate amount of the basin's special biological diversity. Of the 61 Great Lakes dependent, globally significant elements (G1 or G2), 26 percent are supported by coastal shore, while 21 percent are supported by lakeplain systems.

The southern Lake Michigan region of the Great Lakes basin supports both lakeplain and coastal shore systems. The parabolic dunes that stretch along the shoreline from Gary, Indiana to southwestern Michigan are a type of coastal shore system. The exceptionally high levels of biological diversity in these dunes is underscored by the fact that the Indiana Dunes National Lakeshore has the third highest plant diversity of all national parks, despite having less than three percent of the total acreage of either of the top two (National Park Service 1987). The lakeplain system persists in a series of remnants of native landscape, scattered

noted as foraging and nesting habitat for regionally rare birds including; great egret (*Casmerodius albus*), black tern (*Chlidonias niger*) and black-crowned night heron (*Nycticorax nycticorax*).

At the easternmost end of the River, there are three sites that have been inventoried by the Indiana Natural Heritage Program: Miller Beach and Dunes, the USX Site and sections of Marquette Park. All three sites harbor significant natural areas that are associated with the Grand Calumet River Lagoons. All are a part of or are adjacent to the larger Miller Woods unit of the Indiana Dunes National Lakeshore, an approximately 900-acre remnant of native lakeplain landscape. This area is the transition zone between the Tolleston strandplain and the high dunes to the east.

The natural course of the Grand Calumet River was altered to accommodate the building of the USX steel mill in Gary. The relocation of the River channel isolated the section of the River east of the USX facility, and as a result, the Grand Calumet Lagoons formed. Tolleston Beach fans out from a single dune ridge in the east to about 50 dune capped beach ridges south of the Lagoons. The ridges have a linear form that parallels the lakeshore and are capped by moderate size dunes making them higher than those found further west on the strandplain. Windblown sand has divided sections of the swales into separate ponds. High parabolic dunes occur lakeward of the lagoons (Brown 1997). Prior to urban development, Miller Woods graded into the Greater Calumet Wetlands Site. Now the two sites are physically separated by the city of Gary.

The area surrounding the Lagoons can be separated into two units: the area to the north of the Lagoons consists primarily of foredune and dune complex, and the area to the south is a savanna complex (Wilhelm 1990). Natural Heritage data for Miller Woods and for the Grand Calumet Lagoons area is sparse. The northern unit is known to support two globally rare communities: panne and foredune. There is no Natural Heritage data available on community classification in the savanna complex to the south of the Lagoons.

looks at the absolute number of species with high C values that are found at a site (Swink and Wilhelm 1994).

The following is a summary of Native C values and Floristic Quality Index ratings from *Plants of the Chicago Region*:

> Based upon 15 years of application of this assessment system to all types of land in the Chicago region, certain patterns have emerged. We have found that the mean C values in the preponderance of our open land range from 0 to 2. In light of the fact that 89% of our native flora has a C value of 4 or greater, and a mean C value of 7.3, it is evident that the principal elements of our native systems are uninvolved in the Chicago region landscape today.
>
> The vast majority of land in the region registers I values [Floristic Quality Index] of less than 20 and essentially has no significance from a natural area perspective. Areas with I values higher than 35 possess sufficient conservatism and richness to be of profound importance from a regional perspective. Areas registering in the 50's and higher are extremely rare and of paramount importance they represent less than 0.5% of the land area in the Chicago region. (Swink and Wilhelm 1994)

Floristic quality assessments were performed for all the large dune and swale remnants in the Greater Calumet Wetlands Site and included as one component of the *Illinois-Indiana regional Airport Site Selection Report* in 1991 (Mierzwa et al. 1991). The assessments were updated in 1994 (Table 1). The two units of the Miller Woods site were surveyed in August of 1978 and again in August of 1989 (Table 2) (Wilhelm 1990).

The Grand Calumet River corridor

The U.S. Army Corps of Engineers has identified several reaches of the Grand Calumet River that are associated with specific proposed dredging projects. The land adjacent to four of these reaches supports significant pockets of biodiversity. At the easternmost end of the river, Miller Woods and Dunes surround much of the Lagoons reach.

Brunswick Savanna	68	38.81	4.71
Clark & Pine Addition # 1	92	44.00	4.59
Clark & Pine Addition # 2	152	75.03	6.09
Clark & Pine Nature Preserve	277	128	7.7
Clark & Pine East	212	88.58	5.74
Clark Junction	245	101.96	6.51
Clark Junction East	187	76.93	5.63
Cline Ave. Dune & Swale	106	53.52	5.20
DuPont Dune & Swale	226	76.10	5.06
Gibson Woods Nature Preserve	297	103.00	6.0
Ivanhoe Dune & Swale	272	89.62	5.43
Lakeshore Prairie	151	72.02	5.86
Tolleston Ridges	261	101.00	6.1
Tolleston Woods	93	44.59	4.62

Table 2 Floristic Quality Assessments for Miller Section

Miller Woods and Dunes Sites	Native Taxa	Floristic Quality Index	Native C Value
Unit A Foredune and Dune Complex	210	97.00	6.70
Unit B Savanna Complex	179	78.00	5.81

Miller Woods and Dunes

The remnant natural areas in the Miller area cover roughly 1,000 acres and include the Miller Woods and Dunes unit of the Indiana Dunes National Lakeshore, the City of

elements considered rare or limited to a unique niche within the Indiana Dunes National Lakeshore.

Foredune communities occupy the windward exposure of the first line of dunes from the lake shore. Characteristic plants of the foredune include: *Ammophila breviligulata, Andropogon scorparius, Artemisia caudata, Calamovilfa longifolia, Cirsium pitcheri, Cornus stolonifera baileyi, Lathyrus japonicus glaber, Populus detoides, Rhus aromatica arenaria,* and *Solidago racemosa gillamani.*

In the Miller dunes area, panne communities inhabit a series of interdunal depressions that form on the lee sides of the first or second line of dunes. The depressions intersect the groundwater table forming calcareous wetlands and ponds. Pannes are unique in floristic composition, containing species that grow nowhere else in the Chicago region or in the State of Indiana. Plants of the panne community include: *Aster ptarmicoides, Carex garberi,*

Calamagrostis canadensis, Cornus racemosa, Cornus stolonifera, Iris virginica, and *Osmunda regalis spectabilis* grow along the margins of the swales.

In areas with little or no canopy, common species along the crest of the ridge include *Andropogon scorparius, Arabis lyrata, Calamovilfa longifolia magna, Coreopsis lanceolota, Liatris aspera, Liatris cylindracea,* and *Lithospermum croceum*. At midslope *Andropogon gerardii* becomes the more dominant grass, and *Pedicularis canadensis, Senecio pauperculus,* and *Castilleja coccinea* are fairly common as well. *Calamagrostis canadensis* is the dominant grass of wet sand prairie areas. Other common plants *include Eupatorium perfoliatum, Liatris spicata, Oxypolis rigidor, Pycanthemum virginiana, Rosa palustris,* and *Solidago gramnifolia.*

The sand mining operation scraped away the dunes to the water table, creating habitat conditions similar to the natural pannes. Many panne associates are now found growing in these areas, including*: Aster ptarmicoides, Carex viridula, Gentiana crinata, Hypericum kalmianum, Liparis loeselii, Potentilla fruticosa, Rynchospora capillacea,* and *Sabatia angularis.*

The plant species list for Clark and Pine East contains 212 native species and 40 adventives, with a Floristic Quality Index of 78.23 and Native C Value of 5.03 with adventives. These numbers indicate that the site is of extreme importance as a relatively natural area in a region of highly degraded ecosystems.

DuPont Dune and Swale

There are approximately 170 acres of remnant dune and swale included in DuPont's corporate land holdings around its East Chicago plant. Four globally rare communities have been identified at the DuPont natural area: wet-mesic sand prairie, dry-mesic sand prairie, dry sand savanna and sedge meadow.

Approximately 50 acres of the DuPont area comprise a unique formation of dune and swale that has a natural surface water connection with the Grand Calumet River. Marshes along the river curve to the west and grade into linear swales. Near the river, the marshes are generally filled with cattails (*Typha ssp.*), common reed (*Phragmites australis*), and purple loosestrife (*Lythrum salicaria*). The swales support high quality wet prairie and sedge meadow communities. Species that are common throughout the swales include *Aster ptarmicoides, Calamagrostis canadensis, Carex stricta, Chelone glabra, Coreopsis tripteris, Eryngium yuccifolium, Eupatorium maculatum, Liatris spicata, Muhlenbergia glomerata, Pycanthemum virginianum, Scirpus pungens,* and *Scirpus validus creber*. Cattails, common reed, and purple loosestrife are well established in deeper parts of the swales.

The plant species list as of 1993 contains 226 native plant species and 35 adventives. It has a Floristic Quality Index rating of 70.8 and Native C value of 4.38 with adventives. The DuPont property survives as a rare and highly valuable remnant of the ridge and swale habitat-type. Floristic communities of this complexity are extremely rare within the Chicago region.

THE IMPACTS OF THE DREDGING PROCESS

The two major ecological risks of dredging are the possibility for ecologically harmful disposal of sediments and for sloughing of land adjacent to dredged areas. The primary objective in designing restoration measures to accompany dredging is to minimize these risks. Restoration measures of a slightly broader scope also merit consideration in the dredging-planning process. These would enhance the habitat value of high-priority natural areas adjacent to the Grand Calumet River and would serve as preventative measures against further degradation of these areas. Possible sites for various restoration activities are discussed in the section titled "Restoration," included below.

Sloughing of the banks could negatively impact natural areas by destroying existing high quality habitat or by opening degraded habitat to invasion by exotic species. The dunes that encompass the Grand Calumet Lagoons support unique biotic communities that could be negatively impacted by changes in the natural dynamics of dune development. The integrity of the dunes and ridges in this area should be protected during the dredging process. At Clark and Pine East, there is concern for the artificial bank that separates the river channel from the borrow pit at the southern end of the preserve. At DuPont, the high quality sections of the swales are currently separated from the river by a buffer zone of degraded marshes. Changes to these marshes could impact the quality of habitat throughout the swales.

When sediments are removed, their disposal should not impact natural areas or systems. High quality natural areas and key restoration sites should not be used for sediment disposal. Potential impacts to natural processes should also be studied if disposal sites are located near, though not within the borders of, natural areas. Impacts to be studied in this case include changes in run-off patterns and in ground water movement.

RESTORATION

A landscape by design

During the past one hundred years our industrial culture has dramatically restructured the landscape of the Calumet region to fit its needs. Dunes were leveled, wetlands drained and filled, and rivers channelized in order to make the area more suitable for urban development. The destruction of habitat disrupted ecological processes, thereby shattering the natural landscape. Today, only small fragments remain, and these are out of context in their current surroundings. Storm water that once recharged the groundwater table is now urban run-off, that is collected in sewer systems and piped away. Native species no longer range freely across the lakeplain to form and reshape communities. A small number of exotic species thrive in the wake of urban development and dominate the landscape.

In total, over ninety percent of the natural landscape of the Tolleston strandplain has now been destroyed. The remaining fragments are the last refuge for the biotic communities native to this area. Significant ecological interaction is restricted, for the most part, to these fragments. They hold the only biological reserves of native species sufficient to fuel any

future reintroduction efforts, and they are also the last models available of the natural ecological systems of the region function.

Finding relevance for these natural areas in the current landscape is difficult. The natural systems that controlled the ecology of the lakeplain have been eclipsed by conditions created by human activity. The industrial landscape has created a new set of environmental conditions that will shape the future of these remnants as much as the natural systems shaped their past. Restoration of the River as a functional ecosystem presents this basic problem: remnants of the pre-industrial landscape offer the best opportunities for conservation and restoration of ecological health to the system, yet the natural processes that created and kept them dynamic have been changed forever. If we are to conserve our natural heritage for future generations we must account for the long term viability of these sites in our land use planning. Preservation and stewardship of these lands will require a conservation ethic that reaches beyond their immediate borders, making ecological restoration a process of integrating biological diversity into the broader landscape through planning and design.

Until recently, little attention was paid to conservation in land use planning in the Calumet Region. The first attempt to develop a Coastal Zone Management Plan in the mid 1970s produced a list of high quality natural areas in Lake County. These inventories combined with a growing awareness of the value of our biological heritage led to the purchase and dedication of a handful of these sites as Nature Preserves. Until then, these tracts were areas that had escaped urban industrial development by chance. The protection of Hoosier Prairie, Gibson Woods, Tolleston Ridges, Clark and Pine, and Ivanhoe Dune and Swale was the first successful effort at systematic conservation of natural resources in northern Lake County. By establishing these preserves, a means for maintaining biological diversity in the industrialized heart of the Calumet Region was developed.

Nature preserves are created to protect the highest quality examples of natural communities; their intrinsic value as a natural area controls their cultural land use. They are like gardens that operate independent of the surrounding landscape. At present, our best efforts at ecological management keep the natural systems functioning only within the borders of individual preserves. This conservation is severely restricted in its range.

Preservation of natural lands and systems will require a conservation ethic that reaches beyond the immediate borders of designated nature preserves. Such an ethic will challenge the assumption that there is a dichotomy between conservation and human demands on the land. Without ignoring or displacing the human inhabitants of the region, thoughtful land planners will find ways to expand the available habitat options for native species, and will thereby re-introduce natural ecosystem functioning to areas currently devoid of natural-area value. By creating buffers around existing natural areas, developing biological corridors, and replacing exotics with native species on properties not solely dedicated to conservation, we can begin to re-unify our human landscape with the natural ecological system. Such a wide-ranging effort will require that conservation and restoration activities be coordinated with activities that enhance the economic welfare and community development of the region. Plans will be implemented not by narrow constituencies, but through partnerships between government agencies, private landowners, and conservation organizations.

The western extreme of this reach runs through Clark and Pine East Nature Preserve. The eastern end includes parts of the Grand Calumet Lagoons and Miller Woods. The middle section runs through residential and industrial areas, and it currently has little value as a natural area.

USX section of the Grand Calumet Lagoons

The juxtaposition of the "moonscape" of the USX slag piles with the fragile beauty of the natural dune complex in this landscape is a striking reminder of the restoration challenges we face. Gerould Wilhelm (1990) aptly described the problem: "Half of the westernmost pond, still the richest and most stable Panne in existence, has been unceremoniously obliterated by a large slag pile which remains to this day, and should probably continue to remain in place lest the activity of removal obliterate the Panne altogether." Ideally, restoration of this site would include removal of fill, repair of historic degradation, and creation of a buffer between the remaining natural area and the industrial site.

Clark and Pine East Nature Preserve

The preserve serves as a direct link between the River and some of the highest quality natural areas in the state. The cluster of sites around Clark and Pine are legendary among botanists in the Chicago region. Clark and Pine East is owned by the Indiana Department of Natural Resources (IDNR) and managed by the Division of Nature Preserves (DNP). Restoration projects in this area would have to be coordinated with IDNR management objectives for the site.

Restoration of this property is a high priority. Large sections of the preserve have suffered disturbances that have left the ground open for invasive exotic species. Along with common reed and purple loosestrife, European buckthorn is a problem at this preserve. Buckthorn forms dense stands that must be removed manually. Fire suppression has also caused large areas of the preserve to become overgrown. Department of Nature Preserves staff have conducted controlled burns at the site for the past two years. Volunteers have worked under the direction of DNP and Nature Conservancy staff to carry out restoration ovhat dare lDNRve

enhance the potential habitat value of nearby, smaller, more degraded natural areas. Formal protection and management of this site will be crucial to any habitat restoration effort in the DuPont reach of the River.

Outside the boundaries of the DuPont property, but within the DuPont reach, are Siedner Dune and Swale Nature Preserve, the Beamsterboer property, the Harbison Walker property and the USS Lead Refinery. All of these are sites with potential for restoration projects.

The stretch that extends through the East Chicago Sanitary District, Roxanna Marsh, and DuPont reaches of the River holds the largest local complex of riparian wetlands. They are divided only by Kennedy Ave., which crosses the DuPont reach, and Indianapolis Blvd., which crosses between the East Chicago Sanitary District reach and the Roxanna Marsh reach. Most of these are cattail marshes that have become infested with common reed and

brownfield site. Creating a buffer on this degraded land for the adjacent natural area would protect and expand habitat for native species. Restoration activities would include: removal of old roads, parking areas, and building foundations, re-introduction of native plant species, and control of exotics. Currently there are large stands of *Andropogon scorparius*, and smaller patches of *Panicum virgatum, Calamovilfa longifolia, Andropogon gerardii*, and *Sorghastrum nutans* scattered throughout the field. Planting a mixture of these grasses and associated species such as *Asclepias tuberosa, Coreopsis lanceolata, Euphorbia corallata, Heianthus helianthoides, Liatris aspera,* and *Lupinus perennis* would begin the process of re-establishing productive habitat to this part of the site.

The developed areas of the DuPont property are built above the level of the River. The riverbank here is steep. At its base, a series of degraded marshes are found growing along the edge of the River channel. The steep banks are currently subject to high rates of erosion, which could likely be decreased by the planting of native species along these banks. Improvements could also be made to the plant communities of the marshes, which are currently dominated by cattailjT

LITERATURE CITED

Bacone, J.A. October 1979. Shell Oil Dune and Swale: A report on a natural area. Division of Nature Preserves, Indiana Department of Natural Resources, State of Indiana.

Brown, S.E. 1997. Geologic history of the Little Calumet and Grand Calumet Rivers. Indiana Geologic Survey, Indiana University

Crispin, S. and D. Rankin. January 1994. The conservation of biological diversity in the Great Lakes ecosystem: issues and opportunities. The Nature Conservancy Great Lakes Program, Chicago, Illinois, USA.

Moore, P. A. 1959. The Calumet Region: Indiana's last frontier. Indiana Historic Bureau, Indianapolis, Indiana, USA.

National Park Service. 1987. NPFLORA/COMMON: Coverage by Acreage of National Park Service Units, Unpublished NPS document.

Pepoon, H. S. 1927. An annotated flora of the Chicago area. Chicago Academy of Science, Chicago, Illinois, USA.

Swink, F. and G. Wilhelm. 1994. Plants of the Chicago region. Indiana Academy of Science. The Morton Arboretum. Lisle, Illinois, USA.

Mierzwa, K.S., S. Culberson, K.S. King, and C. Ross. 1991. Illinois-Indiana regional airport study: Biotic communities. Technical Paper No. 7, Appendix E, Volume II. TAMS Consultants, Inc. Chicago, Illinois, USA.

Wilhelm, G.S. 1990. Special vegetation of the Indiana Dunes National Lakeshore. The Morton Arboretum. Lisle, Illinois, USA.

NATIVE FLORA OF PRE-SETTLEMENT TIMES

Extirpated species

Peattie (1930), in his publication *Flora of the Indiana Dunes*, estimated that 1,400 plant species occurred in the Indiana Dunes area. Nineteen of these species have not been seen in Indiana since Peattie's sightings, and they have now been classified as extirpated species (species that had not been seen in Indiana for 50 years) by Aldrich et al. (1986). A list of the extirpated species is given below:

Scientific name	Common name
Betula populifolia	gray birch
Carex scabrata	
Corallorrhiza trifida	coral root
Gerardia pedicularia ambigens	clammy flase foxglove
Hemicarpa drummondii	
Hippuris vulgaris	Mare's tail
Lechea stricta	bush pinweed
Lemna perpusilla	least duckweed
Linnaea borealis	twin flower
Lonicera canadensis	American fly honeysuckle
Oryzopsis pungens	short-horned rice grass
Panicum lucidum	bog panic grass
Psilocarya nitens	bald rush
Pyrola secunda	one-sided shinleaf
Scleria reticularis	netted nut rush
Shepherdia canadensis	Russet buffaloberry
Trillium cernuum macranthum	nodding trillium
Utricularia resupinata	small purple bladderwort

Historic plant community types

Bacone et al. (1980) reconstructed the pre-settlement vegetation characteristics of northwestern Indiana by analyzing land survey records that were compiled between 1829 and 1834. Of the community types recognized by this study, aquatic communities, marshes, swamps, bottomland forests, beach communities, and pannes are considered as wetland community types. Summary descriptions of these community types are given below.

Aquatic communities consist of macrophytes and phytoplankton in standing or running water. The 1829-1834 survey record noted *Polygonum* spp., *Nuphar advena*, and *Nymphaea tuberosa* as typical macrophytes (Bacone et al. 1980). Although there is no documented list of phytoplankton in the pre-settlement times, the phytoplankton communities were likely dominated by such genera in the Division Chlorophyta as *Chlamydomonas, Oedogonium, Spirogyra*, and *Volvox*. Since European settlement, species composition of this community has been changed significantly by numerous human activities

such as loading of nutrients, silts and other pollutants, alteration of local hydrology through channelization, drainage and filling of the river and its adjacent wetlands. Current species composition of aquatic community will be discussed later.

Marshes are probably the most prominent plant community type along the Grand Calumet River. Marshes are open (non-forested) wetlands that are dominated by sedges and/or grasses. On the wet end of the moisture gradient, sedge meadow is found directly adjacent to aquatic communities. At the other extreme, wet prairie overlaps with the borders of mesophytic prairies and savanna complex. As is indicated by its name, sedge meadow is characterized by abundant sedges (Cyperaceae); wet prairies are dominated by grasses (Poaceae). Typical plant species in this community type include *Aster puniceus firmus, Bidens coronata tenuiloba, Carex aquatilis altior, Decodon verticillatus, Polygonum punctatum*, and *Scirpus acutus*. Historically, periodic fires, both natural and man-made, have been a crucial factor in maintaining marshes because fires prohibit the invasion of woody shrubs and trees. Since European settlement, however, most marshes have been heavily disturbed by drainage, by invasion of woody species facilitated by artificial fire suppression, and by intentional or accidental introduction of alien species. Shrub carr is a marginal type of marsh community, found along the borders of swamp and marsh. Typical woody species include *Alnus* spp. and *Salix* spp. According to the 1829-1834 survey record, woody species (e.g., *Populus tremuloides* and *Populus grandidentata*) were less abundant in presettlement than the present times. However, significant alterations in the level of the water table, and a long-lasting policy of fire suppression have resulted in significant invasion of overgrown shrubs into marshes, sedge meadows, and wet prairies.

Swamps are forested wetlands where the water level is maintained near or at the surface of the substrate by ground water or by rain. In the land survey records, the swamps in northwestern Indiana were classified into three major types. Coniferous swamps occurred only in dune-swale systems, and were dominated by *Pinus banksiana* and *Thuja occidentalis*. Timbered swamps, now classified as green ash (*Fraxinus pennsylvanica subintegerrima*) swamps, are inhabited by *Populus deltoides, Platanus occidentalis, Fraxinus nigra, Acer saccharinum*, and *Ulmus* spp. This type of swamp occurred along the Kankakee River. However, extensive draining and logging have destroyed much of this community in this century. Shrub swamps overlap with shrub carrs at the edges of marshes and of timbered swamps. The species compositions of these two community types are, as would be expected, quite similar. Shared species include *Alnus rugosa americana* and *Salix* spp. This type of swamp, like the shrub carrs, has become more common than during pre-settlement times as a result of long-term drainage and fire suppression.

Bottomland forests consist of several woody strata underlain by herbaceous cover, and they are located along the banks of water courses. They are characterized by annual deposition of silt during flooding. Major canopy species in these forests include *Acer negundo, Acer saccharinum, Platanus occidentalis, Salix nigra, Ulmus americana,* and *Ulmus rubra*. Small scattered patches of bottomland forest still occur today, although most of these have been heavily disturbed.

Beach communities are narrow specialized strips adjacent to the littoral zone of the lake shore, and they are dominated by *Ammophila breviligulata*. Pannes are moist interdunal

depressions in calcareous sands on the lee-side of a dunes, containing such species as *Aster ptarmicoides* and *Carex* spp.

EXISTING WETLAND PLANT COMMUNITIES

Recent vegetation survey and classification

Bowles et al. (1990) listed ten natural communities within the boundary of Indiana Dunes National Lakeshore (INDU). Six of these communities are considered wetland communities (Table 1). These communities include beach, forested fens, graminoid fens, forested bogs, flatwoods, and graminoid wetlands. Wilhelm (1990) also described 11 community types along the gradients of moisture and arborescent development. Among these communities, swamp complex, bog, marsh complex bottomovBslsurunit1.

Table 3. In this table, for example, Wilhelm's (1990) "marsh complex" is separated into marsh, fen, and sedge meadow. The "wet prairie" is a part of "marsh complex" in the Wilhelm's classification, but the proposed classification places it in the category of "prairie." In addition, "hydromesophytic forest" is a part of Wilhelm's "swamp complex," but it is placed the categories of "forest" and "shrub swamp" in the proposed classification system.

Table 1. Eleven natural plant communities listed by Bowles et al. (1990) in Indiana Dunes National Lakeshore. Wetland types are signified by *italics*.

COMMUNITY	CHARACTERISTICS
Beach/Foredune	Wave actions and sandy substrate Annuals and rhizomatous perennial plants
Dune Complex	Cycle of sand erosion in steep topography & blow-outs
Sand Savanna	Dominant presettlement plant community type on irregular dune topography Open dune conditions with frequent fires
Sand Prairie	Flat topography with frequent burns

	High species diversity in openings and pools
Flatwoods	Wet mineral soils or seepages with high pecies diversity

Figure 1. Eleven plant communities of Indiana Dunes National Lakeshore, depicted by Wilhelm (1990).

Table 2. Forty-nine plant community types of Indiana coastal zone listed by Kurz et al. (1978). Wetland types are signified by *italics*

CATEGORY	COMMUNITY TYPE
Forest	Dry-mesic upland forest Mesic upland forest Wet-mesic upland forest Dry dune forest Dry-mesic dune forest Mesic floodplain forest *Wet-mesic floodplain forest* *Wet floodplain forest* *Flatwoods*
Prairie	

Aquatic (continued)	*Graminoid fen* *Low shrub fen* *Tall shrub fen* *Forested fen* *Sedge meadow* *Panne* *Seep* *Calcareous seep* *Sand seep* *Spring*
Primary	*Beach* Foredune/Blowout

Figure 2. Areas of special conservation needs (Roxanna Marsh, DuPont Tract, Clark and Pine East Nature Preserve-Bongi Ponds, and Grand Calumet River Lagoon in Miller Woods), and fifteen proposed sediment disposal sites.

Special conservation needs

Among the riparian wetlands of the Grand Calumet River, Roxanna marsh, DuPont tract, Clark and Pine East nature preserve (also known as Bongi Pond), and Grand Calumet Lagoons are especially in need of conservation because they are considered as high quality habitats for endangered, threatened and/or rare animals and plants. After a compilation of the existing plant species records (Wilhelm 1990, Mierzwa et al. 1991, IDNR unpublished), a total of 665 plant species were found in DuPont tract, Clark and Pine East, and Miller Woods. Of these 665 species, 16 species were classified as endangered species (inhabit between 1 and 5 extant sites). Another 16 were classified as threatened species (inhabit between 6 and 10 sites), and 21 were classified as rare species (inhabit between 11 and 20 sites) (Aldrich et al. 1986). Key aspects of the areas with high conservation value are discussed below. The possibility of conserving and/or restoring populations of endangered, threatened, and rare species is also discussed individually for each area.

Table 3. Synthesized wetland plant community types under the classification scheme of the Indiana Department of Natural Resources.

Communities by Kurz et al. (1978)	*Communities by Bowles et al. (1990) and Wilhelm (1990)*
Wet-mesic floodplain forest Wet floodplain forest	Bottomland forest* Hydromesophytic forest in the Swamp Complex*
Flatwoods	Flatwoods in the Swamp Complex* Flatwoods**

meadow *Bidens comosa* swamp tickseed *Dryopteris*
madder *Hypericum virginicum fraseri* marsh St.
John’s

DuPont tract, owned by DuPont Chemical Company, is located to the east of Cline Avenue in Hammond, Indiana (Figure 2). Despite extensive industrial development during the 1950s and the 1970s, significant portions of classical "dune and swale systems" are preserved in this property. A total of 205 native species were documented here by TAMS (Mierzwa et al. 1991) and IDNR (unpublished) (Attachment 1), and four of these species are listed as threatened or rare species by IDNR, as follows:

Scientific name	Common name	Status
Carex bebbi	Bebb's sedge	threatened
Eriophorum angustifolium	narrow-leaved cotton grass	threatened
Baptisia leucantha	white wild indigo	rare
Betula papyrifera	paper birch	rare

Clark & Pine East Nature Preserve, owned by the State of Indiana, is located in the southeastern corner of the East Chicago-Gary Regional Airport property (Figure 2). Like the DuPont tract, this area is a classical example of a dune and swale system. TAMS (Mierzwa et al. 1991) and IDNR records for the area (unpublished) list 213 native species (Attachment 1), along with five endangered, five threatened, and ten rare species, as shown below:

Scientific name	Common name	Status
Carex brunnescens		endangered
Carex richardsonii		endangered
Gerardia skinneriana		endangered
Lycopus americanus	common water horehound	endangered
Sceleria pauciflora caroliniana		endangered
Aster ptarmicoides	stiff aster	threatened
Carex bushii		threatened
Carex crawei		threatened
Carex garberi	false golden sedge	threatened
Eleocharis geniculata	panne spike rush	threatened
Arctostaphylos uva-ursi coactilis	arctic bearberry	rare
Aristata intermedia	false arrow feather	rare
Carex aurea	golden sedge	rare
Cypripedium calceolus parviflorum	small yellow lady's slipper	rare
Cypripedium calceolus	large yellow lady's	rare

Tofieldia glutinosa	rare

Aggressive expansions of invasive species, both alien and native, pose serious threats to populations of native species in this reach/tract. The river banks in these natural areas are heavily infested with *Phragmites communis berlandieri*, *Lythrum salicaria*, *Typha angustifolia*, and *Typha latifolia*. Spread of these species is generally facilitated by their effective pollination systems, seed dispersal (mostly by wind), breeding systems (e.g., facultative apomixis), and, in many circumstances, rapid vegetative growth by "root-suckering" or "stem sprouts" (Baker 1986). These characteristics make this group of species very likely to continue expand aggressively in the wetlands of this tract, thereby out-competing native species. Forty alien species have already been found in the natural areas of DuPont and Clark and Pine East tracts (Attachment 6). Invasive species, both native or alien, should be eradicated from these tracts as soon as possible.

Most wetlands of the Grand Calumet Lagoons are located within the Miller Woods area, which is a part of INDU (Figure 2). This is home to what probably is the best-preserved and most diverse flora in northwestern Indiana. Wilhelm (1990) documented 559 species including 11 endangered, 12 threatened, and 12 rare species. These are listed below. The integrity of natural communities in the Grand Calumet Lagoons and Miller Woods tract, like that of natural areas at DuPont and at Clark and Pine East is threatened by the presence of many invasive species. Eighty-six alien species have been documented in the Grand Calumet Lagoons and Miller Woods area (Attachment 6).

Scientific name	Common name	Status
Equisetum variegatum	small scouring rush	endangered
Glyceria borealis	northern manna grass	endangered
Juncus scirpoides	round-headed rush	endangered
Ludwigia sphaerocarpa deamii	round-fruited loosestrife	endangered
Panicum dichotomiflorum	knee grass	endangered

LITERATURE CITED

Abt, S. R., C. C. Watson, J. C. Fisbenich, and M. R. Peters. 1995. Bank stabilization and habitat aspects of low-flow channels. Land and Water January/February:11 - 13.

Aldrich, J. R., J. A. Bacone, and M. A. Homoya. 1986. List of extirpated, endangered, threatened and rare vascular plants in Indiana: an update. Indiana Academy of Science **95**:413-419.

Bacone, J. A., R. K. Campbell, and G. Wilhelm. 1980. Presettlement vegetation of Indiana Dunes National Lakeshore. Pages 156-191 *in* Proceedings of Second Annual Symposium on Scientific Research in National Parks, National Park Service, Washington, D. C., USA.

Baker, H. G. 1986. Patterns of plant invasion in North America. Pages 44-57 *in* H. A. Mooney and J. A. Drake, editors. Ecology of biological invasions of North America and Hawaii. Springer-Verlag, New York, New York, USA.

Bowles, M. L., M. M. DeMauro, N. Pavlovic, and R. D. Hiebert. 1990. Effects of anthropogenic disturbances on endangered and threatened plants at the Indiana Dunes National Lakeshore. Natural Areas Journal **10**:187-200.

Cole, G. A. 1979. Textbook of limnology. The C. V. Mosby Company, St. Louis, Missouri, USA. 426 p.

Cowles, H. C. 1899. Ecological relations of the vegetation on sand dunes of Lake Michigan. Botanical Gazette **27**:95-117, 167-202, 281-308, 361-388.

Driver, T. 1993. The Transco Trail of the Hackensack Meadowlands. Land and Water September/October:28-32.

Indiana State Department of Natural Resources (IDNR). Unpublished. List of endangered, threatened, and rare plants in DuPont tract and Clark and Pine East Nature Preserve. Indiana Department of Natural Resources, Indianapolis, Indiana, USA.

Henderson, N. R., and J. N. Long. 1984. A comparison of stand structure and fire history in two black oak woodlands in northwestern Indiana. Botanical Gazette **145**:222-228.

Kurz, D. R., G. A. Paulson, D. W. Morgan, and J. W. Burling. 1978. Inventory of natural areas in the Indiana coastal zone. Natural Land Institute and Indiana Department of Natural Resources, Indianapolis, Indiana, USA. 119 p.

Mierzwa, K. S., S. D. Culberson, K. S. King, and C. Ross. 1991. Illinois-Indiana Regional Airport - site selection, environmental assessment, and master plan. Technical Paper No. 7, TAMS Consultants, Chicago, Illinois, USA.

Mitsch, W. J., and J. G. Gosselink. 1986. Wetlands. Van Nostrand Reinhold Company, New

Attachment 1. List of plant species in DuPont tract, Clark and Pine East Nature Preserve, and Miller Woods.

Legends: DT - DuPont Tract
CP - Clark and Pine East Nature Preserve (Bongi Ponds)
MW - Miller Woods (Grand Calumet River Lagoon)
A - Alien species
E - Endangered sepces
T - Threatened species
R - Rare species

Scientific Name	Common Name	Location DT	CP	MW	Class
Abutilon theophrasti	velvetleaf			X	A
Acer negundo	boxelder	X		X	
Acer platanoides	Norway maple			X	A
Acer rubrum	red maple		X	X	
Acer saccharinum	silver maple	X	X	X	
Achillia millefolium	yarrow	X		X	A
Agropyron repens	quack grass			X	A
Agropyron smithii	western wheat grass			X	A
Agropyron trachycaulum unilaterale	slender wheat grass		X	X	
Agrostis alba	redtop grass	X	X	X	A
Agrostis hyemalis	tickle grass			X	
Ailanthus altissima	tree-of-heaven	X			A
Alestris farinosa	colic root			X	
Alisma subcordatum	common water plantain		X	X	
Alisma triviale	large-leaved water plantain	X		X	
Alliaria officinalis	garlic mustard			X	A
Allium cernuum	nodding wild onion			X	
Althaea rosea	hollyhock	X			A
Ambrosia artemisiifolia elatior	common ragweed	X		X	
Ambrosia psilostachya coronopifolis	western ragweed			X	A
Ambrosia trifida	giant ragweed	X	X	X	
Amelanchier arborea	serviceberry			X	
Amelanchier interior	dwarf shadbush			X	
Amelanchier laevis	Allegheny shadbush			X	
Ammophila breviligulata	Merram grass			X	
Amorpha canescens	lead plant	X			
Amphicarpa bracteata	hog peanut			X	
Andropogon gerardii	big bluestem	X	X	X	
Andropogon scoparius	little bluestem	X	X	X	
Anemone canadensis	meadow anemone			X	
Anomone cylindrica	timbleweed	X	X	X	
Antennaria neglecta	cat's foot		X		
Antennaria plantaginifolia	pussy toes		X	X	
Anthriscus scandicina	chervil			X	A
Apios americana	ground nut	X		X	
Apocynum androsaemifolium	spearding dogbane			X	
Apocynum cannabinum	Indian hemp	X	X	X	
Apocynum sibiricum	Indian hemp	X	X	X	

Cirsium muticum	swamp thistle	X	X	X	
Cirsium pitcheri	pitcher thistle			X	T
Cirsium vulgare	bull thistle	X	X	X	A
Cladium mariscoides	twig rush	X	X	X	
Comandra richardsiana	false toadflax	X		X	
Commelina communis	common day flower	X			A
Commelina erecta deamiana	savanna day flower			X	
Conium maculatum	poison hemlock	X			A
Convolulus arvensis	field bindweed	X			A
Convolulus sepium	hedge bindweed	X		X	
Coreopsis lanceolata	sand coreopsis	X	X	X	
Coreopsis palmata	prairie coreopsis	X	X		
Coreopsis tripteris	tall coreopsis	X	X	X	
Corispermum hyssopifolium	common bugseed		X		
Cornus obliqua	pale dogwood	X	X	X	
Cornus racemosa	gray dogwood	X	X		
Cornus rugosa	speckled dogwood		X		T
Cornus stolonifera	red-osier dogwood	X	X	X	

Eleocharis compressa	flat-stemmed spike rush			X	
Eleocharis elliptica	golden-seeded spike rush	X	X	X	
Eleocharis engelmanni				X	
Eleocharis geniculata	panne spike rush		X	X	T
Eleocharis intermedia	matted spike rush			X	
Eleocharis olivacea	loose-sheathed spike rush			X	
Eleocharis pauciflora fernaldii			X	X	R
Eleocharis smallii					X
Elodea canadensis	common waterweed				X
Elodea nuttallii					X
Elymus canadensis	Canadian wild rye		X	X	X
Epilobium glandulosum adenocaulon	northern willow herb				X
Equisetum arvense	horsetail	X	X	X	
Equisetum hymale affine	tall scouring rush	X	X	X	
Equisetum hymale intermedium	smooth scouring rush		X		X
Equisetum trachyodon				X	
Equisetum X ferrissii				X	
Equisetum variegatum					

Sisymbrium altissimum	tumble mustard			X	*A*	
Sisyrinchium albidum	common blue-eyed grass	X	X	X		
Sium suave	water parsnip			X	X	
Smilacina racemosa	feathery false Solomon's seal	X		X		
Smilacina stellata	starry false Solomon's seal	X	X	X		
Smilax lasioneura	common carrion flower	X		X		
Smilax rotundifolia	green brier				X	
Smilax tamnoides hispida	bristly green brier	X	X			
Solanum americana	black nightshade			X	A	
Solanum dulcamara	bittersweet nightshade	X	X	X	A	
Solidago altissima	tall goldenrod		X	X	X	
Solidago caesia	blue-stem goldenrod			X	X	
Solidago gigantia	late goldenrod			X	X	
Solidago graminifolia media	smooth grass-leaved goldenrod			X	X	
Solidago graminifolia nutallii	rough grass-leaved goldenrod		X	X	X	
Solidago gymnospermoides	shiny grass-leaved goldenrod			X	X	
Solidago juncea	early goldenrod			X		
Solidago missouriensis fasciculata	Missouri goldenrod			X	X	
Solidago nemoralis	old field goldenrod		X	X	X	
Solidago ohiensis	Ohio goldenrod		X	X		
Solidago ptarmicoides	stiff goldenrod			X	R	
Solidago racemosa gillmani	dune goldenrod			X		
Solidago riddellii	Riddell's goldenrod			X		
Solidago rigida	stiff goldenrod			X		
Solidago rugosa	rough goldenrod		X	X		
Solidago sempervirens	seaside goldenrod		X		A	
Solidago speciosa	showy goldenrod	X	X	X		
Solidago tenuifolia	slender-leaved goldenrod			X		
Solidago uliginosa	bog goldenrod		X			
Sonchus asper	spiny sow thistle			X	A	
Sonchus oleraceus	store-front sow thistle			X	A	
Sonchus uliginosus	common sow thistle				X	A
Sorghastrum nutans	Indian grass		X	X	X	
Sparganium americanum	American bur reed				X	
Sparganium chlorocarpum	simple bur reed			X		
Sparganium eurycarpum	common bur reed			X		
Spartina pectinata	prairie cord grass	X	X	X		
Sphenopholis intermedia	slender wedge grass				X	
Spiraea alba	meadowsweet		X	X	X	
Spirea tomemtosa rosea	steeple bush			X	X	
Spiranthus cernua	nodding lady's tresses			X	X	
Sporobolus asper	rough dropseed			X	A	
Sporobolus cryptandrus	sand dropseed				X	

Talinum rugospermum	fame flower					X	E
Taraxacum officinale	common dendelion			X		X	A
Tephrosia virginiana	goat's rue			X		X	
Teucrium canadense	germander					X	
Thalictrum dioicum	early meadow rue				X		
Tilia americana	American basswood					X	
Tofieldia glutinosa	false asphodel				X		R
Tradescantia ohiensis	spider wort			X		X	
Tragopogon major	sand goat's beard		X		X	A	
Tragopogon pratensis	common goat's beard					X	A
Trifolium hybridum	alsike clover					X	A
Trifolium pratense	red clover					X	A
Trifolium repens	white clover					X	A
Triglochin maritima	common bog arrow grass			X	X		
Triplasis purpurea	sand grass					X	
Triticum aestivm	wheat					X	A
Typha angustifolia	narrow-leaved cattail			X	X	X	A
Typha latifolia	broad-leaved cattail			X	X	X	
Ulmus pumila	Siberian elm					X	A
Utricularia cornuta	horned bladderwort					X	T
Utricularia gibba	humped bladderwort					X	
Utricularia minor	small bladderwort					X	
Utricularia vulgaris	great bladderwort				X		
Vaccinium angustifolium laevifolium	early low blueberry					X	
Vaccinium vacillans	late low blueberry					X	
Vallisneria americana	eel grass				X		
Verbascum thapsus	common mullein		X	X	X	A	
Verbena hastata	blue vervain			X	X	X	
Verbena stricta	hoary vervain			X		X	
Vernonia missurica	Missouri ironweed					X	
Veronicastrum virginicum	Culver's root				X		
Viburnum acerifolium	maple-leaved arrow-wood				X		
Viburnum lentago	nannyberry					X	
Viburnum prunifolium	black haw		X				
Viburnum rafinesquianum	downy arrow-wood					X	
Vicia americana	American vetch		X				
Viola conspersa	dog violet				X		
Viola fimbriatula	sand violet					X	
Viola lanceolata	lance-leaved violet			X			
Viola papilionacea	common blue violet				X		
Viola pedata lineariloba	bird's foot violet				X		
Viola pensylvanica	smooth yellow violet					X	
Viola sagittata	arrow-leaved violet			X			
Viola tricolor	pansy violet		X	A			
Vitis aestivalis	summer grape		X				
Vitis labrusca	fox grape			X			
Vitis riparia	riverbank grape	X	X				
Vitis vulpina			X				
Xanthium strumarium	cocklebur		X	A			
Zizania aquatica	wild rice		X				
Zizania aurea	meadow parsnip	X					

AQUATIC MACROINVERTEBRATES

Laurel L. Last and Richard L. Whitman

Lake Michigan Ecological Research Station
Biological Resources Division
U.S. Geological Survey

caddisfly larvae (*Hydropsyche*). Near the margin, a sandy bottom will include occasional snails (*Goniobasis*, *Pleurocera*, and *Campeloma*), midge larvae, occasional burrowing mayfly larvae, a number of mussels (*Unio gibbosus* and *Quadrula rubiginosa* being most characteristic), and occasionally a long-legged dragonfly larva (*Macromia taeniolata*). A silty bottom will include the mussels *Quadrula undulata* and *Lampsilis siliquoidea*, the burrowing mayfly larva *Hexagenia*, midge larvae, segmented worms, sphaeriid clams, and the mud leech *Haemopis grandis*.

The zone of vegetation formation includes the water scorpion *Ranatra fusca*, the creeping water bug

identified to genus, except the leeches (Hirudinea), earthworms (Oligochaeta), and water mites (Acari), which were not identified any further.

Indiana Department of Environmental Management study (IDEM unpublished data)

Benthic macroinvertebrate data from the Grand Calumet River and Indiana Harbor Canal were collected by the Indiana Department of Environmental Management from 1979 to 1988. The 1986-1988 data have been summarized by Bright (1988). Macroinvertebrates were collected with one to three multi-plate Hester-Dendy artificial substrate samplers per site. The samplers were generally left in the water from six to eight weeks. Two samplers were collected from the East Branch of the Grand Calumet at Virginia Street (IDEM1) in 1987. Three samplers were collected from the East Branch at Bridge Street (IDEM2) in 1986, and two in both 1987 and 1988. Two samplers were collected from the East Branch at Cline Avenue (IDEM3) in both 1986 and 1988, and one in 1987. Three samplers were collected from the East Branch at Kennedy Avenue (IDEM4) in 1986, and two in 1988. Three samplers were collected from the West Branch of the Grand Calumet at Indianapolis Boulevard (IDEM5) in 1986. One sampler was collected from the mouth of the West Branch (IDEM6) in 1987, and two in 1988. Three samplers were collected from Lake George Canal at the railroad bridge (IDEM7) in 1986, and one in 1987. Three samplers were collected from Indiana Harbor Canal at Dickey Rd (IDEM8) in 1979, 1980, and 1981, and two in 1986, 1987, and 1988. Most organisms other than the aquatic earthworms were identified to genus or species if possible; however, the midges (Chironomidae) were usually not identified beyond family from every Hester-Dendy collected during a single sampling.

Illinois Natural History Survey study (Risatti and Ross 1989)

Benthic macroinvertebrate data from Indiana Harbor and Indiana Harbor Canal were collected by the Illinois Natural History Survey on 3 and 4 May 1988. Two petite Ponar grab samples were collected from each site, one for organism enumeration and identification, and one for determination of wet and dry biomass standing crop for the dominant taxa. Each grab sample was washed in a #30 mesh screen bucket and preserved in 10% buffered formalin. One site (INHS1) was located in Lake George Canal, just west of Indianapolis Boulevard. Three sites were located in Indiana Harbor Canal: INHS2, just downstream of Columbus Drive; INHS3, just downstream of Route 912; and INHS4, just downstream of Dickey Road. Two sites were located in Indiana Harbor: INHS5, near the south end of the harbor; and INHS6, near the north end of the harbor. Sexually mature tubificid earthworms were identified to species level; other organisms were identified to family or genus level.

TAMS Consultants, Inc., study (Mierzwa et al. *1991)*

Benthic macroinvertebrate data from the Grand Calumet River were collected by TAMS Consultants, Inc.(Mierzwa et al. 1991), in 1990 and 1991. Three petite Ponar grab samples were collected from each site for each sampling period. Each sample was washed in a #30 mesh screen bucket and then preserved in 10% buffered formalin. The East Branch was sampled at

Route 12 (TAMS1) July 1990, November 1990, and May 1991, and at Cline Avenue (TAMS2) November 1990 and May 1991; the West Branch was sampled at Burnham Avenue (TAMS3) November 1990, May 1991, and July 1991. Most organisms were identified to genus or species level, except the aquatic earthworms and midges, which were identified to family and subfamily, respectively. Numerical data were not published.

U.S. Fish and Wildlife Service study (Sobiech et al. *1994)*

Benthic macroinvertebrate data from the East Branch of the Grand Calumet River were collected by the U.S. Fish and Wildlife Service in 1994. Five multi-plate artificial substrate samplers consisting of five 5.8-cm diameter circular discs were placed at each site on 19 May 1994 and were retrieved on 29 June 1994. Qualitative (non-numerical) sampling was also performed. Only qualitative sampling was possible at the site upstream of Tennessee Street (FWS1). The East Branch was also sampled: FWS2, just downstream of Broadway Avenue; FWS3, just upstream of Interchange 13 entrance/exit ramps of I-90; FWS4, downstream of Bridge Street; and FWS5, at the Wabash railroad trestle. All organisms were identified to family level, except the aquatic earthworms (Oligochaeta), leeches (Hirudinea), scuds (Amphipoda), and crayfish (Decapoda), which were not identified any further.

MACROINVERTEBRATES BY REACH

Lagoons reach

This section includes the Grand Calumet River Lagoons east of USX, and the sites NPS1 and NPS2 (Figure 1). The Lagoons reach is different from the other reaches in that it is connected to the rest of the river by partially-constricted culverts. The most common taxa in this reach were the snail genus *Ferrissia* and the scud genus *Hyalella* in the East Lagoon and *Hyalella*, the midge genus *Glyptotendipes*, and the damselfly genus *Ischnura* in the West Lagoon (Hardy

Physidae, aquatic earthworms, and leeches (Attachment 1). In addition, the snail genus *Ferrissia* was common at IDEM2. At IDEM2, the only site at which midges were identified below family level, the most common midge was *Cricotopus bicinctus*. Invertebrate Community Index (ICI) metric evaluation of the FWS study indicated that the invertebrate community of this reach was severely impaired (Sobiech *et al.* 1994). All sites received a total ICI score of 2 or lower and were classified as having very poor or poor invertebrate biotic integrity. The unbalanced trophic structure of the community, which was dominated by gathering collectors, also indicated degraded environmental conditions.

Gary Sanitary District reach

This reach includes the East Branch of the Grand Calumet River from the USX property to Cline Avenue. Sampling sites from east to west include: TAMS1, TAMS2, and IDEM3 (Figure 1). The most common taxa in this reach were aquatic earthworms, leeches, and the snail family Physidae, plus midges at IDEM3 (Attachment 1). The midges *Cricotopus* (unidentified) and *Cricotopus bicinctus* (possiPICI)

macroinvertebrate habitat, as indicated by its consistently low species richness, diversity, and equitability, and high (low quality) MBI.

Culverts reach

This reach includes the West Branch of the Grand Calumet River from Columbia Avenue to the Illinois/Indiana state line. None of the sampling sites are found within this reach. However, due to industrial and municipal impacts on sediment and water quality in the area, it is unlikely that the macroinvertebrate habitat is better than that in the Roxanna Marsh reach.

Hammond Sanitary District reach

This reach includes the West Branch of the Grand Calumet River from the Hammond/East Chicago boundary to Columbia Avenue. None of the sampling sites are found within this reach. However, due to industrial and municipal impacts on sediment and water quality in the area, it is unlikely that the macroinvertebrate habitat is better than that in the Roxanna Marsh reach.

Roxanna Marsh reach

This section includes the West Branch of the Grand Calumet River from Indianapolis Boulevard to the Hammond/East Chicago boundary, and sampling site IDEM5 (Figure 1). The most common taxa found were the snail genus *Physa*, the midge *Parachironomus abortivus*, and the midge *Chironomus decorus* (Attachment 1). All of these are quite pollution-tolerant (Attachment 2), suggesting very poor habitat. A hydrologic divide occurs at the western end of this reach, so that some of the water flows west to join the Little Calumet River and some flows east to Lake Michigan via the Indiana Harbor Canal.

Lake George reach

This section is the western portion of Lake George Canal, ending approximately 1100 ft west of Indianapolis Boulevard, and it includes the site IDEM7 (Figure 1). The most common taxa found were bryozoans (Bryozoa), aquatic earthworms, the snail genus *Physa*, and *Hydra* (Attachment 1). Most of these are highly pollution-tolerant (Attachment 2), indicating that the macroinvertebrate habitat is probably poor.

Federal Dredging Project reach

The Federal Dredging Project reach, although not specifically addressed in this study, can provide additional information on Grand Calumet macroinvertebrate populations. This section includes the Indiana Harbor and Canal from Lake Michigan to Columbus Drive and the eastern portion of Lake George Canal to approximately 1100 ft west of Indianapolis Boulevard. Sampling sites are: INHS1 in Lake George Canal; INHS2, INHS3, IDEM8, and INHS4 in Indiana Harbor Canal; and INHS5 and INHS6 in Indiana Harbor (Figure 1). The most common taxa found were aquatic earthworms, identified as the family Tubificidae in the INHS sites, the snail family Hydrobiidae and hydras (Hydridae) at INHS5, and bryozoans (Bryozoa) at IDEM8 (Attachment 1).

Many taxa, such as the midges, mayflies (Ephemeroptera), caddisflies (Trichoptera), damselflies, and snails (other than Hydrobiidae), were identified at IDEM8 but not at the INHS sites. These differences likely stem from the different sampling methods used (Hester-Dendy versus petite Ponar) and number of samples collected, rather than real differences in the communities. For example, both IDEM8 and INHS4 are in Indiana Harbor Canal near Dickey Road, yet at least 19 taxa were found at IDEM8 and only four at INHS4. The enumeration and identification data in the INHS study, however, were obtained from a single petite Ponar grab, whereas the IDEM data at this site are drawn from 15 Hester-Dendy artificial substrate collections over nine years. The IDEM data also show a general increase in richness and diversity from 1979-1988, with a peak in 1986 probably caused by historic highs in Lake Michigan water levels (Bright 1988). Although the invertebrate community in this reach is probably degraded, as indicated by the dominance of aquatic earthworms at every site, it may not be as poor as the INHS data suggest.

SPECIES LIST AND DESCRIPTIONS

Sponges (phylum Porifera)

Members of the phylum Porifera, the sponges, are the simplest of the multicellular animals. Of the more than 5000 species of sponges, the vast majority are marine, and only about 27 species occur in the fresh waters of the United States and Canada (Frost 1991). Freshwater sponges are common in unpolluted ponds, lakes, streams, and rivers, and they may be found attached to almost any stable submerged object (Pennak 1989). Most freshwater sponges obtain nourishment through both filtration and large numbers of symbiotic algae living within their cells; they use these sources in different proportions depending on environmental conditions

(Frost 1991). They may be fed upon by fish, crayfish, and possibly snails, but their primary

Roundworms were found in the Gary Sanitary District and DuPont Reaches at TAMS2 (Figure 1 and Attachment 1). They were not identified any further. Freshwater roundworms have been rated quite tolerant of certain natural phenomena and indicative of organically enriched or polluted waters (Attachment 2). However, they are not uniformly sensitive to pollutants (Poinar, Jr. 1991), and a recent study did not rate them as being either pollution-tolerant or intolerant (Attachment 2). It is likely that roundworms also inhabit other areas, but their small size (most < 1 cm in length) would make it easy for them to slip through the #30 sieves or to be unobserved during the separation of the macroinvertebrates from the rest of the samples.

Bryozoans (phylum Bryozoa/Ectoprocta and Endoprocta)

Bryozoans have been called "moss animals" because colonies of some common species can resemble a mat of moss. Approximately 4000 marine species of bryozoans have been described, and only about 50 freshwater species, including about 22 in the United States (Pennak 1989). Freshwater bryozoans attach to submerged surfaces, and will grow on aquatic vegetation and almost any solid, biologically inactive material (Wood 1991). They occur in both still and running water, but are generally restricted to relatively warm water. Freshwater bryozoans use ciliated tentacles for capturing suspended food particles, which may include diatoms, desmids, dinoflagellates, green and blue-green algae, bacteria, rotifers, small roundworms, protozoa, and even microcrustaceans, along with detritus and inorganic materials. However, little is known about bryozoan nutrition other than what they ingest, and items that are ingested are not

1991). The vast majority of aquatic earthworms feed by ingesting bottom sediments and digesting the organic component (Pennak 1989). *Chaetogaster*, one of the few carnivorous freshwater aquatic earthworms, feeds on entomostraca, insect larvae, protozoans, and other aquatic earthworms. *Aeolosoma* feeds on microorganisms and fine particulate debris, using cilia to sweep them toward its mouth. The burrowing activity of aquatic earthworms can contribute greatly to sediment mixing and solute transport across the mud-water interface (Brinkhurst and Gelder 1991).

Aquatic earthworms were found at every site except NPS2 in the Lagoons reach (Figure 1 and Attachment 1). They were identified simply as aquatic earthworms at the NPS, FWS, and IDEM sites. The family Lumbricidae was collected at TAMS1 and 2 (Gary Sanitary District reach), and the family Lumbriculidae was collected at TAMS1, 2, and 3 (Far West reach). The family Tubificidae was identified at the INHS sites in the Federal Dredging Project reach as: *Limnodrilus*, *Limnodrilus cervix*, *Limnodrilus hoffmeisteri*, *Potamothrix vejdovskyi*, and *Quistadrilus multisetosus*. In addition, immature worms without capilliform chaetae were found at all six sites, and those with capilliform chaetae were found at INHS2.

Aquatic earthworms are generally considered quite tolerant of pollution and/or enrichment (Attachment 2). Species composition can be a valuable indicator, however, with a series of species groups inhabiting progressively more polluted stretches of rivers or more eutrophic lakes (Brinkhurst and Gelder 1991). In the Great Lakes, there are basically three species associations of Tubificidae: *T. tubifex, Peloscolex multisetosus*, and several *Limnodrilus* species (dominated by *L. hoffmeisteri* and *T. tubifex*), characteristic of organically polluted bays and harbors; *Aulodrilus*, *Potamothrix*, *Limnodrilus* and *Peloscolex ferox*, characteristic of eutrophic conditions; and *L. hoffmeisteri*, *T. tubifex*, and many species not often found in the other areas, characteristic of "clean" waters (Brinkhurst and Cook 1974).

Leeches (class Hirudenea)

The leeches are predominantly freshwater organisms, with about 60 freshwater species known in the United States (Pennak 1989). They commonly inhabit ponds, marshes, lakes, and slow streams, particularly in the northern half of the country, and the same species may occur in a variety of environments. Leeches are represented in North America by four families: Glossiphoniidae, which either prey upon macroinvertebrates or temporarily parasitize fish, turtles, amphibians, or water birds; Piscicolidae, which parasitize fish and crustaceans; Erpobdellidae, which primarily prey upon macroinvertebrates and zooplankton; and Hirudinidae, which either prey upon macroinvertebrates or suck the blood of amphibians and mammals (Davies 1991). Predators of leeches include fish, birds, garter snakes, newts, salamanders, insects, snails, and scuds.

Leeches were identified at all sites except FWS1 in the USX reach; TAMS3 in the Far West reach; and INHS1, 3, 4, 5, and 6 in the Federal Dredging Project reach (Figure 1 and Attachment 1). No leeches were identified below class at the NPS or FWS sites. The family Erpobdellidae was identified to: family, *Dina microstoma* (uncertain), *Dina parva*, *Erpobdella punctata*, *Mooreobdella*, *Mooreobdella fervida*, and *Mooreobdella microstoma*. The family Glossiphoniidae was represented by *Helobdella*, *Helobdella stagnalis*, and *Placobdella*.

Cocoons were identified at IDEM4, 5, and 7. It is possible that the lack of leeches found at the

of small particles including bacteria, algae, ciliates, and small rotifers. In addition, chydorids gather food by scraping, and *Leptodora*, *Polyphemus*, and *Bythotrephes* prey upon protozoans, rotifers, small midge larvae, and small crustaceans. Water fleas are an important food source for fish; in addition, they are eaten by hydras and immature and mature insects (Pennak 1989).

Water fleas were found at only one site--IDEM7 in the Lake George reach (Figure 1 and Attachment 1). Two organisms were identified in 1986, and they were not identified to a lower taxonomic level. Water fleas have been rated quite tolerant of certain natural phenomena (Attachment 2). Pollution tolerances vary among species, but most (19 out of 22) of the water flea species identified in the Delaware and Neches Estuaries and the Flint River in New England were rated characteristic of natural conditions by Patrick and Palavage (1994). It is likely that water fleas also inhabit other areas, but their small size would make it easy for them to slip through the #30 sieves or to be unobserved during the separation of the macroinvertebrates from the samples.

Class Malacostraca

The water mites include five distantly related groups in the subclass Acari (mites and ticks), with the most numerous and diverse group, by far, being the Hydrachnida (or Hydrachnellae, Hydracarina, or Hydrachnidia) (Smith and Cook 1991). Water mites inhabit many different habitat types, including springs, rapidly flowing areas of streams and rivers, sand and gravel deposits, cool pools, lakes, permanent ponds, wetlands, and temporary pools. Larval water mites parasitize many kinds of aquatic insects, and pre-adult and mature water mites prey on insect and fish eggs, insect larvae, ostracods, cladocerans, and copepods, scavenge on dead organisms, or parasitize bivalves. Water mites are eaten by a wide variety of aquatic invertebrates, particularly coelenterates and insects (Pennak 1989), and occasionally form a significant part of fish and turtle diets (Smith and Cook 1991).

Water mites were found at NPS2 in the Lagoons reach (Figure 1 and Attachment 1). They were not identified any further than Acari (formerly Acarina). Water mites are excellent indicators of environmental quality; their diversity is greatly reduced in chemically polluted or physically disturbed habitats (Smith and Cook 1991). Water mites have been rated as quite tolerant of certain natural phenomena and indicative of clean unpolluted waters (Attachment 2).

Subphylum Uniramia of phylum Arthropoda

Insects (class Insecta)

Mayflies (order Ephemeroptera)

The mayflies all have aquatic larvae which may be found in streams, rivers, lakes, and temporary or permanent ponds and marshes (Hilsenhoff 1991). Almost all mayfly larvae are herbivores or detritivores, but a few species prey on other invertebrates; the adults do not feed. Often, mayfly larvae are an important food source for fish in streams.

Mayflies were found at NPS1 and 2 in the Lagoons reach and IDEM8 in the Federal Dredging Project reach (Figure 1 and Attachment 1). The family Baetidae (small minnow mayflies) was represented by *Baetis* at NPS2 and IDEM8. The family Caenidae (small squaregills) was represented by *Caenis* at NPS1 and 2. The family Heptageniidae (flatheaded mayflies) was represented by *Stenonema* (*pulchellum* group) at IDEM8. Mayflies as a group are very important biological indicators for water quality because many species are very susceptible to water pollution or occur in predictable habitat types (McCafferty 1983).

Baetidae is a widespread and abundant family and can be found in a variety of streams, permanent and temporary ponds, and lake shallows (Hilsenhoff 1991). The family has been used as an indicator of clean unpolluted waters and possible slight organic pollution (Attachment 2). Habitats of *Baetis* include “torrential mountain streams, warm, meandering rivers, and still waters of northern Canada” (McCafferty 1983). *Baetis* has been rated moderately tolerant of certain natural phenomena, slightly pollution tolerant, and characteristic of natural conditions (Attachment 2).

Caenidae larvae are widespread and common in a wide variety of habitats, including streams, spring seeps, marshes, swamps, ponds, and lakes (Hilsenhoff 1991). They are generally more tolerant of low levels of dissolved oxygen than mayflies in other families. Caenidae is considered an indicator of clean, unpolluted waters, and *Caenis* has been rated moderately

(Attachment 2). *Hydropsyche* is considered quite tolerant of certain natural phenomena, somewhat pollution-tolerant, and characteristic of natural conditions (Attachment 2). *H. orris* has been rated slightly pollution-tolerant, indicative of some organic pollution, and characteristic of natural conditions (Attachment 2). *H. simulans* is considered somewhat pollution tolerant and indicative of significant organic pollution (Attachment 2).

Hydroptilidae larvae may be found in a wide variety of habitats and feed on algae and other plant material (Hilsenhoff 1991). They have been rated quite tolerant of certain natural phenomena and indicative of possible slight organic pollution (Attachment 2). *Neotrichia* has been considered quite tolerant of certain natural phenomena, slightly pollution-tolerant, indicative of no apparent organic pollution, indicative of clean, unpolluted streams, and characteristic of natural conditions (Attachment 2). *Orthotrichia* has been considered pollution intolerant and indicative of clean to slightly enriched streams (Attachment 2).

Leptoceridae larvae occur in a variety of permanent aquatic habitats (Hilsenhoff 1991). Most are omnivore-detritivores, but *Oecetis* species are predators, and some *Ceraclea* feed on freshwater sponges (McCafferty 1983). They have been rated somewhat tolerant of certain natural phenomena, indicative of clean, unpolluted streams, and indicative of possible slight organic pollution (Attachment 2). *Leptocerus* has been rated slightly pollution-tolerant (Attachment 2). *Nectopsyche* has been considered slightly pollution-tolerant, indicative of no apparent organic pollution, and characteristic of natural conditions (Attachment 2). *Oecetis* has been considered somewhat tolerant of certain natural phenomena, somewhat pollution-tolerant, indicative of very significant organic pollution, and characteristic of natural conditions (Attachment 2).

Most Polycentropodidae larvae inhabit streams, but they also occur in a variety of other habitats (Hilsenhoff 1991). Most species are predators, but a few are herbivores. Polycentropodidae has been considered moderately tolerant of certain natural phenomena, indicative of likely substantial organic pollution, and indicative of clean, unpolluted streams (Attachment 2). *Cyrnellus* has been rated somewhat pollution tolerant (Attachment 2). *C. fraternus* has been considered indicative of very significant organic pollution (Attachment 2). *Neureclipsis* has been considered slightly pollution tolerant, indicative of significant organic pollution, and characteristic of natural conditions (Attachment 2).

Water bugs (order Hemiptera, suborder Heteroptera)

Heteropterans, the true bugs, are primarily terrestrial, but about 8.5% of the species are aquatic, with 217 species living in North American fresh waters and 107 species living on them (Hilsenhoff 1991). Water bugs are remarkably diverse and occupy many different habitat types, including saline ponds, mountain lakes, hot springs, and large rivers (Polhemus 1984). Most species are predators; however, many genera of the water boatmen (Corixidae) are primarily collectors, feeding on detritus. They can be important predators of mosquito larvae and adults; however, some species bite people or eat small fish in hatcheries, thereby becoming a nuisance. Most water bugs seem to be resistant to predation, possibly due to their characteristic scent glands. However, the water boatmen are preyed upon by fish, used as food for pet fish and turtles, and relished by people in Mexico, and the giant water bugs (Belostomatidae) are considered a delicacy in Asia.

Water bugs were identified at NPS1 and 2 in the Lagoons reach and FWS5 in the USX reach (Figure 1 and Attachment 1). The family Belostomatidae was identified to *Lethocerus* at NPS2. The family Corixidae was identified to family at FWS5 and *Sigara* at NPS1. The family Pleidae (pygmy backswimmers) was identified to *Plea* at NPS2. Water bugs are more tolerant of environmental extremes than most insects, except the water beetles and flies (Roback 1974).

Giant water bugs inhabit permanent standing water habitats (*Belostoma* and *Lethocerus*), especially weedy ponds, lake margins, marshes, or streams (*Abedus*), among aquatic plants or under rocks in riffles (Hilsenhoff 1991). They are powerful predators and will capture and kill fish, frogs, tadpoles, and other insects. Giant water bugs have been rated moderately tolerant of certain natural phenomena and indicative of clean, unpolluted streams; *Lethocerus* has been rated moderately tolerant of certain natural phenomena (Attachment 2).

Water boatmen are good water quality indicators in standing waters (Polhemus 1984). They are found in most permanent aquatic habitats, and frequently temporary ones as well (Hilsenhoff 1991). They feed primarily on detritus, algae, protozoans, and other very small animals; a few species will capture and eat larger insects such as mosquito larvae. *Sigara* are notable as herbivores (McCafferty 1983). Water boatmen have been rated quite tolerant of certain natural phenomena and indicative of clean, unpolluted streams, and *Sigara* has been rated quite tolerant of certain natural phenomena (Attachment 2).

Pygmy backswimmers inhabit vegetation, primarily in permanent ponds but also in lake shallows, stream backwaters, and swamps (Hilsenhoff 1991). They feed on small invertebrates. Pygmy backswimmers have been considered to be indicative of clean unpolluted streams (Attachment 2).

Water beetles (order Coleoptera)

Only about 3% of Coleoptera species have an aquatic life stage, but since it is the largest insect order, this amounts to more than 1100 aquatic species in North America (Hilsenhoff 1991). Three suborders have aquatic representatives: Adephaga, represented by five families in which all species have aquatic larvae and adults; Myxophaga, represented by a single species in North America in which both larvae and adults are aquatic; and Polyphaga, represented by species in several families in which larvae, adults, or both are aquatic. Water beetles inhabit a broad range of aquatic environments, and most are usually found on surfaces rather than in open water (White *et al*. 1984). Feeding habits of water beetles are extremely variable.

Water beetles were found at NPS2 in the Lagoons reach, IDEM1 in the USX reach, and IDEM6 in the East Chicago Sanitary District reach (Figure 1 and Attachment 1). The family Dytiscidae (predaceous diving beetles) was identified as *Dytiscus* at IDEM1 and *Laccophilus* at NPS2. The family Gyrinidae (whirligig beetles) was identified as *Dineutus* at NPS2 and *Gyrinus* (uncertain) at IDEM6. The family Haliplidae (crawling water beetles) was identified as *Haliplus* at NPS2. Water beetles are more tolerant of environmNorth Am(0 Tw 37iNnt 1rS37iNnling water beetles) was

They have been considered moderately tolerant of certain natural phenomena and indicative of clean, unpolluted streams (Attachment 2). *Dytiscus* has been considered moderately tolerant of certain natural phenomena, and *Laccophilus* has been considered moderately tolerant of certain natural phenomena and pollution-tolerant (Attachment 2).

Whirligig beetles are widespread and often abundant (Hilsenhoff 1991). Larvae are predators, feeding mostly on other invertebrates; adults are scavengers on dead animals or predators of small invertebrates. They have been considered quite tolerant of certain natural phenomena and indicative of clean, unpolluted streams (Attachment 2). *Dineutus* has been considered slightly pollution-tolerant, and *Gyrinus* has been rated quite tolerant of certain natural phenomena and pollution-tolerant (Attachment 2).

Crawling water beetles are often abundant in shallow vegetation-choked habitats (Hilsenhoff 1991). Both larvae and adults are herbivores, feeding on algae or aquatic plants. They (and *Haliplus*) have been rated somewhat tolerant of certain natural phenomena (Attachment 2).

Flies and midges (order Diptera)

Biting midges (suborder Nematocera, family Ceratopogonidae)

Although Diptera (flies and midges) is mostly a terrestrial order, it is the dominant order of insects in the aquatic environment (Hilsenhoff 1991). The order is divided into two suborders: Nematocera, which dominates the aquatic fauna, and Brachycera. Biting midge larvae live in a variety of aquatic habitats, including tree holes, marshes, swamps, ponds, lakes, and streams. Most larvae are carnivores, and others are herbivores or detritivores. Adults of some aquatic species feed on people; most others (including *Palpomyia*) feed on small insects (McCafferty 1983). Biting midges were found at NPS1 and 2 in the Lagoons reach, FWS5 in the USX reach, and IDEM6 in the East Chicago Sanitary District reach (Figure 1 and Attachment 1). They were identified to family at FWS5 and IDEM6, and to *Palpomyia* at NPS1 and 2. Biting midges have been rated quite tolerant of certain natural phenomena, somewhat pollution-tolerant, indicative of clean unpolluted streams, and indicative of likely substantial organic pollution (Attachment 2). *Palpomyia* has been considered moderately pollution-tolerant, indicative of likely substantial organic pollution, and pollution-tolerant (Attachment 2).

Phantom midges (suborder Nematocera, family Chaoboridae)

Phantom midges, so calltiv331 Tf1Tw 18MCID 10 >> Tuted(Paly Chaoboridae))]TJETp cyoboridairers are

Procladius has been rated moderately pollution-tolerant, indicative of severe organic pollution, and characteristic of natural conditions (Attachment 2). *Thienemannimyia* group has been rated moderately pollution-tolerant and characteristic of natural conditions (Attachment 2).

Mosquitoes (suborder Nematocera, family Culicidae)

Mosquito larvae inhabit a variety of shallow, standing waters, including tree holes, artificial containers, catch basins, pitcher plants, swamps, shallow temporary or permanent ponds and marshes, and heavily vegetated margins of lakes and streams (Hilsenhoff 1991). Larvae feed primarily on small aquatic animals, algae, and detritus; larvae of the genus *Toxorhynchites* and some *Psorophora* are predaceous, often feeding on other species of mosquito larvae (Newson 1984). The adult females of a great majority of species feed on the blood of various cold- and warm-blooded animals, including humans. The adult males feed on plant juices and nectar. Mosquitoes are vectors of many human and animal diseases.

Mosquito larvae were identified at IDEM6 in the East Chicago Sanitary District reach (Figure 1 and Attachment 1). They were not identified to a lower taxonomic level. Mosquito larvae have been rated quite tolerant of certain natural phenomena, moderately pollution-tolerant, and indicative of organically polluted streams (Attachment 2).

Crane flies (suborder Nematocera, family Tipulidae)

The vast majority of species in Tipulidae, the largest family of Diptera, are not aquatic, but several genera have species with aquatic larvae (Hilsenhoff 1991). Crane fly larvae are found in nearly every kind of aquatic environment, including slow-flowing rivers, torrential mountain streams, margins of ponds and lakes, marshes, and tree holes (Byers 1984). Many larvae eat organic detritus--such as decaying leaves, plant fragments, and associated microorganisms--that accumulates on the bottoms of ponds or in backwaters of streams. Larvae of several species in the subfamily Limoniinae are active predators. Larval and adult crane flies are an important food source for other invertebrates, birds, mammals, fish, amphibians, and reptiles.

Crane fly larvae were common at IDEM6 in the East Chicago Sanitary District reach (Figure 1 and Attachment 1). They were not identified to a lower taxonomic level. Crane fly larvae have been rated moderately tolerant of certain natural phenomena, slightly pollution-tolerant, indicative of unlikely organic pollution, and indicative of clean unpolluted waters (Attachment 2).

Dance flies (suborder Brachycera, family Empididae)

Most species of dance flies are terrestrial or semiaquatic, but some have aquatic larvae (Hilsenhoff 1991). Most aquatic dance fly larvae and pupae live on the rocky bottoms of rapid streams, but some live on the margins of ponds and streams; larvae are predaceous (McCafferty 1983). One dance fly larva was found at FWS5 in the USX reach (Figure 1 and Attachment 1). It was not identified below family. Dance fly larvae have been rated quite tolerant of certain natural phenomena, moderately pollution-tolerant, indicative of likely substantial organic pollution, and indicative of clean, unpolluted streams (Attachment 2).

Soldier flies (suborder Brachycera, family Stratiomyidae)

Although most soldier fly larvae are terrestrial or semiaquatic, there are many species that live in shallow, vegetated standing waters (Hilsenhoff 1991). Aquatic larvae feed on detritus and algae; adults feed mostly on flowers. Soldier fly larvae were identified at FWS4 in the USX

depletion. Improving sediment quality will be necessary to restore macroinvertebrate habitat in the Grand Calumet River, and one way to achieve this is by cleaning up the contaminated sediments.

Sediment clean-up options include removal (dredging), capping with clean materials, and in-place treatment. Although it is theoretically feasible, not enough is known about in-place treatment (*e.g.*, fixation/solidification or biodegradation) to consider it seriously at this point (USACE 1994a). Dredging--the excavation of bottom sediments from a waterway--may be performed with a variety of equipment (USACE 1994b). The two basic types of dredges are mechanical dredges, which remove sediments using a large bucket or shovel, and hydraulic dredges, which remove and transport the sediments in a water slurry. The particular method selected depends on reach-specific factors that will not be discussed here.

The positive impacts of dredging on the aquatic macroinvertebrates, provided that it would remove the total depth of contaminated sediments, would be the removal of the contaminants (and thus their direct and indirect negative impacts) from the system, and the uncovering of uncontaminated sediments for recolonization. However, both of these benefits would be greatly reduced without sediment source controls (see below). Removal of only part of the contaminated sediments would be of little benefit unless water depth was maintained with capping.

The negative impacts of dredging would include the removal of existing benthic macroinvertebrates and rooted vegetation, potentially severe adverse habitat impacts due to changes in channel morphology, and temporary, localized degradation of water and habitat quality. Although little could be done about the first impact, the macroinvertebrate communities that would be removed are degraded, and the newly exposed sediments would be recolonized and revegetated over time. The other two negative impacts could be minimized by taking certain steps during and after dredging. The placement of weirs up- and downstream of the dredging operation would help maintain water flow and surface levels and could increase localization of turbidity during dredging. Care should be taken not to dig the channel too deep or the banks too steep in order to encourage the re-establishment of rooted vegetation and minimize flow pattern changes and riverbank sloughing. In reaches where the contaminated sediment is quite deep or the River is already too channelized, the bottom could be filled in with clean sediment.

Capping is the covering of contaminated sediment by clean materials (USACE 1994a). The contamination remains in the waterway, but its availability to the water column and aquatic life is reduced. The cap must isolate sediment contamination from the overlying water, prevent penetration by benthic or burrowing organisms, and be resistant to scour. Cap design depends on various factors, including the hydraulic system, sediment characteristics, and types of contamination. One concern with leaving the sediment in place is that groundwater may still interact with the contaminants.

The positive impacts of capping on the aquatic macroinvertebrates, provided that it would isolate and stabilize the contaminated sediments, would be the removal of the possibility of direct and indirect negative impacts from the contaminants and the availability of new, uncontaminated sediments for recolonization. However, both of these benefits would be greatly reduced without sediment source controls.

The negative impacts of capping would include the covering of existing benthic macroinvertebrates and rooted vegetation and possible adverse habitat impacts due to water

depth reduction in shallow areas. Although little could be done about the first impact, the macroinvertebrate communities that would be buried are degraded, and the newly exposed sediments would be recolonized and revegetated over time. In shallow areas, partial dredging prior to capping could compensate for water depth loss.

Sediment source controls

Source controls, which reduce the quantity and contamination level of sediments entering the River, will be very important in improving sediment quality and macroinvertebrate habitat in the Grand Calumet River and Indiana Harbor Canal. If done properly, they could only impact the macroinvertebrate communities positively. Reductions in the quantity of sediment entering the River would improve habitat by decreasing siltation and turbidity, both of which can be very detrimental to some macroinvertebrates. Without reductions in contamination levels of sediments entering the river, sediment clean-up would only provide a temporary solution, since uncontaminated sediment would simply be covered and replaced by more contaminated sediment (USACE 1994a).

There are three major sources of sediments to the Grand Calumet River and Indiana Harbor Canal: municipal and industrial point discharges, combined sewer overflows (CSOs), and urban runoff. Point sources include three municipal wastewater treatment plants and over 40 outfalls for discharges from industries and manufacturers. Over 90% of the system's dry-weather flow originates as treated municipal and industrial wastewater (McCown *et al*. 1976). Point discharges are regulated under the Clean Water Act (NPDES permit program); effects of this regulation can be seen in the 56% reduction of suspended solids loadings from point sources

Transport controls reduce the resuspension and transport of sediments that have already been deposited on the river bottom. Reductions in sediment resuspension and transport would improve macroinvertebrate habitat by reducing turbidity, erosion, and the exposure of the organisms and the water column to sediment contaminants (USACE 1994a). Sediment impacts on water quality and aquatic organisms are directly related to the sediment surface area exposed, and when sediments are in suspension, surface area is greatly increased. Sediment resuspension

constraints. Restoration would require the cooperation of federal and state agencies, local governments, industries, municipal wastewater treatment plants, and the public, and many compromises would be necessary. Industrial and residential development have severely altered the landscape and the River itself, yet there remains a great potential for improving the health of the River and the whole ecosystem. Let us take the challenge, and future generations will surely thank us for it.

LITERATURE CITED

Brannon, J.M., D. Gunnison, D. Averett, J.L. Martin, R.L. Chen, and R.F. Athow, Jr. 1989. Analysis of impacts of bottom sediments from Grand Calumet River and Indiana Harbor Canal on water quality. Misc. Paper D-89-1. U.S. Army Corps of Engineers, Waterways Experiment Station, Vicksburg, MS. Not read; cited in USACE 1994a.

Bright, G.R. 1988. Recent water quality in the Grand Calumet River as measured by benthic invertebrates. Proceedings of the Indiana Academy of Science **98**:229-233.

Brinkhurst, R.O., and D.G. Cook. 1974. Aquatic earthworms (Annelida: Oligochaeta). Pages 143-156 *in* C.W. Hart, Jr., and S.L.H. Fuller, editors. Pollution ecology of freshwater invertebrates. Academic Press, New York, New York, USA.

Brinkhurst, R.O., and S.R. Gelder. 1991. Annelida: Oligochaeta and Branchiobdellida. Pages 401-435 *in* J.H. Thorp and A.P. Covich, editors. Ecology and classification of North American freshwater invertebrates. Academic Press, New York, New York, USA.

Brown, K.M. 1991. Mollusca: Gastropoda. Pages 285-314 *in* J.H. Thorp and A.P. Covich, editors. Ecology and classification of North American freshwater invertebrates. Academic Press, New York, New York, USA.

Bushnell, J.H. 1974. Bryozoans (Ectoprocta). Pages 157-194 *in* C.W. Hart, Jr., and S.L.H. Fuller, editors. Pollution ecology of freshwater invertebrates. Academic Press, New York, New York, USA.

Byers, G.W. 1984. Tipulidae. Pages 491-514 *in* R.W. Merritt and K.W. Cummins, editors. An

IDEM. 1991. The Remedial Action Plan for the Indiana Harbor Canal, the Grand Calumet River and the nearshore Lake Michigan, Stage One. Indiana Department of Environmental Management, Indianapolis, Indiana, USA.

Illinois EPA. 1985. Illinois Environmental Protection Agency macroinvertebrate tolerance list.

Kenk, R. 1974. Flatworms (Platyhelminthes: Tricladida). Pages 67-80 *in* C.W. Hart, Jr., and S.L.H. Fuller, editors. Pollution ecology of freshwater invertebrates. Academic Press, New York, New York, USA.

Ketcham, M.L., V.P. Kunchakarra, and C.T. Jafvert. 1992. Urban targeting of nonpoint source pollution in the Grand Calumet River watershed. Report to USDA Soil Conservation Service, Lake County Soil and Water Conservation District, Crown Point, Indiana, USA.

Kolasa, J. 1991. Flatworms: Turbellaria and Nemertea. Pages 145-171 *in* J.H. Thorp and A.P. Covich, editors. Ecology and classification of North American freshwater invertebrates. Academic Press, New York, New York, USA.

McCafferty, W.P. 1983. Aquatic entomology: the fishermen's and ecologist's illustrated guide to insects and their relatives. Jones and Bartlett Publishers, Inc., Boston, Massachussetts, USA. (Orig. pub. 1981 by Science Books International, Inc.)

McCown, D.L., W. Harrison, and W. Orvosh. 1976. Transport and dispersion of oil-refinery water in the coastal waters of southwest Lake Michigan (experimental design--sinking-plume condition). Argonne National Laboratory, Argonne, Illinois, USA. Not read; cited in USACE 1994a.

Patrick, R., and D.M. Palavage. 1994. The value of species as indicators of water quality. Proceedings of the Academy of Natural Sciences of Philadelphia **145**:55-92.

Pennak, R.W. 1989. Fresh-water invertebrates of the United States: Protozoa to Mollusca, 3rd edition. John Wiley & Sons, Inc., New York, New York, USA.

Poinar, G.O., Jr. 1991. Nematoda and Nematomorpha. Pages 249-283 *in* J.H. Thorp and A.P. Covich, editors. Ecology and classification of North American freshwater invertebrates. Academic Press, New York, New York, USA.

Polhemus, J.T. 1984. Aquatic and semiaquatic Hemiptera. Pages 231-260 *in* R.W. Merritt and K.W. Cummins, editors. An introduction to the aquatic insects of North America, 2nd edition. Kendall / Hunt Publishing Co., Dubuque, Iowa, USA.

Risatti, J.B., and P. E. Ross, editors. 1989. Chemical, biological and toxicological study of sediments from Indiana Harbor, Canal and adjacent Lake Michigan. Final Rep. to USACE, Chicago District, Chicago, Illinois. 83 p.

Roback, S.S. 1974. Insects (Arthropoda: Insecta). Pages 313-376 *in* C.W. Hart, Jr., and S.L.H. Fuller, editors. Pollution ecology of freshwater invertebrates. Academic Press, New York, New York, USA.

Sawyer, R.T. 1974. Leeches (Annelida: Hirudinea). Pages 81-142 *in* C.W. Hart, Jr., and S.L.H. Fuller, editors. Pollution ecology of freshwater invertebrates. Academic Press, New York, New York, USA.

Shelford, V.E. 1977. Animal communities in temperate America: as illustrated in the Chicago region. Arno Press, New York, New York, USA. (Orig. pub. 1937 by the University of Chicago Press, Chicago, IL.)

Slobodkin, L.B., and P.E. Bossert. 1991. The freshwater Cnidaria--or coelenterates. Pages 125-143 *in* J.H. Thorp and A.P. Covich, editors. Ecology and classification of North American freshwater invertebrates. Academic Press, New York, New York, USA.

Smith, I.M., and D.R. Cook. 1991. Water mites. Pages 523-592 *in* J.H. Thorp and A.P. Covich, editors. Ecology and classification of North American freshwater invertebrates. Academic Press, New York, New York, USA.

Sobiech, S.A., T.P. Simon, and D.W. Sparks. 1994. Pre-remedial biological and water quality assessment of the East Branch Grand Calumet River Gary, Indiana, June 1994. USFWS, Bloomington, Indiana, USA 112 p.

INTRODUCTION

The Grand Calumet River - Indiana Harbor Canal fish community structure and function have undergone drastic changes since the turn of the century. Changes in stream depth, water quality, and land use eliminated indigenous species during the middle of the century. Dredging and channelizing altered the stream from a riverine wetland to a narrow channel and significantly increased the flow velocity. During the early 1970's, few if any species of fish were documented in either the East or West Branches of the Grand Calumet River. Young-of-the-year of lake dwelling transient species were present only seasonally in the Indiana Harbor Canal. Subchronic toxicity was documented at a few of the major dischargers to the system. Documented improvements in water quality and the fish community in this decade suggest that the possibility of restoring the Grand Calumet River is not only a dream but a distinct possibility. The river

The change in the fish community of the Grand Calumet River is a direct result of the flow changes during the early- to mid 1800's. Prior to these changes, the Grand Calumet was comprised of palustrine wetlands, a series of shallow pools connected by narrow flowages between pools. These pools and flowages enabled movement of species between Lake Michigan and the important wetland nursery and spawning habitat provided by the slow flowing River. Shelford (1937) compared the physical status of the Grand Calumet River to other Lake Michigan tributaries such as the lower St. Joseph River, lower Galien, and Dead River. Although Shelford did not specifically describe the condition of the Grand Calumet River, it can be supposed that the river possessed characteristics similar to the other streams he mentioned.

These rivers characteristically possess bottoms of either fine sand or fine organic materials with little or no rock. The rivers originate in wetland habitats characterized by emergent rooted macrophytes. Differentiation of pools and riffles are not common. Instead, deep pools and narrow, shallow connections (termed flowages) enable passage of water between pools. The flow of water is mostly determined by the depth of Lake Michigan and reflects annual climatologic and hydrologic conditions. The deeper portions of these streams support rooted aquatic macrophytes such as various pondweeds (*Potemogeton* sp.), and the water lilies, *Nuphar* and *Nelumbo*, thrive. Bullrushes (*Scirpus* sp.) and cattails (*Typha*) line the shoreline and gradually sloping banks along back bays and side channel margins.

Meek and Hildebrand (1910) evaluated the distribution of fish species within a fifty mile radius from Chicago and indicated that as many as 22 species occurred in the Grand Calumet River drainage (Table 1).

Shelford documented 12 species of fish during 1909 including species such as blacknose shiner (*Notropis heterolepis*) lake chubsucker, northern pike, redhorse, and tadpole madtom (*Noturus gyrinus*) (Table 2). The Dead River has a much smaller drainage area than the Grand Calumet River, but unlike the Grand Calumet River, it has not been extensively modified. The Dead River is a designated nature preserve contained within the Illinois Beach State Park near Zion, Illinois that is one of the few remaining areas of dune and swale topography in Illinois. When we sampled the Dead River in 1996 we captured 14 species, six of which were present in the 1909 sample. Of the 20 total species collected from the Dead River during the two events, less than 30% were found in both the 1909 and 1996 collections.

TABLE 1. Fishes collected by Meek and Hildebrand (1910) from the Grand Calumet River drainage and deposited at the Field Museum of Natural History.

Common Name	Species Name
bowfin	*Amia calva*
carp	*Cyprinus carpio*
golden shiner	*Notemigonus crysoleucas*
emerald shiner	*Notropis atherinoides*
spottail shiner	*N. hudsonius*
bluntnose minnow	*Pimephales notatus*

white sucker	*Catostomus commersoni*
channel catfish	*Ictalurus punctatus*
black bullhead	*Ameiurus melas*
yellow bullhead	*A. natalis*
brown bullhead	*A. nebulosus*
tadpole madtom	*Noturus gyrinus*
northern pike	*Esox lucius*
grass pickerel	*E. americanus*
central mudminnow	*Umbra limi*
green sunfish	*Lepomis cyanellus*
pumpkinseed	*L. gibbosus*
bluegill	*L. macrochirus*
black crappie	*Pomoxis nigromaculatus*
yellow perch	*Perca flavescens*
logperch	*Percina caprodes*
freshwater drum	*Aplodinotus grunniens*

Until the 1960's, the Grand Calumet River suffered from chemical and physical degradation caused by municipal and industrial pollution. Untreated sewage and waste from packing plants and heavy industry eliminated most of the natural aquatic communities. Surveys of the river during the early 1960's found only 20-108 oligochaetes/m^2 in the Indiana Harbor Canal mouth. By the early 1970's, the abundance of oligochaetes had increased to 2,400-500,000/m^2. The increase in biomass was considered sufficient to support native fish populations.

TABLE 2. Species of Fish collected from the Dead River in 1909 (Shelford 1937) and 1996 (Moy and Simon, unpublished data).

		Year	
Common Name	Species Name	1909	1996
gizzard shad			

pumpkinseed	*Lepomis gibbosus*	--	X
warmouth	*Lepomis gulosus*	--	X
bluegill	*Lepomis macrochirus*	X	X
white crappie	*Pomoxis annularis*	X	X
black crappie	*Pomoxis nigromaculatus*	--	X
smallmouth bass	*Micropterus dolomieui*	--	X
largemouth bass	*Micropterus salmoides*	X	X

the screening procedure but could not be resampled due to a plant shutdown in 1986. East Chicago Sanitary District effluent did not elicit a chronic response for hatching or survival during definitive testing. However, statistically

proportion of tolerant species, and an increase in the use of the River by migrating transient species, including young-of-the-year yellow perch and adult rainbow trout.

Simon et al. (1988) evaluated seven locations in the East Branch between 1985-1988. The number of species increased at stations sampled multiple times. Generally, the dominant species comprising the East Branch fish community were carp, goldfish, and golden shiner. Collections of pumpkinseed, largemouth bass, black crappie, bluegill, central mudminnow, and rainbow smelt have increased between 1985 and 1988 (Table 4). The lack of adult representatives of most species indicate the area served as nursery habitat and as an intermittent food and habitat resource.

Simon (1991) developed biological integrity expectations for northwestern Indiana as part of an evaluation of the Central Corn Belt Plain. An index of biotic integrity was developed and calibrated for the Lake Michigan subdivision of the Lake Michigan drainage. Three stations in the East Branch were sampled as a part of this investigation. Although species lists were not reported by station, individual IBI assessments were calculated for each station. The results did not differ significantly from the 1988 results (Simon et al. 1988). The 1990 survey was a high water year so many species, including pumpkinseed and yellow perch which were not collected during the 1988 drought, were found again as they began to return to the area. An increase in abundance of the rudd (*Scardinius erythrophthalmus*), a European exotic, was noted in the East and West
Branches of the Grand Calumet River. Exotic species have become the majority of the fish species collected (Table 3).

TABLE 3. Summary of fish species collected from the Indiana Harbor Canal segment ([1] Polls and Dennison 1984; [2] Simon et al. 1988; [3] Risatti and Ross 1989).

			List of sites										
								1983					
	1986	1987	1988	1988	1994	1995	1996						
Species			A[1]	B[1]	C[1]	D[1]	E[1]	F[1]	G[2]	G[2]	G[2]	D, E, F[3]	B[3]
Gizzard shad, *Dorosoma cepedianum*			41	8	705	47	3	72		65		25	14

Black crappie,

TABLE 4. Summary of fish collections completed in the East Branch of the Grand Calumet River ([1] CDM/Limnetics; [2] IDEM unpublished data; [3] Simon et al. 1988; [4] Simon 1990; and [5] Sobiech et al. 1994).

	List of sites									
Species	1976 A[1]	1980 B,C,D[2]	1982 B,C,D[2]	1984 B,C,D[2]	1985 B,C,E[3]	1986 B,C,D[2]	1987 B,C,D[2]	1986 B,C,D[3]	1987 B-G[3]	1988 B,C,D[3]
Gizzard shad, *Dorosoma cepedianum*							96		157	
Alewife, *Alosa psuedoharengus*	2						1		2	8
Central mudminnow, *Umbra limi*	1			3				16	1	
Chinook salmon, *Oncorhynchus tschawytscha*								1		
Rainbow trout, *Oncorhynchus mykiss*						11				
Carp, *Cyprinus carpio*	5	18	5	31	111	19	59	165	36	1
Goldfish, *Carassius auratus*					34		113	157	140	161
Carp x Goldfish hybrid										
Rudd, *Scardinius erythrophthalmus*										
Spotfin shiner, *Cyprinella spiloptera*		1					1			12
Golden shiner, *Notemigonus crysoleucas*			1	10	124	64	25	222	140	239
Golden shiner x Rudd hybrid										
Emerald shiner, *Notropis atherinoides*							2			14
Bluntnose minnow, *Pimephales notatus*						92	21	1	5	3
Fathead minnow, *Pimephales promelas*					1		1			1
Black bullhead, *Ameiurus melas*				1			1			
White sucker, *Catostomus commersoni*										
Rainbow smelt, *Osmerus mordax*								1	1	
Pumpkinseed, *Lepomis gibbosus*						48	14	59	41	1
Bluegill, *Lepomis macrochirus*							2	6	2	2
Green sunfish, *Lepomis cyanellus*					3			6		
Largemouth bass, *Micropterus salmoides*								2		
Black crappie, *Pomoxis nigromaculatus*								1		
Yellow perch, *Perca flavescens*		1				3	142		4	
unidentified darter, Percidae species				1						
Total Number	2	8	19	16	197	281	471	470	691	519

In 1988, a fish kill occurred in the upper five miles of the East Branch of the Grand Calumet River due to an upper River oil spill which caused degr

TABLE 5. Summary of fish collected from sites in the West Branch Grand Calumet River ([1] Simon et al. 1988; [2] Simon 1990; [3] Simon, unpublished data).

	List of sites		
	1985	1986	1987

In 1990, most of northwestern Indiana was affected by significant flooding. The surface elevation of Lake Michigan and the depth of the West Branch increased. Samples from the Indianapolis Boulevard area taken that year produced the greatest number of fish from the West Branch representing seven species (Simon 1991). The first Indiana collection of the eurasian rudd (*Scardinius erythrophthalmus*) was included with this catch. The water quality of the discharge canal at East Chicago enabled several very tolerant species such as green sunfish and bluntnose minnow to inhabit the West Branch. The fish community downstream of Indianapolis Boulevard was numerically dominated by carp and goldfish, however, the proportion of exotic species was significantly reduced as the number of native species such as bluntnose minnow and green sunfish increased (Table 5). Fish at this site had high proportions (4.5%) of deformities, eroded fins, lesions, and tumors (DELT anomalies); fish were collected which had eroded fins, fungus, and lesions.

Simon (unpublished data) sampled the West Branch in the vicinity of Indianapolis Boulevard and Roxanna Marsh again during 1994. One site extended from the east side of Indianapolis Boulevard to an area approximately 50 m upstream from the junction of the East and West Branches. The East Chicago Sanitary District discharge canal was included in this sampling site. The second location included the area from Indianapolis Boulevard west along the edge of Roxanna Marsh to the I5).till0, tTable e 5 -1.1ethe east side of

Central mudminnow, *Umbra limi*	x			1			
Grass pickerel, *Esox americanus*	x	x		6		x	
Northern pike, *Esox lucius*	x						
Carp, *Cyprinus carpio*	x	18	2	17			
Goldfish, *Carassius auratus*	x			1			
Carp x Goldfish hybrid							
Common shiner, *Luxilus cornutus*	x						
Golden shiner, *Notemigonus crysoleucas*	x	x			2		
Emerald shiner, *Notropis atherinoides*		x					
Blackchin shiner, *Notropis heterolepis*	x				51		
Bluntnose minnow, *Pimephales notatus*							
Lake chubsucker, *Erimyzon sucetta*	x	x	1	13			
Black redhorse, *Moxostoma duquesnei*	x						
Black bullhead, *Ameiurus melas*	x		1				
Yellow bullhead, *Ameiurus natalis*	x	x		1			
Brown bullhead, *Ameiurus nebulosus*	x						
Tadpole madtom, *Noturus gyrinus*	x			1			
Green sunfish, *Lepomis cyanellus*	x	x	2	12	15		
Pumpkinseed, *Lepomis gibbosus*	x	x	1	70	32	x	x
Warmouth, *Lepomis gulosus*	x	x		20	2		
Bluegill, *Lepomis macrochirus*	x	x	5	113	27		
Largemouth bass, *Micropterus salmoides*	x	x	2	26	94		
White crappie, *Pomoxis annularis*	x						
Black crappie, *Pomoxis nigromaculatus*		x	1	1			
Yellow perch, *Perca flavescens*	x	x		9	25		
Iowa darter, *Etheostoma exile*		8	4				
Total Number	--	--	30	297	281	--	--
Total Species	20	14	7	14	11	1	1
% Tolerant Species	30	30	70	5.4	30.6	0	0
% Exotic Species	0	14.3	60	0.1	6.4	0	0
Number of Collections	--	--	1	3	3	2	2

A = Dunal ponds identified by Shelford speculated to be Grand Calumet Lagoons; B = Middle Lagoon; C= West Lagoon; D = East Pond; E = West Pond.

Other assessment categories were used at the stations that had no fish. At these stations the IBI scores differed by more than 10 IBI points. An Ohio study found the largest departures in IBI scores were due to large scale disturbance (Yoder and Rankin 1995). In the Grand Calumet River, the wider fluctuations in IBI points also reflected highly disturbed conditions. The largest difference observed in IBI points within any single year occurred at the site to the east of Indianapolis Boulevard during 1987, when a difference of 6 IBI points was observed (Table 7).

FISH CONSUMPTION ADVISORIES

The Indiana State Department of Health (ISDH 1997) states that fish from the Grand Calumet River should not be consumed, primarily due to contamination by PCB's and mercury. Table 8 lists the fish species and sizes included in the consumption advisory. The advisory is less restrictive for the Marquette Park Lagoon, recommending only that largemouth bass more than 12 inches long be avoided. There is a state-wide advisory on carp consumption: 1 meal/ month of carp 15-20" long, 1 meal/2months for carp 20-25" in length and no carp over 25" should be consumed.

Numerous species in the nearshore of Lake Michigan also are restricted for consumption. Factors which would limit complete removal of consumption advisories and eventual recovery and delisting of the Grand Calumet River are: 1) Lake Michigan species which enter the River during migration include wide ranging salmonids, 2) resident species in the Grand Calumet River would require several generations to reduce body burdens, and 3) stocking of non-indigenous species and migration of species from other more contaminated areas into the river will require more time to see decline in contaminant body burdens. Recovery and restoration objectives for fish communities in the Grand Calumet River should initially strive toward the goal of reaching the same levels as Lake Michigan.

Fish bioaccumulation, body burdens, and contaminant levels

Steffek (1989) evaluated three abandoned hazardous waste dump sites located in the Grand Calumet River watershed. Steffek collected a variety of organisms representing various trophic levels and feeding regimes. Earthworms, turtles, crayfish, fish, and small mammals were tested for body burdens. Compounds found at elevated levels in whole body tissue samples included acetone, 2-butanone, benzene, trichloroethane, toluene, ethylbenzene, total xylenes, aluminum, copper, chromium, lead, manganese, and silver. Values for lead were above the national levels obtained from the national biomonitoring program (Lowe et al. 1985). Taxa collected as environmental indicators showed various levels of effect. Earthworms were absent from many of the sites but showed mixed results as a result of illegal dumping and mixed contamination. Fish, mudpuppy, and crayfish provided important bioaccumulation information for inorganic and volatile compounds, while turtles did not show significant results.

Sparks and Hudak (1996) reviewed available information on environmental impacts associated with the dredging of the Indiana Harbor Canal. High sediment levels of polycyclic aromatic hydrocarbons (PAHs), phenols, cyanide, metals, and ammonia have been detected in the

TABLE 7. Index of Biotic Integrity values for the Grand Calumet Watershed from 1985 to 1990 (Simon et al. 1988; Simon 1991). Sites indicated below are: 1) Western Lagoon; 2) East Branch Grand Calumet River, Broadway; 3) East Branch, Bridge Street;
East Branch, Grant Street; 5) East Branch, Cline Avenue; 6) East Branch, Kennedy Ave; 7) East Branch, Dupont de Nemours
outfall; 8) Junction of East and West Branch Grand Calumet River; 9) West Branch Grand Calumet River, Indianapolis Blvd.;
10) West Branch, I-90 bridge; 11) West Branch, Columbia Ave., 12) Indiana Harbor Canal, Dickey Road.

	Index of Biotic Integrity Score											
Year	1	2	3	4	5	6	7	8	9	10	11	12
1985												
October			24			24		24			24	0
1986												
June	32		26		24	24			22			24
October			28		30	28			20			26
1987												
April		30	32	24	22		22	22	24	24		
April			24		24	26			22			28
November			32		30	30			0			34
1988												
May			26		22	24			0			
July			28		32	26			0			24
1990												
July			24		20	32			21			16
Avg IBI	32	27	27.5	24	25.3	27.1	23	22	13.6	24	0	25.3
Minimum	32	24	24	24	20	24	22	22	0	24	0	16
Maximum	32	30	32	24	32	32	24	22	24	24	0	34
SD	--	3	1.0	--	1.33	1.14	1.0	--	3.0	0	--	3.0
N	1	2	8	1	9	7	2	1	8	2	1	6

Rainbow trout	<22	PCB's	1
	22+	PCB's	0.5
Walleye	17-26	PCB's	1
	26+	PCB's	0.5
Whitefish	<23	PCB's	1
	23+	PCB's	0.5
White sucker	15-23	PCB's, Mercury	1
	23+	PCB's	0.5

Flow velocity in the Grand Calumet River varies with effluent discharge volumes. Channel depth through the length of the study area is highly variable ranging from 8 to 10 feet in the Grand Calumet Lagoons to one foot or less in portions of the West Branch near the state line. Removal or isolation of contaminated sediment and consequent reduction of sediment resuspension may improve water quality after sediment remediation is complete. However, deepening of the channel will create a more pool-like habitat resulting in slowed current, and possibly reduced dissolved oxygen concentrations. The additional water depth may prohibit the reestablishment of rooted vegetation which provides substrate for invertebrates, cover for fish and is a source of oxygen.

The dredging and sediment clean-up process may temporarily degrade water quality and reduce aqueous habitat quality. Dredging suspends sediment in the water column which can increase turbidity and the dissolved contaminant concentration and reduce dissolved oxygen concentrations. These impacts tend to be temporary, and ambient levels return shortly after dredging ceases. Discharge from the dewatering site must be treated to reduce the concentration of contaminants in order to meet state water quality standards. Care must be taken to assure that the discharge does not adversely affect instream dissolved oxygen levels and that concentrations of toxic chemicals such as ammonia are within acceptable limits. Weirs surrounding the dredge site can help to contain water quality impacts within the immediate area of dredging.

Long-term habitat impacts would result as dredging converts shallow, littoral habitat to more pelagic habitat with steep banks and a narrow littoral zone. The vegetation in the littoral zone forms important foraging and nursery habitat for young fish. Water depths and clarity after dredging must be such that light can penetrate to the bottom of the channel to assure reestablishment of this important aquatic habitat component.

Suggestions for aquatic habitat improvements

The U.S.EPA in 1985 identified factors which reduced the quality of biological habitat in portions of the Grand Calumet River system. Slow flowing or stagnant water in the West Branch warms causing decreased dissolved oxygen concentrations. Particulates settle over detrital matter and suffocate benthic organisms. The U.S.EPA (1985) stated that these conditions are typical of many Great Lakes stream mouth environments, but they are exaggerated in the West Branch of the Grand Calumet River.

Flow reversals due to fluctuations in Lake Michigan water levels have less impact on the East Branch and main stem due to the high lakeward flow rate sustained by industrial effluent, particularly at USX. Without a high volume of industrial discharges, the majority of the Grand Calumet River system would probably be much less suitable for fish and aquatic life (assuming adequate control of pollutants in the effluent) (U.S.EPA 1985).

An additional obstacle to restoration is the intensity of urban and industrial development within the watershed. Most precipitation falling to the basin is captured on non-porous surfaces, i.e., roof drains, parking lots and roadways, and is then diverted to the River via storm drains. This reduces the opportunity for soil filtration of dissolved and suspended pollutants, exaggerates peak flows and depresses low flows, reduces dilution of groundwater contaminants, and slows dispersion from origin to point of discharge along the river.

Establishment of aeration stations, either using injected air or elevated sidestream aerators such as those present on the Cal-Sag Channel and Chicago River in Illinois, could help to improve the aquatic habitat by maintaining adequate dissolved oxygen concentrations, as well as by assisting in the breakdown of organic debris.

After the contaminated sediment is removed, care should be taken to assure that the remaining channel profile is designed to allow establishment of a littoral zone. The rooted vegetation that grows in the littoral zone provides forage and cover for invertebrates and young fish, as well as ambush sites for predatory species. Creation of a littoral zone may involve replacement of dredged material with clean material in some portions of the river. Replacement with gravel or other substrate will provide substrate heterogeneity suitable for lithophilic spawners and nest builders such as sucker, darters, and sunfish, and it may facilitate the reestablishment and maintenance of these populations. Woody debris such as brush piles, stumps, or logs placed along the channel border can form current breaks, nesting cavities, hard surfaces for attachment of adhesive eggs, and habitat for invertebrates. Half logs can be placed in the center of the channel to provide cover and shaded areas for species that previously would have used aquatic vegetation. Wetland, bog, and dune areas adjacent to the river may serve as spawning and rearing areas for many important fish species (U.S.EPA 1985). Creation of artificial wetlands and shallow side channels would form more of this important limiting habitat.

Innovative dredging techniques could be used to create a side channel through the heavily choked *Typha, Phragmites*, and purple loosestrife stands adjacent to the River. These side-channels could be staggered and would be parallel to the channel to emulate a braided wetland channel. The entrance to these areas could be kept open by installing deflector logs to scour and divert flow from the main channel. In addition, many depth profiles could be established by refilling with clean sand. Also, side-channel created wetlands could be created by dredging perpendicular to the channel and then filling with clean sand to create a shelf zone. This area could be planted with native emergent wetland plants which would serve as fish nursery habitat, and as foraging and resting areas for wading birds, reptiles, and mammals. A rigorous effort to control exotic, invasive plant species would have to be implemented to control reinvasion of exotic plant species after initial efforts are implemented.

After sediment clean up, water quality of industrial and municipal discharges will need to continue to meet NPDES permit requirements. Further treatment and design improvements are needed to reduce impacts attributable to thermal pollution, nutrient enrichment, combined sewer overflows, and other non-point source episodes. A single episode of oxygen depletion could be sufficient to eliminate an entire year class of young fish. Without continued emphasis on meeting designated uses and NPDES permit limits, little or no improvement of the fish community can be expected.

LITERATURE CITED

Baumann, P.C. 1989. PAH, metabolites, and neoplasia in feral fish populations. Pp. 269-289 *in* U. Varanasi, editor. Metabolism of polycyclic aromatic hydrocarbons in the aquatic environment. CRC Press, Boca Raton, Florida, USA. 341 p.

of Engineers Chicago District. Metropolitan Sanitary District of Greater Chicago, Chicago, Illinois, USA.

Risatti, J.B. and P. Ross. eds. 1989. Chemical, biological and toxicological study of sediments from Indiana Harbor Canal and adjacent Lake Michigan. Final Report for U.S. Army Corps of Engineers, Chicago District, Chicago, Illinois, USA.

Robertson, B. 1986. Evaluation of 1984 supplementa

Protection: Science, Politics, and Common Sense. Vancouver, British Columbia, Canada. p. 281 (Abstract).

U.S. Environmental Protection Agency. 1984. EPA master plan for improving water quality in the Grand Calumet River/Indiana Harbor Canal. U.S. Environmental Protection Agency, Water Division, Chicago, Illinois, USA. EPA 905/9-84/003A.

U.S. Fish and Wildlife Service. 1987. Polycyclic aromatic hydrocarbon hazards to fish, wildlife, and invertebrates: a synoptic review. U.S. Department of the Interior, Biological Report 85(1.11). Contaminant Hazard Reviews Report 11.

Yoder, C.O. and E.T. Rankin. 1995. Biological criteria program development and implementation in Ohio. Pp. 109-144 *in* W.S. Davis and T.P. Simon, editors. Biological Assessment and Criteria: Tools for Water Resource Planning and Decision Making. Lewis Press: Boca Raton, Florida, USA.

AMPHIBIANS AND REPTILES

Kenneth S. Mierzwa

TAMS Consultants
180 N. Stetson, Suite 3200
Prudential Plaza
Chicago, Illinois 60601

Spencer A. Cortwright

Biology Department, Marram Hall
Indiana University Northwest
3400 Broadway
Gary, Indiana 46408-1197

David Beamer

INTRODUCTION

The ridge and swale area surrounding the Grand Calumet River is home to one of the more diverse assemblages of amphibians and reptiles in northwestern Indiana. Here, as a result of a series of recent geological and climatic events, species more typical of areas to the north, south, east, and west come together and occur in close proximity. Surprisingly, despite the long and intensive industrial history of the region, several relatively pristine natural areas have survived along with most of their salamander, frog, turtle, lizard, and snake species. The survival of these natural areas and their associated faunas provides unique opportunities for preservation and restoration.

We have looked beyond the immediate riparian area, in part because much more is known about the contiguous ridge and swale sites. Since amphibians and reptiles are less mobile than many other vertebrates, and less likely to colonize remote or isolated habitats, it is important to inventory existing centers of biodiversity and factor that information into management decisions.

We have largely relied on information available from recent inventories and older museum specimens and publications, interpreted in the context of our own field experience in the area.

Nomenclature follows Collins (1990), except for the recent elevation of *Bufo fowleri* to specific status (Sullivan et al. 1996).

PRESETTLEMENT CONDITIONS

Origins of the herpetofaunal assemblages

Presumably, amphibians and reptiles entered what is now the Grand Calumet River area shortly after the end of the Wisconsinan glaciation. Remains of the cold-tolerant turtles *Chelydra serpentina* and *Chrysemys picta* have been found in association with mastodon bones in southeastern Michigan and east central Indiana (Holman and Andrews 1994), at sites dated to 13,000-11,000 years before present (ybp). Other species which today have northern distributions must have also been present in the boreal forest which covered northwestern Indiana at that time. However, the present day site of the Grand Calumet River was still covered by the waters of post-glacial Lake Chicago. Subsequent fluctuation of lake levels (Chrzastowski and Thompson 1992; Chrzastowski et al. 1991) and climate change (Ahearn and Kapp 1984; Ebbers 1984) profoundly influenced the sequence and location of later colonization events.

The diverse herpetofauna includes northern elements such as blue-spotted salamanders, *Ambystoma laterale,* and Blanding's turtles, *Emydoidea blandingii;* and eastern species like eastern newts, *Notophthalmus viridescens,* and green frogs, *Rana clamitans*. These species may

Combining early published sources, specimens in area museum collections, and knowledge based on the best remaining natural areas and examples of particular habitat types, the following list of potential presettlement amphibians and reptiles has been compiled:

Mudpuppy, *Necturus maculosus*
Blue-spotted salamander, *Ambystoma laterale*
Tiger salamander, *Ambystoma tigrinum*
Eastern newt, *Notophthalmus viridescens*
Redback salamander, *Plethodon cinereus*
American toad, *Bufo americanus*
Fowler's toad, *Bufo fowleri*
Northern cricket frog, *Acris crepitans*
Spring peeper, *Pseudacris crucifer*
Western chorus frog, *Pseudacris triseriata*
Gray tree frog, *Hyla versicolor*
Bullfrog, *Rana catesbeiana*
Green frog, *Rana clamitans*
Northern leopard frog, *Rana pipiens*
Snapping turtle, *Chelydra serpentina*
Musk turtle, *Sternotherus odoratus*
Painted turtle, *Chrysemys picta*
Spotted turtle, *Clemmys guttata*
Blanding's turtle, *Emydoidea blandingii*
Map turtle, *Graptemys geographica*

While the Grand Calumet River and bordering riparian wetlands support a depauperate herpetofauna, several natural areas contiguous with the river are noted for their species richness. Some parcels of lesser quality also support amphibians or reptiles tolerant of habitat modification. These sites are described individually below, on a reach-by-reach basis, from east to west.

Grand Calumet Lagoons reach

The lagoons at the easternmost end of the Grand Calumet River are partially bordered on the north and south by Miller Woods, which are a part of Indiana Dunes National Lakeshore. Areas

Musk turtle, *Sternotherus odoratus*
Painted turtle, *Chrysemys picta*
Blanding's turtle, *Emydoidea blandingii*
Six-lined racerunner, *Cnemidophorus sexlineatus*
Eastern hognose snake, *Heterodon platirhinos*
Brown snake, *Storeria dekayi*
Western ribbon snake, *Thamnophis proximus*
Common garter snake, *Thamnophis sirtalis*

9.20k Storeria dekayi epitanc -0.00011 Tw 9.47501 0 T6.09502therus odoratus

The Clark and Pine complex of seven sites is located north of the Grand Calumet River at the eastern end of the Gary Sanitary District reach, and extends slightly into the adjacent USX reach. The seven natural areas include perhaps the highest concentration of rare and endangered plant and animal species remaining in the state of Indiana (Bowles 1989). Although only Clark and Pine East directly borders the Grand Calumet River, several of the areas are divided from each other only by railroad tracks or roads, and to some extent they probably still function as a unit. Several of the ponds studied by Shelford (1913) were within or near the Clark and Pine complex. Others who collected in the general area included Hubbs and Meek, and specimens are available from as early as 1902.

Clark and Pine East, sometimes referred to as the Bongi site, borders the Grand Calumet River just east of the U.S. Route 12 bridge and extends to the north. The approximately 102 hectare site was acquired by the Division of Nature Preserves, Indiana Department of Natural Resources, in 1993. The area was originally a classic example of ridge and swale habitat. Several of the ridges were mined for sand at various times in the recent past. The resulting low, wet, flat areas subsequently revegetated with mostly native species, including an unusual panne-like assemblage. Relatively undisturbed but overgrown sand savanna alternates with swales in the east-central part of the site. Management of Clark and Pine East is now underway, and consists largely of brush clearing and controlled burning. Amphibians and reptiles were inventoried during 1990 and 1991 by Mierzwa et al. (1991) as part of the Illinois-Indiana Regional Airport Study. The most intensive inventory effort, including quantitative drift fence sampling using pitfall and funnel traps (see Heyer et al. 1994 for a detailed discussion of drift fence techniques), was focused within a complex of dry-mesic sand savanna, mesic to wet sand prairie, marsh, and shrub swamp in the least disturbed portion of the site. Frequent searches were also carried out in other parts of the site.

The 17 hectare Clark and Pine Nature Preserve is located just to the northwest of Clark and Pine East. The two sites are separated by Clark Street. The nature preserve has been under state ownership for some time, and it has been intensively managed. It is generally wetter and more open than the preceding site, with extensive areas of pond, marsh, sand prairie, and open sand savanna. The sand savanna includes both black oak (*Quercus velutina*) and jack pine (*Pinus banksiana*); plant communities were described in detail by Bowles (1989). Clark and Pine Nature Preserve was sampled for amphibians and reptiles by Resetar (1988), using a combination of drift fence and random search techniques.

Other sites within the complex are privately owned. Lakeshore Railroad Prairie, a small site located just to the north of Clark and Pine East, has pond, marsh, panne, sand prairie, and sand savanna communities, the latter dominated by jack pine. Clarke Junction East and Clarke Junction West are located to the northwest of Clark and Pine Nature Preserve. These sites also include pond, marsh, panne, and sand prairie communities, but suffer from shrub encroachment. Limited information on the herpetofauna of these sites, based on only a few visits, is included in Mierzwa et al. (1991). Morgan and Burling (1979) visited several Clark and Pine complex sites and reported the presence of some amphibians and reptiles, but their lists include at least one apparent misidentification and therefore should be used with caution.

A list of amphibians and reptiles observed by Mierzwa et al. (1991) and Resetar (1988) within the Clark and Pine complex is included below. Eighteen species are known to occur within the complex at present, an unusually high number for a relatively small site. Unusual

species include the northern cricket frog, a once common species which has nearly disappeared from northern Indiana, and the state threatened spotted turtle.

Tiger salamander, *Ambystoma tigrinum*

of the Grand Calumet River watershed. The same survey recorded snapping turtles and painted turtles as common in open water areas. The marsh habitat is of relatively low quality, and it probably supports few species of amphibians and reptiles.

Hammond Sanitary District reach

Extant species

Blue-spotted salamander, *Ambystoma laterale*. This species occurs in wooded communities including relatively open savanna, woodland, mesic forest, and swamp forest. Breeding likely takes place in the swales. Blue-spotted salamanders are uncommon within the study area. Werth (1990) found one specimen at Miller Woods. Mierzwa et al. (1991) collected two in the western part of Ivanhoe Dune and Swale. Morgan and Burling (1979) reported one from the Clark and Pine area. The blue-spotted salamander is a northern species, and it is near the southern limit of its range here.

Tiger salamander, *Ambystoma tigrinum*. Although seldom seen because of burrowing habits, the tiger salamander is relatively widespread and common in the vicinity of the Grand Calumet River. The species breeds in marsh or shrub swamp within swales, and resides for the remainder of the year in dry-mesic sand savanna on the dune ridges. Individual salamanders are often found in surprisingly open locations; one Illinois population of tiger salamanders increased dramatically as savannas were opened up with brush clearing and prescribed burns (Mierzwa in press). Adult tiger salamanders usually do not travel far from breeding sites; recent studies in areas of sandy soil in New York have indicated that up to 80 percent of the population remains within 100 meters of the breeding wetland with a few individuals moving as far as 300 meters. Tiger salamanders spend most of their time in burrows created by shrews or other small mammals (Madison 1993), with limited surface activity on rainy nights.

Eastern newt, *Notophthalmus viridescens*. Newts are aquatic as adults, but they are capable of terrestrial movements if ponds dry. The adult phase may be preceded by a terrestrial eft stage lasting up to several years. Newts are common at Miller Woods (Werth 1990) and uncommon but present in the Clark and Pine area, where they were reported by Shelford (1913) and Mierzwa et al. (1991). In both areas newts inhabit pond and marsh communities in permanent and semi-permanent swales.

American toad, *Bufo americanus*. American toads are common in the western part of the study area, and they occur at Clark and Pine, DuPont, Ivanhoe Dune and Swale, near the Hammond Sanitary District treatment plant and at Burnham Prairie (Mierzwa et al. 1991). They have not been reported at Miller Woods. Elsewhere in northwestern Indiana, populations frequently occur on river floodplains. Water quality may constrain the toads' use of Grand Calumet riparian areas, but some have been observed in swales immediately adjacent to the river at DuPont.

Fowler's toad, *Bufo fowleri*. In northwestern Indiana this species, more characteristic of areas well to the south, is restricted to sandy soil. In the dunes area it typically breeds in blowout ponds and pannes in the immediate vicinity of Lake Michigan (Breden 1988). A population at Miller Woods, to the north of the Grand Calumet Lagoons, may be the westernmost one extant in the dunes area. Historically, the species occurred at Pine, where in 1917 a specimen was collected at the pond closest to Lake Michigan.

Northern cricket frog, *Acris crepitans*. Cricket frogs are associated with the margins of permanent ponds and streams, and they must have been common at one time along the banks of the Grand Calumet River. Sometime in the late 1970s or early 1980s, this once abundant frog disappeared from most localities in the northern part of its range. A population at the Clark and

Pine complex (Mierzwa et al. 1991) is the only one known to persist in Lake County. Cricket frogs are quite common at both Clark and Pine Nature Preserve and Clark and Pine East, and they are frequently seen along swale edges during the summer months.

Spring peeper, *Pseudacris crucifer*. Spring peepers are common at Miller Woods and in

in some wetlands near the river. Werth (1990) found musk turtles at Miller Woods, and a single specimen sited crossing a road in Hammond in 1996 turned in at Gibson Woods Nature Center.

Painted turtle, *Chrysemys picta*. There are numerous records of this species from the study area, as early as 1909 and as recently as 1996. It is one of the few species still occasionally found in the main channel. Permanent swales close to the river support sometimes dense populations, especially at Miller Woods, DuPont, and the Clark and Pine complex. Painted turtles inhabit both ponds and rivers, and they are easily observed because of their tendency to bask on logs or other objects.

Spotted turtle, *Clemmys guttata*. This small, attractive turtle is threatened in Indiana and endangered in Illinois. It is characteristic of shallow, sedge dominated wetlands and adjacent marsh borders. The Clark and Pine complex supports one of the larger spotted turtle populations in northwestern Indiana, but it has recently suffered from illegal collecting. Suitable habitat is present in swales very close to the Grand Calumet River at Clark and Pine East. This locality is nearly at the western distributional limits of the species.

Blanding's turtle, *Emydoidea blandingii*. This is a federal "species at risk" (formerly "Category 2"), and it is included on special concern or watch lists in most states in which they occur. Recent attention has focused on the long life span (> 60 years) and late age of maturity (14-20 years) of this species (Congdon et al. 1993). Because of these life-history characteristics, loss of reproductive females through highway mortality, over collecting, or other causes may have significant long-term effects on population viability. Blanding's turtles occur throughout the study area, with recent reports from Miller Woods, Clark and Pine East, Clark and Pine Nature Preserve, Lakeshore Railroad Prairie, Ivanhoe Dune and Swale, and DuPont. Populations appear to be at low density at most sites. Blanding's turtles are most common in swales which include pond and marsh communities, and particularly those with considerable submerged

study area are disjunct and near the northern limits of the distribution, and they may represent remnants from warmer or drier times.

Eastern hognose snake, *Heterodon platirhinos*. The only recent reports of hognose snakes have been from Miller Woods (Werth 1990). The species historically occurred at Pine (Richards 1987). This is an upland species found in sand prairie and sand savanna; it may occasionally forage for toads near wetland margins.

Milk snake, *Lampropeltis triangulum*. Milk snakes are uncommon in the study area.

Clark and Pine East (Lake Michigan Federation 1984). Extensive sampling there by several workers, including two with considerable massasauga experience, failed to confirm the record (Mierzwa et al. 1991). The habitat at Clark and Pine is also quite different from that typically utilized by the species. The report may be based on a sighting of the superficially similar northern water snake.

POTENTIAL

Several sites identified by the Indiana Natural Areas Inventory, including parts of the Clark and Pine complex, remain in private ownership and are at risk of degradation. Acquisition of these sites would help to preserve a number of rare species.

Acquisition of buffer areas

Acquisition of land contiguous to core natural areas, in conjunction with restoration activities discussed below, could greatly improve the long-term viability of sites which currently support high levels of biodiversity.

Restoration of core natural areas

The best quality sites are in need of woody vegetation management, including manual removal of exotic shrubs and the regular use of controlled fire. Enforcement activities to discourage illegal dumping and poaching would also be helpful.

Restoration of degraded sites

Because most of the landscape around the Grand Calumet River once consisted of alternating ridges and swales, some of the amphibians and reptiles in the area are capable of utilizing both upland and wetland habitat. Restoration efforts which address both habitat components are likely to support the greatest diversity of wildlife, especially if restoration areas are adjacent to the existing core natural areas. Early efforts can serve to create buffer areas for existing preserves. Over time these buffers could be extended to serve as connections between preserves. The challenges to this process will be many in this complex urban landscape, but opportunities should be pursued as they present themselves.

Wetlands with a direct river connection could contribute to water quality renovation and potentially provide habitat for some semi-aquatic amphibians and reptiles such as mudpuppies, bullfrogs, green frogs, snapping turtles, painted turtles, and northern water snakes. Turtles could benefit from opening the existing dense cattail marshes in riparian areas. Deepening of selected areas in conjunction with dredging operations could improve interspersion of marsh and open water habitats.

Many amphibians do not breed in riparian wetlands because of fish predation on eggs and larvae. Creation of a few palustrine emergent wetlands isolated from direct contact with the river could provide habitat for these species, including tiger salamanders, American toads, western chorus frogs, and northern leopard frogs. Because groundwater generally flows toward larger water bodies such as the Grand Calumet River (Watson et al. 1989), wetlands of excellent quality have been able to persist at DuPont, Clark and Pine East, Miller Woods, and other sites, only meters from the main channel. Restored or created wetlands separated from the channel by narrow areas of upland may have a reasonable chance of maintaining good water quality and supporting a variety of plants and animals.

In addition to traditional enhancement and restoration techniques such as woody vegetation control or wetland excavation, complex but innovative opportunities are possible near the Grand Calumet River. Some areas of degraded habitat are at present of minimal value to amphibians

LITERATURE CITED

Ahearn, P. J., and R. O. Kapp. 1984. Pollen analysis and vegetational history associated with archaeological sites in Berrien County, Michigan. Pages 113-123 *in* E. B. Garland, editor. Late

Breden, F. 1988. Natural history and ecology of Fowler's toad, *Bufo woodhousei fowleri* (Amphibia: Bufonidae) in the Indiana Dunes National Lakeshore. Fieldiana Zoology, New Series **49**:1-16.

Chrzastowski, M. J., and T. A. Thompson. 1992. Late Wisconsinan and Holocene coastal evolution of the southern shore of Lake Michigan. Quaternary coasts of the United States: Marine and lacustrine systems. SEPM Special Publication **48**:397-413.

Chrzastowski, M. J., F. A. Pranschke, and C. W. Shabica. 1991. Discovery and preliminary investigations of the remains of an early Holocene forest on the floor of southern Lake Michigan. Journal of Great Lakes Research **17**:543-552.

Collins, J. T. 1990. Standard common and current scientific names for North American amphibians and reptiles. Third edition. Society for the Study of Amphibians and Reptiles Herpetological Circular **19**. iii + 41p.

Congdon, J. D., A. E. Dunham, and R. C. van Loben Sels. 1993. Delayed sexual maturity and demographics of Blanding's turtles (*Emydoidea blandingii*): Implications for conservation and management of long-lived organisms. Conservation Biology **7**:826-833.

Cowles, H. C. 1901. The plant societies of Chicago and vicinity. Geographic Society of Chicago Bulletin **2**:1-76.

Ebbers, B. C. 1984. Reconstruction of the floristic environment in central Berrien County, Michigan, 2000 B.C. to nineteenth Century A.D. Pages 83-107 *in* E. B. Garland, editor. Late Archaic and early Woodland adaptation in the lower St. Joseph River valley, Berrien County, Michigan. Unpublished Report to the State of Michigan Department of Transportation.

Griggs, D., and S. Balzano. Personal communication, May 1990.

Heyer, W. R., M. A. Donnelly, R. W. McDiarmid, L. C. Hayek, and M. S. Foster. 1994. Measuring and monitoring biological diversity: Standard methods for amphibians. Smithsonian Institution Press, Washington, D.C., USA.

Holman, J. A., and K. D. Andrews. 1994. North American Quaternary cold-tolerant turtles: Distributional adaptations and constraints. Boreas **23**:44-52.

Harding, J. H., and J. A. Holman. 1992. Michigan frogs, toads, and salamanders. Michigan State University Museum.

Indiana Department of Natural Resources. 1992. Endangered, threatened, and rare species and high quality natural communities and natural areas documented from or adjacent to the Grand Calumet River on the Highland and Gary quadrangles in Lake County, Indiana. Database printouts. Division of Nature Preserves, Indianapolis, Indiana, USA.

Lake Michigan Federation. 1984. The Grand Calumet: Exploring the rivers potential. The Lake Michigan Federation, Chicago, Illinois, USA.

Madison, D. M. 1993. Tiger salamander habitat use. Unpublished report to the New York Department of Environmental Conservation, Endangered Species Unit. Delmar, New York, USA.

Mierzwa, K. S. 1993. Amphibians and reptiles of two macrosites in McHenry County, Illinois. Unpublished Report to the McHenry County Conservation District.

Mierzwa, K. S. In press. The status of Chicago region amphibians. *in* M. J. Lannoo, editor. Status and conservation of Midwestern amphibians. University of Iowa Press, Iowa City, Iowa, USA.

Mierzwa, K. S., S. Culberson, K. S. King, and C. Ross. 1991. Illinois-Indiana regional airport study: Biotic communities. Technical Paper No. 7, Appendix E, Volume II. TAMS Consultants, Inc., Chicago, Illinois, USA.

Minton, S. A., Jr. 1972. Amphibians and reptiles of Indiana. Indiana Academy of Science Monograph **3**, Indianapolis, Indiana, USA.

Morgan, D., and J. Burling. 1979. Listing of vertebrates seen on selected Lake County natural areas during August, 1978 with comments on preservation values. *in* Indiana Department of Natural Resources and Natural Land Institute. An inventory of natural areas in the Indiana Coastal Zone study area. Indiana State Planning Services Agency Technical Report **302**.

Resetar, A. 1988. Distribution and ecology of amphibians and reptiles in selected areas of Lake, Porter, LaPorte and Newton Counties, Indiana with emphasis on state-listed species. Unpublished Report to the Indiana Department of Natural Resources, Nongame and Endangered Wildlife Program, Indianapolis, Indiana, USA.

Richards, R. L. 1987. The vertebrate collection of the Indiana State Museum: "Old" specimens and records. Proceedings of the Indiana Academy of Science **97**:547-570.

Rossman, D. A., N. B. Ford, and R. A. Seigel. 1996. The garter snakes: Evolution and ecology. University of Oklahoma Press, Norman, Oklahoma, USA.

Schmidt, K. P., and W. L. Necker. 1935. Amphibians and reptiles of the Chicago region. Chicago Academy of Science Bulletin **5**:57-77.

Seibert, H. C. 1950. Population density of snakes in an area near Chicago. Copeia **1950**:229-230.

Shelford, V. E. 1913. Animal communities in temperate America, as illustrated in the Chicago

BIRDS

Kenneth J. Brock

Biology Department, Marram Hall
Indiana University Northwest
3400 Broadway
Gary, Indiana 46408-1197

INTRODUCTION

Birdlife, and especially breeding species of birds, constitutes a very sensitive environmental barometer. It is therefore surprising that, despite years of abuse, the Grand Calumet River system

Great Blue Heron, *Ardea herodias*. Several authors, including Butler (1898), suggest that the Great Blue Heron was a common nesting species across the northern half of Indiana. There is no evidence of local nesting, but *A. herodias* almost certainly occupied the Grand Calumet River floodplain during late summer dispersal.

Green Heron, *Butorides striatus*. Both Butler (1898) and Woodruff (1907) list this species as a common summer resident. It very likely inhabited scrubby trees along the River.

Black-Crowned Night Heron, *Nycticorax nycticorax*. Woodruff (1907) gives the following statement from E.W. Nelson, "The first of July, 1874, I saw a few young of the year in the Calumet Marshes." Birds present at this date could represent either locally fledged individuals or post-nesting dispersals from colonies outside the Grand Calumet River system.

Mallard,

Wilson's Phalarope, *Phalaropus tricolor*. According to Butler (1898), *P. tricolor* was a common breeding species in the extreme northwestern part of Indiana. The last recorded nesting was in 1941 (Mumford and Keller 1984).

Black Tern, *Chlidonias niger*. Black Terns were summer residents in marshes north of the Kankakee River; they bred commonly at some locations (Butler 1898). Mumford and Keller (1984) indicate that 40 nests were found on nearby Wolf Lake in 1924, rendering it likely that some nested within the Grand Calumet River system.

Marsh Wren, *Cistothorus palustris*. Butler (1898) referred to this species as an abundant resident of Indiana's marshes. *C. palustris* surely nested at every site that provided the requisite cattail stands.

Gray Catbird, *Dumetella carolinensis*.Woodruff (1907) deemed *D. carolinensis* to be an abundant summer resident in the Chicago region. It probably nested in dense shrubbery on the River floodplain.

Cedar Waxwing, *Bombycilla cedrorum*. A common summer resident in the Chicago area (Woodruff 1907), *Bombycilla cedrorum* very likely foraged, and perhaps nested, on the Grand Calumet River floodplain.

Yellow Warbler, *Dendroica petechia*. This species was an abundant summer resident of the Chicago region (Butler 1898). Undoubtedly, *D. petechia* nested along the Grand Calumet River.

Common Yellowthroat, *Geothlypis trichas*. Butler (1898) referred to this species as a common summer resident of the Chicago region. It almost certainly nested in cattail marshes on the Grand Calumet River floodplain.

Indigo Bunting, *Passerina cyanea*. According to Woodruff (1907), *P. cyanea* was a common summer resident within the Chicago area. It probably nested in scrubby trees on the River floodplain.

Nelson's Sharp-tailed Sparrow,en45
sumon
r ywherteinfloodplain. st certainly >BDC BT /TT0 1 Tf 0.0000 Tw 12 0 0 12 72 471.06128 Tm ( )Tj ET

Swamp Sparrow, *Melospiza georgiana*. Woodruff (1907) deemed this species to be a common summer resident of the Chicago area; it almost certainly nested in the floodplain marshes.

Red-Winged Blackbird, *Agelaius phoeniceus*. This widespread and common species (deemed to be an abundant summer resident by Butler (1898)), almost certainly nested in cattail marshes along the River.

Yellow-Headed Blackbird, *Xanthocephalus xanthocephalus*. According to Butler (1898) *X. xanthocephalus* was a summer resident of some localities in northwestern Indiana. It is likely that this species inhabited some of the cattail marshes along the Grand Calumet River.

Common Grackle, *Quiscalus quiscula*. Butler (1898) considered *Q. quiscula* a common summer resident; it probably foraged on the River floodplain.

Brown-Headed Cowbird, *Molothrus ater*. According to Butler (1898), *M. ater* was a common summer resident in Indiana. There is little doubt that it was present on the floodplain.

American Goldfinch, *Carduelis tristis*. This species, deemed to be a permanent resident of the Chicago region by Butler (1898), almost certainly bred in the Grand Calumet River marshes.

Pre-settlement migrants

The elongate geometry of Lake Michigan imposes a major geographical limitation on southbound migrants, directing thousands to the southern tip of Lake Michigan and across the Grand Calumet River (Brock 1986). As occurs today, untold numbers surely passed over the Grand Calumet River during historical migrations. Those species preferring habitats provided by the River floodplain probably stopped to feed and rest.

The literature provides little first-hand evidence as to which species were actually observed on the Grand Calumet River; indeed specific references were obtained for only the two following species:

Osprey, *Pandion haliaetus*. Woodruff (1907) reports seeing migrants on both Lake Calumet and Berry Lake. Osprey probably hunted over waters of the Grand Calumet River.

Le Conte's Sparrow, *Ammodramus leconteii*. Woodruff (1907) quotes E.W. Nelson, who refers to a specimen collected in May 1875, as follows, "...The specimen in my possession was flushed from a small depression in the prairie near the Calumet River (in Illinois)..."

Least Bittern, *Ixobrychus exilis. I. exilis*, an Indiana endangered species, is a rare summer

10, 1982. Counts of downy young at Roxanna include: seven on July 12, 1984, six on June 1, 1985, 18 on June 5, 1987, and 20 on July 3, 1987.

Blue-Winged Teal, *Anas discors*. *A. discors* is uncommon during the summer months. The peak summer count was four at Roxanna Pond on July 26, 1986, and a mean of one bird per year was recorded at that location during the past twelve years. Though it is quite likely that it occurs, breeding in the Grand Calumet River system has not been confirmed.

Ruddy Duck, *Oxyura jamaicensis*. Though there are several nesting records for this species in the Calumet Region, the only summer observation within the Grand Calumet River system involves a breeding plumed male sited at Roxanna Pond on June 24, 1983. Recent nesting has occurred on the Little Calumet River floodplain (Brock 1986).

Red-tailed Hawk, *Buteo jamaicensis*. During the summer period, this species is regularly seen flying above the floodplain. No nests have been found.

American Kestrel, *Falco sparverius*. This diminutive falcon is seen regularly in the Calumet Region during the summer months. Breeding on the floodplain has not been confirmed, but small numbers may nest.

Ring-necked Pheasant, *Phasianus colchicus*. This introduced species has been rec
orded regularly throughout the summer months at Roxanna Pond. It was also noted on the DuPont property in June by Mierzwa et al. (1991).

Virginia Rail, *Rallus limicola*. Tough rarely seen, *R. limicola*, which is listed among Indiana's "species of special concern," is a fairly common breeder in cattail marshes on the floodplain. Summer period birds were observed at Roxanna Pond on July 20, 1982 (seen by the author) and in the DuPont wetlands during the summer of 1991. Specific records at the latter site included a pair seen June 15, and a family group, two adults and two young, observed July 11, 1991 (Mierzwa et al. 1991).

Sora, *Porzana carolina*. Distribution of the Sora is similar to that of the Virginia Rail. June and July records include an adult with three young in the DuPont Marsh on July 11, 1991, an adult at Ivanhoe on June 11, 1991, and a single bird in a Clark & Pine swale on June 8, 1991 (Mierzwa et al. 1991). Additionally, fully feathered juveniles have been recorded at Roxanna Pond on four occasions during the fall; the largest count consisted of five, seen on September 16, 1995.

Common Moorhen, *Gallinula chloropus*. An abundant nesting species at Roxanna Pond during the 1980s, *G. chloropus* has declined in numbers markedly during the past six years. The mean summer count between 1980-1989 was 17.5 per year; in contrast, the average for annual June-July counts from 1990-1995 was only 1.3. Peak counts of young during the halcyon 1980s included 50 on June 28, 1984, and 13 on July 26, 1986.

American Coot,

Eastern Kingbird, *Tyrannus tyrannus*. The Kingbird probably breeds locally on the floodplain where it is seen regularly during the summer months. Breeding period records include eight birds on the DuPont property during the summer of 1991 (Mierzwa et al. 1991) and singletons at Roxanna Pond in 1985, 1986, and 1991.

Purple Martin, *Progne subis*. This large swallow is occasionally observed foraging over the floodplain during the summer months. Additionally, late July pre-migratory flocks are sometimes noted on power-line wires at Grand Calumet Lagoons (e.g., 20 birds seen on July 25, 1995, observed by the author).

Tree Swallow, *Tachycineta bicolor*. A few *T. bicolor* very likely nest locally on the floodplain wherever the requisite cavities or nest boxes are available. Many others forage over the waterways and wetlands of the Grand Calumet River. Summer period records include seven birds at DuPont, four at Roxanna, and one at the Cline Avenue bridge.

Barn Swallow, *Hirundo rustica*. *H. rustica* is an abundant summer resident of the floodplain; the species may nest beneath bridges that span the channel. The peak summer count is 70, recorded at Roxanna Pond on June 23, 1990.

Blue Jay, *Cyanocitta cristata*. This widespread corvid is regularly noted on the floodplain during the summer months.

American Crow, *Corvus brachyrhynchos*. Crows occasionally forage on the floodplain during the summer months. The largest count was four birds, seen at Roxanna Pond on June 23, 1990.

House Wren, *Troglodytes aedon*. This species breeds sparingly in oak savannah and scrubby second growth areas of the floodplain.

Marsh Wren, *Cistothorus palustris*. *C. palustris*, a threatened species in Indiana, is a common nesting species in cattail stands along the channel and in adjacent marshes. Summer records include 31birds (over 14 years) at Roxanna Pond and 35 at the DuPont marsh in 1991. Birds performing display flights were observed at the latter site on July 11, 1991 (Mierzwa et al. 1991).

American Robin,

Brown Thrasher, *Toxostoma rufum.* The thrasher is a local breeder that frequents the oak savannah habitat. Mierzwa et al. (1991) recorded this species regularly on the DuPont property.

Cedar Waxwing, *Bombycilla cedrorum*. *B. cedrorum* is a fairly common summer resident and sporadic breeder on the floodplain. A peak summer count of four was recorded in 1991 by Mierzwa et al. (1991).

European Starling, *Sturnus vulgaris*. This invasive species, first recorded in the Chicago area in

Red-throated Loon,

American Avocet, *Recurvirostra americana*. A singleton was observed at Roxanna Pond on May 11, 1980 (seen by the author).

Hudsonian Godwit, *Limosa haemastica*. Eight birds were seen at Roxanna Pond on May 11, 1978. This record constitutes the largest number ever recorded in Indiana (Mumford and Keller 1984).

Marbled Godwit, *Limosa fedoa*. One bird was seen at Roxanna Pond on May 10 and 11, 1978 (Brock 1986).

Ruddy Turnstone, *Arenaria interpres*. A flock of 15 birds was seen at Roxanna Pond on May 26, 1988 (seen by the author).

Red Knot, *Calidris canutus*. This species is rarely recorded away from the sandy beaches of Lake Michigan. A juvenile was seen along the muddy channel banks in the Hammond Sanitary

Red-Necked Phalarope, *Phalaropus lobatus*. Only one record of this species exists for the Grand Calumet River system; a single bird was seen at Roxanna Pond on July 15, 1982 (seen by the author).

Franklin's Gull, *Larus pipixcan*. An adult bird was observed on the Grand Calumet Lagoons, on November 5, 1991 (seen by the author).

White-Winged Tern, *Chlidonias leucopterus*. Indiana's only record of this Eurasian species is a bird seen at Roxanna Pond on July 19, 1979 (Brock, 1983).

Golden-winged Warbler, *Vermivora chrysoptera*. This Indiana endangered species was recorded at the DuPont wetlands on August 23, 1991 (Mierzwa et al. 1991).

Northern Parula, *Parula americana*. This warbler, which is rare in the Calumet Region, was observed in cottonwoods along the Grand Calumet River channel in the Hammond Sanitary District reach on September 27, 1980 (seen by the author).

Connecticut Warbler, *Oporornis agilis*. On May 24, 1991, a migrant was flushed from the oak savannah at DuPont by Mierzwa et al. (1991).

Contemporary winter residents

Birds that winter in the Grand Calumet River corridor are primarily aquatic species that take advantage of open water created by effluent from local industries. All species listed below have been recorded on or above the floodplain during the winter months (December throughd o,/MCID the winter mo

*Common Moorhen, *Gallinula chloropus*
American Coot, *Fulica americana*
Ring-billed Gull, *Larus delawarensis*
Herring Gull, *Larus argentatus*
*Glaucous Gull, *Larus hyperboreus*
Rock Dove, *Columba livia*
Mourning Dove, *Zenaida macroura*
Belted Kingfisher, *Ceryle alcyon*
Downy Woodpecker, *Picoides pubescens*
Blue Jay, *Cyanocitta cristata*
American Crow, *Corvus brachyrhynchos*
Cedar Waxwing, *Bombycilla cedrorum*
European Starling, *Sturnus vulgaris*
Northern Cardinal, *Cardinalis cardinalis*
American Tree Sparrow, *Spizella arborea*
*Fox Sparrow, *Passerella iliaca*
Song Sparrow, *Melospiza melodia*
Swamp Sparrow, *Melospiza georgiana*
Dark-Eyed Junco, *Junco hyemalis*
Red-Winged Blackbird, *Agelaius phoeniceus*
Common Grackle, *Quiscalus quiscula*
Brown-Headed Cowbird, *Molothrus ater*
House Finch, *Carpodacus mexicanus*
*Common Redpoll, *Carduelis flammea*
American Goldfinch, *Carduelis tristis*

IMPACT OF DREDGING

Considerable evidence suggests that the muddy substrate of the Grand Calumet River channel is contaminated (see USFWS 1996 for a discussion of the contaminants and potential effects on birdlife). The detection of Hg and PCB residues in failed eggs at the East Chicago Peregrine Falcon nest site (USFWS 1996) indicates that some contaminants have entered the avian food web.

The presence of pollutants in the muddy substrate poses an unassessed potential risk for birds that feed in the channel. Many birds spend extensive time foraging in this contaminated environment. On December 30, 1995, for example, 970 Mallards, plus other waterfowl species, were counted in the Roxanna marsh and Hammond Sanitary District reaches. These birds winter on the River, apparently feeding exclusively on channel vegetation. Although the long-term effects of prolonged exposure to Grand Calumet River pollutants on birds are unknown, lengthy exposure is certainly undesirable.

Perhaps at even greater risk are the migrant shorebirds that consume invertebrate infauna living within the substrate. Most migrant shorebirds depart within a few days, rendering their exposure brief, but the potential for accumulating toxins is high because they feed on organisms extracted directly from the contaminated sediment.

Thus, it is very likely that removal of contaminated sediment through dredging will have a positive long-term effect on the avifauna. A major concern is that the dredging operation itself might increase contamination through the resuspension of sediment during dredging. This possibility should be examined carefully within the context of the dredging mechanism employed.

OPPORTUNITIES

Roxanna Pond lies within the Grand Calumet River meander to the north of Roxanna Street and about 0.4 km to the west of Indianapolis Boulevard (Roxanna Reach of the Grand Calumet River). Water levels in this shallow pond fluctuate widely, and they are seemingly more dependent upon the rate of effluent discharge from local industries than on precipitation rates. Low water levels expose extensive mudflats, which provide feeding habitat for migrant shorebirds. When water levels are appropriate, the pond serves as a resting and feeding area for hundreds of spring and fall shorebird migrants. High water conditions, however, entirely eliminate the mudflats; on these occasions shorebirds cannot land at the pond.

At least 29 different shorebird species have been recorded at Roxanna Pond. The most cothesy.rnd species ar: Les se Yeallolegns,LleastSlanpipver,landPfecorcalSlanpiper. Swevecalra(re)]TJ0 -1.164

July through October), 2) that pre-migration water levels are appropriate for development of an adequate invertebrate infauna, on which shorebirds can feed, and 3) controlling vegetation to maintain mudflat habitat. Vegetation will be controlled by flooding during the non-migratory periods.

LITERATURE CITED

Brock, K. J. 1983. Indiana's first White-winged Black Tern: an Inland Sight Record. American Birds **37**:109-111.

Brock, K.J. 1986. Birds of the Indiana Dunes. Indiana University Press, Bloomington, Indiana, USA.

Buskirk, W. B. 1993. Unpublished Minutes of the July 23, 1993 IDNR Nongame Bird Technical Advisory Committee.

Butler, Amos W. 1998. The Birds of Indiana. 22nd Annual Report, Indiana Department of Geology and Natural Resources, pp. 575-1187.

Ford, E. R., C. C. Sanborn, and C. B. Coursen. 1934. Birds of the Chicago Region. Chicago Academy of Science, Chicago, Illinois, USA.

Helmers, D. L. 1992. Shorebird Management Manual. Western Hemisphere Shorebird Reserve Network. Manomet, Massachusetts, USA.

Kleen, V. M. 1979. The Changing Seasons: Spring 1978. American Birds **32**:1014.

Mlodinow, S. 1984. Chicago Area Birds. Chicago Review Press, Chicago, Illinois, USA.

Mumford, R. E., and C. E. Keller. 1984. The Birds of Indiana. Indiana University Press, Bloomington, Indiana, USA.

Nelson, E. W. 1875. Proceedings of the Boston Society of Natural History XVII.

Peterjohn, B. G. 1987. The Changing Seasons: Fall 1986. American Birds **41**:96.

Mierzwa, K.S., S. Culberson, K.S. King, and C. Ross. 1991. Illinois-Indiana regional airport study: Biotic communities. Technical Paper No. 7, Appendix E, Volume II. TAMS Consultants, Inc., Chicago, Illinois, USA.

U.S. Fish and Wildlife Service. 1996. Endangered Species Act, Consultation- Biological Opinion: Dredging of Indiana Harbor and Canal. 46p.

West, H. C. 1956. The Status of the Grebe Family in Indiana. Indiana Audubon Quarterly **34**:42-55.

Woodruff, F. M. 1907. The Birds of the Chicago Area. Natural History Survey, Bulletin No VI, Chicago Academy of Sciences, Chicago, Illinois, USA.

Attachment 1

Migrant Birds

Common Goldeneye, *Bucephala clangula*		X
Bufflehead, *Bucephala albeola*	X	X
Hooded Merganser, *Lophodytes cucullatus*	X	X
Common Merganser, *Mergus merganser*	X	X
Red-Breasted Merganser, *Mergus serrator*	X	X
Ruddy Duck, *Oxyura jamaicensis*	X	X
e Osprey, *Pandion haliaetus*	X	X

	Historical	Modern
e Northern Harrier, *Circus cyaneus*	X	X
Cooper's Hawk, *Accipiter cooperii*	X	X
Red-tailed Hawk, *Buteo jamaicensis*	X	X
American Kestrel, *Falco sparverius*	X	X

White-Winged Tern, *Chlidonias leucopterus*		X
e Black Tern, *Chlidonias niger*	X	X
Mourning Dove, *Zenaida macroura*		X

	Historical	Modern
Yellow Warbler, *Dendroica petechia*	X	X
Magnolia Warbler, *Dendroica magnolia*	X	X
Cape May Warbler, *Dendroica tigrina*	X	X
Yellow-Rumped Warbler, D*endroica coronata*	X	X
Black-Throated Green Warbler, *Dendroica virens*	X	X
Blackburnian Warbler, *Dendroica fusca*	X	X
Palm Warbler, *Dendroica palmarum*	X	X
Bay-breasted Warbler,		

Birds Distribution by Reach

The following bird species were identified within the Grand Calumet River corridor. It must be emphasized that the data are based on limited sampling; data for each reach were collected at only those few sites accessible from land.

Culverts reach

American Black Duck, *Anas rubripes*
Mallard, *Anas platyrhynchos*

Hammond Sanitary District reach

Pied-Billed Grebe, *Podilymbus podiceps*
Black-Crowned Night Heron, *Nycticorax nycticorax*
Canada Goose, *Branta canadensis*
Green-Winged Teal, *Anas crecca*
American Black Duck, *Anas rubripes*
Mallard, *Anas platyrhynchos*

American Robin, *Turdus migratorius*
Yellow-Rumped Warbler, D*endroica coronata*
Blackpoll Warbler, *Dendroica striata*
Song Sparrow, *Melospiza melodia*
Dark-Eyed Junco, *Junco hyemalis*
Red-winged Blackbird, *Agelaius phoeniceus*
Common Grackle, *Quiscalus quiscula*
Brown-Headed Cowbird, *Molothrus ater*
House Finch, *Carpodacus mexicanus*
American Goldfinch, *Carduelis tristis*

Roxanna Marsh reach

Pied-Billed Grebe, *Podilymbus podiceps*
Horned Grebe, *Podiceps auritus*
Eared Grebe, *Podiceps nigricollis*
Double-Crested Cormorant, *Phalacrocorax auritus*
Least Bittern, *Ixobrychus exilis*
Great Blue Heron, *Ardea herodias*
Great Egret, *Ardea albus*
Cattle Egret, *Bubulcus ibis*
Green Heron, *Butorides striatus*
Black-Crowned Night Heron, *Nycticorax nycticorax*
Mute Swan, *Cygnus olor*
Canada Goose, *Branta canadensis*
Snow Goose, *Chen caerulescens*
Wood Duck, *Aix sponsa*
Green-Winged Teal, *Anas crecca*
American Black Duck, *Anas rubripes*
Mallard, *Anas platyrhynchos*
Northern Pintail, *Anas acuta*
Blue-Winged Teal, *Anas discors*
Northern Shoveler, *Anas clypeata*
Gadwall, *Anas strepera*
American Wigeon, *Anas americana*
Canvasback, *Aythya valisineria*
Redhead, *Aythya americana*
Ring-necked Duck, *Aythya collaris*
Greater Scaup, *Aythya marila*
Lesser Scaup, *Aythya affinis*
Bufflehead, *Bucephala albeola*

Osprey, *Pandion haliaetus*
Northern Harrier, *Circus cyaneus*
Cooper's Hawk, *Accipiter cooperii*
Red-tailed Hawk, *Buteo jamaicensis*
American Kestrel, *Falco sparverius*
Peregrine Falcon, *Falco peregrinus*
Virginia Rail, *Rallus limicola*
Sora, *Porzana carolina*
Common Moorhen, *Gallinula chloropus*
American Coot, *Fulica americana*
Black-Bellied Plover, *Pluvialis squatarola*
American Golden Plover, *Pluvialis dominicus*
Semipalmated Plover, *Charadrius semipalmatus*
Killdeer, *Charadrius vociferus*
American Avocet, *Recurvirostra americana*
Greater Yellowlegs, *Tringa melanoleuca*
Lesser Yellowlegs, *Tringa flavipes*
Solitary Sandpiper, *Tringa solitari*a
Spotted Sandpiper, *Actitis macularia*
Hudsonian Godwit, *Limosa haemastic*a
Marbled Godwit, *Limosa fedoa*

Common Nighthawk, *Chordeiles minor*
Chimney Swift, *Chaetura pelagica*
Belted Kingfisher, *Ceryle alcyon*
Yellow-Bellied Sapsucker, S*phyrapicus varius*
Northern Flicker, *Colaptes auratus*
Willow Flycatcher, *Empidonax traillii*
Eastern Phoebe, *Sayornis phoebe*
Eastern Kingbird, *Tyrannus tyrannus*
Purple Martin, *Progne subis*
Tree Swallow, *Tachycineta bicolor*
N. Rough-winged Swallow, *Stelgidopteryx serripennis*
Bank Swallow, *Riparia riparia*
Cliff Swallow, *Hirundo pyrrhonota*
Barn Swallow, *Hirundo rustica*
Blue Jay, *Cyanocitta cristata*C
American Crow, *Corvus brachyrhynchos*
Black-Capped Chickadee, *Parus atricapillus*
Brown Creeper, *Certhia americana*
House Wren, *Troglodytes aedon*
Winter Wren, *Troglodytes troglodytes*
Marsh Wren, *Cistothorus palustris*
Golden-Crowned Kinglet, *Regulus satrapa*
Ruby-Crowned Kinglet, *Regulus calendula*
Gray-cheeked Thrush, *Catharus minimus*
Swainson's Thrush, *Catharus ustulatus*
Hermit Thrush, *Catharus guttatus*
American Robin, *Turdus migratorius*
Gray Catbird, *Dumetella carolinensis*
American Pipit, *Anthus spinoletta*
Cedar Waxwing, *Bombycilla cedrorum*
European Starling, *Sturnus vulgaris*
Warbling Vireo, *Vireo gilvus*
Red-eyed Vireo, *Vireo olivaceus*
Tennessee Warbler, *Vermivora peregrina*
Orange-crowned Warbler, *Vermivora celata*
Nashville Warbler, *Vermivora ruficapilla*
Northern Parula, *Parula american*
Yellow Warbler, *Dendroica petechia*
Magnolia Warbler, *Dendroica magnolia*
Cape May Warbler, *Dendroica tigrina*
Yellow-Rumped Warbler, *Dendroica coronata*
Black-Throated Green Warbler, *Dendroica virens*
Palm Warbler, *Dendroica palmarum*
Bay-breasted Warbler, *Dendroica castanea*
Blackpoll Warbler, *Dendroica striata*

American Redstart, *Setophaga ruticilla*
Northern Waterthrush,

American Woodcock, *Scolopax minor*
Ring-billed Gull, *Larus delawarensis*
Herring Gull, *Larus argentatus*
Black Tern, *Chlidonias niger*
Mourning Dove, *Zenaida macroura*
Common Nighthawk, *Chordeiles minor*
Chimney Swift, *Chaetura pelagica*
Belted Kingfisher, *Ceryle alcyon*
Downy Woodpecker, *Picoides pubescens*
Northern Flicker, *Colaptes auratus*

Blackpoll Warbler, *Dendroica striata*
American Redstart, *Setophaga ruticilla*
Ovenbird, *Seiurus aurocapillus*
Northern Waterthrush, *Seiurus noveboracensis*S
Connecticut Warbler, *Oporornis agilis*
Mourning Warbler, *Oporornis philadelphia*
Common Yellowthroat, *Geothlypis trichas*
Wilson's Warbler, *Wilsonia pusilla*
Canada Warbler, *Wilsonia canadensis*
Common Yellowthroat, *Geothlypis trichas*
Indigo Bunting, *Passerina cyanea*
Eastern Towhee, *Pipilo erythrophthalmus*
Fox Sparrow, *Passerella iliaca*
Song Sparrow, *Melospiza melodia*
Swamp Sparrow, *Melospiza georgiana*
White-throated Sparrow, *Zonotrichia albicollis*
White-crowned Sparrow, *Zonotrichia eucophrys*
Dark-Eyed Junco, *Junco hyemalis*
Red-winged Blackbird, *Agelaius phoeniceus*
Eastern Meadowlark, *Sturnella magna*
Common Grackle, *Quiscalus quiscula*
Brown-Headed Cowbird, *Molothrus ater*
House Finch, *Carpodacus mexicanus*
American Goldfinch, *Carduelis tristis*

Gary Sanitary District reach

Double-Crested Cormorant, *Phalacrocorax auritus*
Great Blue Heron, *Ardea herodias*
Great Egret, *Ardea albus*
Mallard, *Anas platyrhynchos*
Lesser Scaup, *Aythya affinis*
American Kestrel, *Falco sparverius*
Semipalmated Plover, *Charadrius semipalmatus*
Killdeer, *Charadrius vociferus*
Lesser Yellowlegs, *Tringa flavipes*
Solitary Sandpiper, *Tringa solitaria*
Spotted Sandpiper, *Actitis macularia*
Semipalmated Sandpiper, *Calidris pusilla*
Western Sandpiper, *Calidris mauri*
Least Sandpiper, *Calidris minutilla*
Baird's Sandpiper, *Calidris bairdii*
Pectoral Sandpiper, *Calidris melanotos*
Stilt Sandpiper, *Calidris himantopus*
Ruff, *Philomachus pugnax*

Short-Billed Dowitcher, *Limnodromus griseus*
Common Snipe, *Gallinago gallinago*
Chimney Swift, *Chaetura pelagica*
Swamp Sparrow, *Melospiza georgiana*

Lagoons reach

Red-throated Loon, *Gavia stellata*
Common Loon, *Gavia immer*
Pied-Billed Grebe, *Podilymbus podiceps*
Horned Grebe, *Podiceps auritus*
Red-Necked Grebe, *Podiceps grisegena*
American White Pelican, *Pelecanus erythrorhynchos*
Double-Crested Cormorant, *Phalacrocorax auritus*
Great Blue Heron, *Ardea herodias*
Cattle Egret, *Bubulcus ibis*
Green Heron, *Butorides striatus*
Tundra Swan, *Cygnus columbianus*
Mute Swan, *Cygnus olor*
Canada Goose, *Branta canadensis*
Wood Duck, *Aix sponsa*
Green-Winged Teal, *Anas crecca*
American Black Duck, *Anas rubripes*
Mallard, *Anas platyrhynchos*
Blue-Winged Teal, *Anas discors*
Gadwall, *Anas strepera*
Canvasback, *Aythya valisineria*
Redhead, *Aythya americana*
Ring-necked Duck, *Aythya collaris*
Black Scoter, *Melanitta nigra*
White-Winged Scoter, *Melanitta fusca*
Common Goldeneye, *Bucephala clangula*
Bufflehead, *Bucephala albeola*
Hooded Merganser, *Lophodytes cucullatus*
Common Merganser, *Mergus merganser*
Red-Breasted Merganser, *Mergus serrator*
Ruddy Duck, *Oxyura jamaicensis*
Cooper's Hawk, *Accipiter cooperii*
Red-tailed Hawk, *Buteo jamaicensis*
American Coot, *Fulica americana*
Franklin's Gull, *Larus pipixcan*
Ring-billed Gull, *Larus delawarensis*
Herring Gull, *Larus argentatus*
Chimney Swift, *Chaetura pelagica*
Belted Kingfisher, *Ceryle alcyon*

Yellow-Bellied Sapsucker, S*phyrapicus varius*
Northern Flicker, *Colaptes auratus*
Purple Martin, *Progne subis*
Barn Swallow, *Hirundo rustica*
Blue Jay, *Cyanocitta cristata*C
American Crow, *Corvus brachyrhynchos*
Cedar Waxwing, *Bombycilla cedrorum*
European Starling, *Sturnus vulgaris*
Cape May Warbler, *Dendroica tigrina*
Yellow-Rumped Warbler, *Dendroica coronata*
White-crowned Sparrow, *Zonotrichia eucophrys*
Dark-Eyed Junco, *Junco hyemalis*
Red-winged Blackbird, *Agelaius phoeniceus*
House Finch, *Carpodacus mexicanus*
American Goldfinch, *Carduelis tristis*

MAMMALS

John O. Whitaker, Jr.

Department of Life Sciences
Indiana State University
Terre Haute, Indiana 47809

INTRODUCTION

The objectives of this chapter are to describe the pre-settlement and present mammal communities of the Grand Calumet River basin, and to discuss how dredging operations may affect these communities. A further objective is to produce recommendations that might be implemented during the dredging operations to enhance the mammal populations of the area.

Fisher, *Martes pennanti*. Rand and Rand (1951) found fisher remains in the region, indicating that they were clearly present. The last report of a fisher in Indiana was in 1859 (Mumford and Whitaker, 1982).

River otter, *Lutra canadensis*. The last record of the river otter in northwestern Indiana is from 1900. The otter was first reintroduced into Indiana at Muscatatuck National Fish and Wildlife Refuge in 1995. Additional releases were made in other areas in 1996 and 1997, and more are planned. The otter perhaps could be reintroduced eventually into northwestern Indiana.

Mountain lion, *Felis concolor*. Panthers were reported in the 1660's, and the last mountain lion recorded from Indiana was seen in 1830, although there is some question as to whether it was identified correctly.

Canada lynx, *Felis lynx*. The last Canada lynx in the region was reportedly killed by Hunter Green in 1873 at Tremont, although the identification of this species in Indiana has been questioned. The last record in the state is from 1880.

Three papers present data from areas actually within the Grand Calumet River basin: Whitman (1986), Mierzwa et al. (1991), and Whitaker et al. (1994).

Whitman et al. (1990) found 16 species of mammals at Miller Woods: opossum, short-tailed shrew, masked shrew, eastern mole, cottontail, white-footed mouse, prairie deer mouse, meadow vole, muskrat, fox squirrel, gray squirrel, thirteen-lined ground squirrel, red squirrel, raccoon, long-tailed weasel, and white-tailed deer.

Mierzwa et al. (1991) studied mammals at 15 sites in five different study areas in northeastern Illinois and northwestern Indiana as possible sites at which to establish an airport. Two of the areas were in the Grand Calumet River area: one at an area in Gary, the other near Lake Calumet. However, they reported only ten species of mammals in these two areas (Attachment 2): Virginia opossum, masked and short-tailed shrews, gray and Franklin's ground squirrels, deer and white-footed mice, meadow vole, house mouse and Norway rat.

Whitaker et al. (1994) included information on Miller Woods, but otherwise they did not study areas within the Grand Calumet River basin. However, the habitats, and therefore the mammals, of the Grand Calumet River basin are similar to those of the Indiana Dunes National Lakeshore. Therefore, information from that work and from other papers on the Indiana Dunes were used extensively in this work.

The first significant publication on the mammals of the Indiana Dunes was by Lyon (1923). Lyon reported 22 species: opossum, short-tailed shrew, eastern mole, eastern red bat, eastern cottontail, eastern chipmunk, woodchuck, thirteen-lined ground squirrel, fox squirrel, red squirrel, prairie deer mouse, white-footed mouse, prairie vole, woodland vole, muskrat, Norway rat, house mouse, red fox, raccoon, long-tailed weasel, mink, and striped skunk. He did not personally observe specimens of eastern mole, muskrat, Norway rat, raccoon (a few were taken

Brennan (1923) relates early reports of bison, black bear (last seen in the region in 1871, when it was probably driven there from Michigan by the great fire; the entire east shore of Lake

ten species of carnivores (coyote, red and gray foxes, raccoon, long-tailed and least weasels, mink, American badger, striped skunk, and feral cat), and white-tailed deer. Gray fox brought to 38 the number of species known to occur at the Lakeshore.

The diverse habitats of the Lakeshore thus create a home for approximately 38 of the 57 species of mammals presently known to occur in Indiana (including the river otter which was reintroduced in Indiana in 1995). The beaver and white-tailed deer were extirpated from the state during the last century, but both occur there now. Franklin's ground squirrel was probably present at the Lakeshore through the 1940's, but then it apparently disappeared from there. However, it was recorded in the Grand Calumet River basin by Mierzwa et al. (1991). Feral dogs and cats are present, and they may partially fill the predator niche. Bobcat (state endangered) signs have been reported in the Heron Rookery area of the Lakeshore, but their presence has not been verified.

Species of mammals suspected to currently reside in the Grand Calumet River basin are discussed below and are indicated in Attachment 1.

Didelphidae, the opossums

Virginia opossum, *Didelphis virginiana* Kerr. The opossum is common in the Lakeshore area, and likewise it is surely common throughout the Grand Calumet River region. It was found in ten of the 24 habitats sampled at the Indiana Dunes National Lakeshore and also 93 were recorded as roadkill (Whitaker et al. 1994). Texas Instruments Inc. found this species in all six of the terrestrial habitats they sampled, and Whitman et al. (1990) found it to be common at Miller Woods. Mierzwa et. al. (1991) recorded one in Eggers Woods, near Wolf Lake.

Insectivora, shrews and moles

The Insectivora consists of the moles and shrews, six species of which conceivably could occur in the Grand Calumet River area, four species of shrews and two of moles. Only three species of shrews (masked, short-tailed and least) and one mole (eastern) have been confirmed to occur there.

Northern short-tailed shrew, *Blarina brevicauda*. The short-tailed shrew is one of the most common mammals in northwestern Indiana, and it is common in the Grand Calumet River basin (Mumford and Whitaker 1982, Krekeler 1981, Texas Instruments 1975-80, Whitman et al. 1990, and Whitaker et al. 1994). Whitaker et al. (1994) took short-tailed shrews in 17 of the 24 habitats studied. They were most abundant in upland terrestrial shrubland, wet prairie, old field, ephemeral lowland forest, and mixed deciduous savanna. Mierzwa et al. (1991) trapped nine short-tailed shrews: four at the DuPont area, four at the Burnham Prairie/Powderhorn Prairie area, and one at the Clark & Pine area.

Least shrew, *Cryptotis parva* (Say).The least shrew is a small brownish short-tailed shrew, much smaller than *Blarina*. Its total length is only about 63 to 88 mm, its tail only 11 to 20 mm. It usually occurs in fairly dry open fields. It occurs throughout Indiana, but it is not taken often. There are few records in the northern part of the state. Lyon trapped a least shrew in "subdunal

woods" on October 31, 1924. It was apparently from Tremont, as Sanborn (1925) reported that Lyon took one there in the fall of 1924. The specimen was deposited in the U.S. National Museum (#240630). Whitaker et al. (1994) did not take it at the Indiana Dunes National Lakeshore, but it most likely occurs sparingly in dry fields in the Calumet River area.

Masked shrew, *Sorex cinereus* Kerr. The masked shrew is common in several habitats at the Indiana Dunes National Lakeshore and in the Grand Calumet River area, especially in wet areas. Mierzwa et al. (1991) captured 35 individuals in their Lake Calumet study area: four in the Big Marsh/Indiana Ridge area, 20 in the Burnham Prairie/Powderhorn Prairie area, and 11 at Egger's Woods. Mumford and Whitaker (1982) found masked shrews in several habitats east of the Bailey Generating Station, and 124 of 178 mammals (69.7 %) taken at Cowles and Pinhook Bogs and at Trail Creek Fen were masked shrews. Whitman et al. (1990) reported masked shrew from Miller Woods. Whitaker et al. (1994) took only 35 during their study of mammals at the Indiana Dunes National Lakeshore. This included individuals from eight habitats, although they were most abundant in wet prairie (1.83 per 100 trap-nights) and marsh (1.17 per 100 trap-nights). Shrew populations were apparently low at the time of this study. The masked shrew lives in areas where the soil retains moisture sufficient to maintain burrows 100% saturated. Because of this moisture requirement, the species often lives in dense vegetation or in mossy areas.

Pygmy shrew, *Sorex (Microsorex) hoyi* Baird. The pygmy shrew has long been thought to be exceedingly rare, but pitfall trapping has shown it is much more abundant than was previously realized. The pygmy shrew could occur in the Calumet River basin, but it has not been found at the Indiana Dunes National Lakeshore or elsewhere in northern Indiana. It does not occur in the southern portion of the lower peninsula of Michigan either (Baker 1983). There are records from Wisconsin including one at the extreme southeast corner, or less than 80 miles from the Grand Calumet River area. Also, one was taken in mid-winter in a garage at Palatine which is northwest of Chicago in Cook County, Illinois, and about 50 miles from the Grand Calumet. No pygmy shrews were taken in extensive trapping at Bailly, Cowles', or Pinhook Bogs (Mumford and Whitaker 1982) and none were taken in other areas at the Indiana Dunes National Lakeshore (Whitaker et al. 1994). Thus, this shrew's occurrence in the Grand Calumet River area is unlikely. However, the unicuspid teeth of all long-tailed shrews (*Sorex*) from that area should be carefully examined. (*Sorex cinereus* has four "large" unicuspids and one small unicuspid, all easily visible from the side. *Sorex hoyi* has the third and fifth extremely reduced, thus only 3 unicuspids are readily visible from the side).

Talipidae, the moles

Eastern mole, *Scalopus aquaticus* (Linnaeus). The eastern mole is common at Indiana Dunes National Lakeshore (Krekeler 1981, Whitman et al. 1990, Whitaker et al. 1994) and in the Grand Calumet River region. Whitaker et al. (1994) found burrows of the eastern mole in ten of 24 habitats examined at the Indiana Dunes National Lakeshore. This species was most abundant

Grand Calumet River basin. The eastern mole is common in many of the drier habitats at the Lakeshore.

Star-nosed mole, *Condylura cristata* (Linnaeus). The star-nosed mole is a species that lives in muckland habitats. Its burrows are usually quite evident, since they are in muck, whereas the burrows of eastern moles are usually in drier areas. The star-nosed mole has been documented only in the northeast portion of the state, and its range has apparently contracted in this century.

July of 1987 at the West Beach bathhouse; 2) a red bat observed flying at Indiana Dunes State Park on August 27, 1988; and 3) two individuals netted (1 male, 1 female) over Dunes Creek, Indiana Dunes State Park, on September 23, 1988. The red bat is solitary and hangs in trees during the daytime. It is one of the most common bats in wooded areas of northwestern Indiana, and it should occur in areas of the Grand Calumet River basin where enough trees are present.

Little brown myotis, *Myotis lucifugus* (LeConte). Neither Lyon (1936) nor Mumford and Whitaker (1982) recorded the little brown myotis in Lake, Porter, or LaPorte Counties. The only specific record of this species at INDU is that of Texas Instruments (1975-80), but this record needs verification. The author has often seen small bats flying about over openings in Indiana Dunes State Park and vicinity which I think are this species and/or perhaps the northern myotis, *Myotis septentrionalis*. The little brown myotis migrates to the karst regions of southern Indiana where it hibernates in caves.

Silver-haired bat, *Lasionycteris noctivagans* (LeConte). Hahn (1909) reported this species from Michigan City, LaPorte County. Mumford and Whitaker (1982) reported three individuals taken at the Indiana Dunes State Park, a female taken on September 24, 1928 by W.A. Weber, and two individuals taken on May 3, 1936 by J. Schmidt. Whitaker et al. (1994) netted one individual on September 9, 1988, about one mile northeast of the visitors center at the INDU.

The silver-haired bat is a migratory solitary bat. It spends the summer and has its young to the north of Indiana and then migrates southward. A few individuals hibernate in caves or mines in southern Indiana, but most winter in states to the south. This species is fairly common in Indiana during migration from about April 18 to May 28 and from about August 29 to November 6, when it should be relatively common at the Lakeshore (Mumford and Whitaker 1982).

Evening bat, *Nycticeius humeralis* (Rafinesque). Russell E. Mumford shot an evening bat two miles to the northwest of Porter (Porter County) on August 5, 1958. However, populations of this species have decreased greatly in Indiana in recent years, and we doubt that it presently occurs in northwestern Indiana. For that matter, the single record could have been a stray.

Bat species probably present but unsupported by records

Northern myotis, *Myotis septentrionalis* (Merriam). This species has often been referred to as Keen's bat, *Myotis keenii septentrionalis*. However, the populations in the eastern regions of the United States are currently recognized as a separate species from the populations in the western U.S. (Van Zyll de Jong 1979). The eastern species is known as the northern myotis, *Myotis septentrionalis*. There are no records of this species for any of the Lake shore counties (Mumford and Whitaker 1982). However, since wooded habitat is abundant, and since it is a northern species, the author suspects further work will reveal it to be present. It forms small summer colonies under the bark of trees or in buildings, it then migrates to caves and mines where it hibernates individually rather than in groups. Kurta (1982) found it was relatively uncommon in southern Michigan, and Long (1974) reported it as less common than the little brown myotis in the Lake Michigan drainage.

Hoary bat, *Lasiurus cinereus* (Palisot de Beauvois). Like the red and silver-haired bats, this is a solitary, migratory species which roosts in trees. It is the largest and one of the most colorful bats of Indiana. It occurs throughout the state, but it is nowhere common. It probably occurs in the Grand Calumet River basin in areas with adequate trees.

Indiana myotis, *Myotis sodalis* (Miller and Allen). The Indiana myotis is listed as endangered primarily because huge numbers congregate in very few caves in winter. There are no records for the northwestern 15 or so counties of the state (Mumford and Whitaker 1982); however, Kurta (1982) recorded this species, and Kurta et al. (1993) later recorded a maternity colony of this species from southern Michigan. The Indiana myotis is probably present in northwestern Indiana, including the Grand Calumet River area, where enough forest with large trees is present. It forms small summer colonies under the bark of dead trees, often along watercourses. It hibernates in large numbers in a very few caves, some of which are found in southern Indiana.

Lagomorpha, rabbits and hares

Eastern cottontail, *Sylvilagus floridanus* (Allen). The eastern cottontail is the only lagomorph in northwestern Indiana. Lyon (1923), Krekeler (1981), Texas Instruments, and Whitaker et al. (1994) all listed it as common in the area of the Indiana Dunes National Lakeshore. It was observed occasionalThe 079 Tc -0.00079 Tw 18.wTmeDdtieID iy6udnumm

gray squirrel and chipmunk are obvious where common. The flying squirrel may often be common, but it is seldom observed because it is nocturnal. The thirteen-lined ground squirrel has a spotty distribution, but it is found at several Indiana Dunes National Lakeshore localities (Whitaker et al. 1994). Gray and Franklin's ground squirrels have been documented within the Grand Calumet River basin (Mierzwa et al. 1991). The various squirrels are discussed below.

Eastern chipmunk, *Tamias striatus* (Linnaeus). The chipmunk is common in many of the wooded habitats of northwestern Indiana, although it is sporadic in occurrence. Lyon (1923) stated that "chipmunks do not appear to be very common in the dunes." Krekeler (1981) listed it as abundant in open woods, thickets, and suburbs. Texas Instruments (1975-80) recorded numerous captures of chipmunks in several habitats. Whitman et. al. (1990) did not mention them in their Miller Woods report. Whitaker et. al. (1994) captured 24 chipmunks in nine habitats at the Indiana Dunes National Lakeshore, and sign or sight observations were made in 11 plots in seven habitats. Seventeen were seen dead on the road, and numerous individuals were seen elsewhere at INDU. The chipmunk should be fairly common in the Grand Calumet River basin, but Mierzwa et al. (1991) did not report it.

Woodchuck, *Marmota monax* (Linnaeus). Lyon (1923) and Whitaker et al. (1994) found woodchucks to be common in several habitats at the Indiana Dunes National Lakeshore. They were observed up to October 4. Whitman et al. (1990) does not mention them from Miller Woods, but Mierzwa noted them from DuPont and Burnham Prairie.

Thirteen-lined ground squirrel, *Spermophilus tridecemlineatus* (Mitchill). Lyon (1923) found this species "not uncommon along the Chicago, Lake Shore and South Bend Railway just south of the dunes," and reported one "just north of Oak Hill Station and a few feet above the subdunal swamp." Krekeler (1981) listed it as "common" at INDU and indicated its habitat as "pastures, road borders, dunes, weedy or cultivated fields." Texas Instruments (1975-80) reported three individuals: two from the Cowles Marsh area and one from the transmission corridor. Whitman et al. (1990) saw this species along the railroad bed "at the north end of the study area." Whitaker et al. (1994) found a roadkilled individual on Wagner Road just north of Route 20, and two additional roadkills just south of route 20. Fifteen individuals were trapped in nine plots in four habitats, four in dry prairie, four in terrestrial shrubland, three in oak savanna, and four in coniferous savanna. Ten of the 15 were at West Beach. Mierzwa et al. (1991) saw this species only at Wolf Lake where it was very common on the Illinois side of the lake on grassy roadsides. The species should be present in dry open areas with little ground cover near the Grand Calumet River.

Franklin's ground squirrel, *Spermophilus franklinii* (Sabine). Neither Lyon (1923), Krekeler (1981), Texas Instruments (1975-80), nor Whitaker et al. (1994) reported Franklin's ground squirrels from INDU. Three Franklin's ground squirrels were taken by Alex Bognar, from "Miller" in Lake County, Indiana in 1947. The specimens apparently are in the Field Museum (CNHM #'s 73872, 73873, and 73874). We assume these were from or near Miller Station, which is about half a mile south of Miller Woods. In 1986-87, Scott Johnson and other Indiana DNR personnel used two sets of ten live traps to examine the area along the railroad at Miller

Station for Franklin's ground squirrels. This locality would appear to be very close to the locality where Bognar collected this species, but no Franklin's ground squirrels were seen or taken there.

Mierzwa et al. (1991) trapped two Franklin's ground squirrels on low dunes between swales: one at the Dupont area and one at the Big Marsh/Indian Ridge area. Also, they found it to be common at Powderhorn Lake, and present at Burnham Prair -0a.2e5 at 1 Cala numet.s. pr or

Gray squirrel, *Sciurus carolinensis* (Gmelin). Lyon (1923) did not report this species from the Lakeshore area; however, Krekeler (1981) recorded it as common in woods and suburbs, and Texas Instruments (1975-80) reported 23 from three habitats at the Lakeshore. Whitaker et al. (1994) recorded it in upland oak forest and in oak savanna, and also reported several road killed individuals. Also, a number were seen in black oak forest at Miller Woods and one was observed in black oak forest at Dune Acres. Whitman et al. 1990 also observed gray squirrels at Miller Woods.

Beaver, *Castor canadensis* Kuhl. Beavers had nearly disappeared from Indiana by about 1840 (Lyon 1936). They were reintroduced into Jasper-Pulaski and Kankakee Fish and Wildlife Areas in 1935 from Wisconsin and Michigan, and later introductions were made into other areas. Brooks (1955) knew of 326 colonies in 43 counties of Indiana. The majority were in portions of northwestern Indiana and some were in Lake County. There is a photograph taken in 1968 of a beaver lodge at Cowles Bog (Lindsey et al. 1969). Krekeler (1981) listed the beaver as uncommon at INDU, but he did say that it had caused high water problems on the road leading into Dune Acres (at Cowles marsh). There is presently a beaver colony on the Salt Creek watershed, and signs of beaver were seen by Whitaker et al. (1994) on the lower portions of the Little Calumet River. There had been a colony on the Little Calumet north of Chesterton, but all of the beavers there have been trapped. Tom Sobat reported fresh beaver cuttings along Derby Ditch on October 17, 1990. Mierzwa et al. (1991) did not mention beaver in their studies of the Grand Calumet River area. The beaver should be a prominent species in wooded portions of a clean Grand Calumet River.

Muridae, mice and rats

Most mice and rats are currently placed in the family Muridae, which contains three subfamilies: the Sigmodontinae (previously the Cricetinae), the Arvicolinae (previously the Microtinae) and the Murinae. The Murinae are the Old World rats and mice, *Mus* and *Rattus*.

Sigmodontinae

The Sigmodontinae are the native rats and mice and constitute constitute one of the largest groups of mammals in North America both in number of species and in number of individuals. However, there are only two (or possibly three) species in this group in the Grand Calumet River basin, although one of them, *Peromyscus leucopus*, is the most abundant mammal there.

Western harvest mouse, *Reithrodontomys megalotis* (Baird). The western harvest mouse was probably first taken in Indiana in 1969 at Willow Slough State Fish and Wildlife area in Newton County (Whitaker and Sly 1970) when it moved into Indiana. By 1975 it had extended its range to include at least seven counties of northwestern Indiana (Ford 1975), but it had not crossed the Kankakee River to the north into Lake and Porter Counties. By 1994 (Whitaker unpublished), it had extended its range to the south into Vigo County. By 1995 it had crossed the Wabash River into Clay County, and by 1997 it had crossed the Kankakee. Now that that barrier has been crossed, harvest mice should become part of the fauna of the rest of northwestern Indiana.

Deer mouse, *Peromyscus maniculatus* (Wagner). The deer mouse occupies a variety of habitats ranging from woods to dry open areas. Its prime habitat in Indiana is in cultivated fields where it lives even when the fields are bare (Whitaker 1967). However, only the prairie deer mouse, *P. m. bairdii* occurs in Indiana. It lives in dry open areas. It is never found in woods, although it can be abundant in savanna with open sand. Unlike all of the other species of small mammals that occur in Indiana, its habitat is inversely related to plant cover, and it can live in areas with little

or even no herbaceous vegetation, such as recently plowed or harvested fields or open sandy areas. Its prime habitat in Indiana is in cultivated fields (Whitaker 1967; Mumford and Whitaker 1982). Its tracks are often obvious in loose sand on dunes, or in light snow in plowed fields, where it emerges from one burrow and enters another. It probably occurred in areas of open sand such as on dunes in presettlement times.

Hahn (1909) recorded this species from only five counties, and did not think it was very common in Indiana. Evermann and Butler (1920) stated that it was abundant on the dunes along Lake Michigan. Lyon (1923) found numerous footprints of this species in the loose sand on the dunes but took few specimens and concluded that a few mice made many tracks. Krekeler (1981) recorded it as abundant in foredunes, grasslands, thickets, and old fields. Whitman et al. (1990) recorded it from Miller Woods. Whitaker et al. (1994) recorded *Peromyscus maniculatus* from ten habitats, but it was most common in dry prairie and on right-of-ways at the Indiana Dunes National Lakeshore. Mierzwa et al. (1991) recorded 30 individuals in open sand on the low dunes of the Grand Calumet River area, and stated that they were "so common at Lakeshore Railroad Prairie that we were able to find them under boards and debris without even trapping for them." This site is the most open one in the Clark and Pine complex (Mierzwa, pers. comm.).

White-footed mouse, *Peromyscus leucopus* (LeConte). *Peromyscus leucopus* is the most abundant small mammal at the Indiana Dunes National Lakeshore (Lyon 1923, Texas Instruments 1975-80, Krekeler 1981, Whitman et al. 1990). Whitaker et al. (1994) took a total of 445 individuals in 22 of 24 habitats they studied and in 98 of the 168 (58.3%) plots in which they trapped. The only habitats from which the white-footed mouse was not taken were barren grounds and excavated areas, both habitats lacking plant cover. It was taken at its greatest abundance in ephemeral lowland forest, oak upland forest, mixed deciduous savanna, mixed deciduous upland forest, coniferous savanna, and pine plantations. It is probably also the most abundant species in the area of the Grand Calumet River, as Mierzwa et al. 1991 took 67 individuals of this species, and found it in every one of their study sites (Attachment 2).

Arvicolinae, the voles

All five species of microtines present in Indiana are probably present both at the Lakeshore and in the area of the Grand Calumet River.

Meadow vole, *Microtus pennsylvanicus* (Ord). The meadow vole is found throughout northern Indiana where it is usually the most abundant small mammal of moist grassy meadows and wet prairies. In dry fields with sparse vegetation, it tends to be replaced by the prairie vole, *M. ochrogaster*. Lyon (1923) took prairie voles and also pine voles at INDU, but inexplicably, he did not take meadow voles. Krekeler (1981) and Texas Instruments (1975-80) indicated it as abundant. Whitman et al. (1990) trapped them in low-lying areas around one pond at Miller Woods. Whitaker et al. (1994) trapped 171 individuals in 28 plots in 13 habitats. These animals were most abundant in old field and upland terrestrial shrubland. Thirty-four were taken in one plot in this latter habitat. The cover here was heavy and mainly of forbs, but it included dogwood, poisen ivy, roses, some young pines, and few grasses. The soil was quite moist.

Mierzwa et al. (1991) captured 25 meadow voles in five of their six study areas near the Grand

Rattus norvegicus and *Mus musculus* are introduced exotics and are usually found in habitats created or disturbed by humans, such as in buildings or cultivated fields.

Norway rat, *Rattus norvegicus* (Berkenhout). This is the common rat associated with garbage dumps, barns, grain storage units, and suburban warehouses. It is a major pest almost everywhere that it occurs. It ruins foods and other materials and carries disease. It is common in larger cities and on farms in northwestern Indiana. Lyon (1923) did not trap or see any rats but said that residents reported them. Krekeler (1981) reported them as common around farms suburbs and ditches. None were reported during the Texas Instruments studies. Whitaker et al. (1994) found rat droppings in abandoned buildings. Mierzwa et al. (1991) took one Norway rat at the Big Marsh/Indian Ridge area.

House mouse, *Mus musculus* Linnaeus. This mouse has been transported around the world by humans. In Indiana, it is very common in cultivated fields when adequate cover is available but, unlike the prairie deer mouse, it vacates immediately once the cover is removed. The prairie deer mouse and house mouse are the primary small mammals of the Indiana corn and soybean fields (Whitaker 1967), although *P. leucopus* is sometimes present as well. The species also invades beaches, offshore islands, and estuarine areas.

Lyon (1923) took two individuals, both in foredune areas, and Texas Instruments (1975-80) took five from young foredunes. Surprisingly, none were taken in any of the plots in the 24 habitats studied by Whitaker et al. (1994). Their only records were sight records at residential areas and some from Chellberg Farm. Two individuals were taken among 70 mammals from snap traps at Pinhook Bog by Whitaker and Mumford in 1978 (Mumford and Whitaker 1982). It is clear that the house mouse is not abundant at the Indiana Lakeshore, probably because of the lack of agricultural land present. The same is probably true in the Grand Calumet River basin, as Mierzwa et al. (1991) recorded only one house mouse. It was found at the Big Marsh/Indian Ridge site.

Dipodidae (previously Zapodidae)

Meadow jumping mouse, *Zapus hudsonius* (Zimmermann). Lyon (1923) concluded that *Z. hudsonius* was almost certainly present in the Lakeshore region. Texas Instruments (1975-80) reported four individuals from young foredunes and 23 from the transmission corridor. Whitaker et al. (1994) took 15 individuals from three habitats, 13 of which came from one plot in aquatic shrublands. Whitaker and R.E. Mumford took one at Cowles Bog and five at Pinhook Bog in the fall of 1978, and one from a flat depression behind the foredunes east of the Bailley Generating Station in October of 1974. The meadow jumping mouse was not taken by Mierzwa et al. (1991), but it is undoubtedly present in the Grand Calumet River area.

Carnivora

The raccoon is abundant and obvious, but the other carnivores of the Grand Calumet River area are difficult to assess and count. Being relatively large, they are much less abundant

than small mammals, and they are usually among the first to disappear as humans develop the land, because of habitat loss, trapping, and hunting. They are often thought of as problem animals and killed by the general public. For some species, roadkills may give us the best assessment of status.

The carnivores of northwestern Indiana that may be present in the Grand Calumet River basin are grouped in four families, the Canidae (coyote, two species of foxes, domestic dog), Procyonidae (raccoon), Mustelidae (two species of weasels, mink, badger, skunk), and the Felidae (bobcat, housecat).

Canidae

Coyote, *Canis latrans* Say. The coyote has always been present in Indiana, but in recent years its populations have increased. There are early reports of wolves and/or coyotes at INDU; however, wolves are long gone and there was no confirmed record of coyotes prior to the work of Whitaker et al. (1994). These authors report that one was seen by Noel Pavlovic at Tolleston Dunes on August 7, 1990, and that Lakeshore rangers have seen them several times near the Heron Rookery, starting in June and July of 1990. Dan Fagre saw one in a cornfield just south of INDU in 1991. It is not clear yet whether coyotes have taken up residence at the Lakeshore, but if not, it appears to be only a matter of time until they do. Unfortunately, Rand and Rand (1951) found no canid bones which could have helped determine whether coyotes or wolves or both inhabitated the dunes in presettlement times. Mierzwa et al. (1991) did not report coyotes, and it is not clear whether they now occur in the Grand Calumet area.

Red fox, *Vulpes vulpes* Linnaeus. Red and gray foxes are often confused, primarily because the gray fox has some reddish coloration. However, the red fox is easily identified because it is all red above with a white tail tip. Lyon (1923) reported red foxes at INDU. Whitaker et al. (1994) reported them from five different habitats at the Lakeshore, and one was found dead along the roadside in 1984-85. Four were seen from a helicopter during the 1991 Lakeshore survey for deer. Mierzwa (pers. comm.) saw a red fox dead on the road on route 312 just east of Cline Avenue on September 27, 1990, and he saw a skull found at Clark and Pine East on April 23, 1991. It is not clear how common this species is in the Grand Calumet River area.

Gray fox, *Urocyon cinereoargenteus* (Schreber). The gray fox is a resident of the woods and should occur in the forests of INDU. Lyon (1923) did not mention this species, but Texas Instruments (1975-80) recorded it on the basis of tracks. Whitaker et al. (1994) reported it from Howes Prairie on August 11, 1987, and more recently, Dan Fagre saw two in the area just east of Dune Acres. The gray fox could occur in the Grand Calumet River region in a wooded area such as in Miller's Woods, but it is less likely to occur than the red fox because of the scarcity of mature woods.

Procyonidae

Raccoon, *Procyon lotor* Linnaeus. The raccoon is obvious at the Indiana Dunes National Lakeshore and at the Indiana Dunes State Park because it is abundant in campgrounds and at

the Indiana Dunes National Lakeshore (Whitaker et al. 1994). However, its claws had been removed, probably indicating that it had been transported there from some other locality. A

introduced into 22 counties. Population estimates for the state were 900 in 1943, 1200 in 1944, and more than 2900 by 1946. A deer season was opened in 1951 when the deer populataion of the state was estimated at 5000. By 1966, deer were probably present in all counties, and they have continued to multiply, thereby becoming very abundant in recent years.

Texas Instruments (1975-1990), Krekeler (1981), and Whitaker et al. (1994) all indicated that white-tailed deer were common at the Indiana Dunes National Lakeshore. Deer were counted by aerial census in 1982, 1984, 1988, 1989, 1991, and 1992. The counts were 85, 29, 214, 349, 166, and 207 deer in those respective years. The actual populations were probably larger because it is not known what proportion of the deer were observed, and only about 75% of the Lakeshore was flown. Whitman et al. (1990) and Mierzwa (pers. comm.) saw deer tracks during their respective studies.

THE HABITATS

At the southern end of Lake Michigan, east of Chicago and east of the Gary Hammond area, lies the Indiana Dunes National Lakeshorre. The Chicago and Gary areas are some of the more highly developed and heavily polluted regions of the world. The entire area was once composed of extensive series of dunes, and classic studies of plant and animal succession were done there (Cowles 1899; Shelford 1912 a, b); however, by the middle of the 20th century, the entire area was being developed. Senator Paul Douglas was instrumental in establishing the Indiana Dunes National Lakaeshore in the area between Gary and Michigan City. Since then, much land has been acquired, many buildings have been razed, and much of the land is being converted to resemble its original condition.

Some of the larger species are not much restricted ecologically, but they could occur in any of the terrestrial habitats. Others are more restricted or are characteristic constituents of only a few of the habitats. Some of the mammals with relatively little ecological restriction, at least within an area as small and as varied as the Grand Calumet River basin, are the following:

Eastern cottontail, *Sylvilagus floridana*
Raccoon, *Procyon lotor*

White-footed mouse, *Peromyscus leucopus*
Eastern cottontail, *Sylvilagus floridanus*

Upland terrestrial shrub

There is a fair amount of upland terrestrial shrubland in the area of the Grand Calumet River. Four plots sampled by Whitaker et al. (1994) in upland terrestrial shrubland all were in highly disturbed transition stages between grassy oldfield or savanna and wooded habitats. One plot was highly productive, yielding 44 individuals of four mammal species. The nine shrews and 33 meadow voles taken there clearly reflected the former field habitat rather than the present transitional shrub stage, and these forms will undoubtedly disappear from this area as the transition towards more woody vegetation continues in this plot. A total of 14 species (including the domestic dog) was found in this habitat, but most were in low numbers. The author suspects that the lack of a well-developed community in shrubland is due to the ephemeral nature of the habitat.

Mammals likely in upland terrestrial shrubland in the Grand Calumet River basin

White-footed mouse, *Peromyscus leucopus*
Eastern cottontail, *Sylvilagus floridanus*
White-tailed deer, *Odocoileus virginiana*

Oak savanna

Oak savanna is a common habitat in the Grand Calumet River basin, and fifteen oak savanna plots were studied by Whitaker et al. (1994) at the Lakeshore. All had a canopy of nearly pure black oak, usually thin. Both the shrub and herb layer varied considerably. Some of the more abundant shrubs were blackberry, blueberry, rose, and Japanese honeysuckle. Abundant herbaceous plants were *Andropogon*,

Mixed deciduous savanna

Three plots in mixed deciduous savanna at the Lakeshore all had scattered cottonwoods with little bluestem as the principal herbaceous species (Whitaker et al. 1994). Major grasses in this habitat were old little bluestem, old witch grass and brome grass, and grape and aromatic sumac were among the more abundant shrubs. Cover was fair to good in these plots due to the grass. The white-footed mouse was the most abundant mammal, being taken at all three plots with a total of 20 individuals.

Mammal species likely in mixed deciduous savanna in the Grand Calumet River basin

White-footed mouse, *Peromyscus leucopus*
Prairie white-footed mouse, *Peromyscus maniculatus bairdii*
Prairie vole, *Microtus ochrogaster*
Eastern cottontail, *Sylvilagus floridanus*
Thirteen-lined ground squirrel, *Spermophilus tridecemlineatus*
Franklin's Ground Squirrel, *Spermophilus franklinii*
Eastern mole, *Scalopus aquaticus*

Upland forest

Upland forest, mostly oak, is widespread at INDU. (Whitaker et. al. 1994). There is relatively little mature forest in the Grand Calumet River basin, but scrubby black oaks occur on much of the forest in the Grand Calumet River basin where they grade into scrubby black oak savanna. The shrub layer at INDU was often dense and diverse in this habitat, but it often contained blueberry, *Vaccinium vacillans*. The herbaceous layer was again diverse, although often thin and depauperate, most often providing poor to fair cover. The dominant herbaceous plants there were most often Pennsylvania sedge, *Carex pennsylvanicus*, and bracken fern, *Pteridium aquilinum*. The white-footed mouse was the most abundant small mammal in upland oak forest.

Mammal species likely to be found in upland forest in the Grand Calumet River basin

White-footed mouse, *Peromyscus leucopus*
Eastern chipmunk, *Tamias striatus*
Southern flying squirrel, *Glaucomys volans*
Gray squirrel, *Sciurus carolinensis*
Fox squirrel, *Sciurus niger*
Raccoon, *Procyon lotor*
Red squirrel, *Tamiasciurus hudsonicus*
Northern short-tailed shrew, *Blarina brevicauda*
Pine vole, *Microtus pinetorum*
Gray fox, *Urocyon cinereoargenteus*

Lowland forest

Lowland forest was divided into perennial (contains water more than six months of the year) and ephemeral wet lowland forest and was the second largest habitat after upland forest in the Indiana Dunes National Lakeshore (Whitaker et al. 1994). The canopy was usually quite dense but varied in species composition. Common trees were silver or red maple followed by oak, ash, aspen, elm, and sasafrass. The shrub layer varied from sparse to dense, and it consisted mostly of seedlings of the trees mentioned above plus spicebush, *Viburnum*, *Cornus*, *Ilex*, *Rubus*, blueberries and others. Ground cover varied from fair to excellent, and species composition varied greatly between plots, with graminoid plants ferns being common. The herb and shrub layers were often clumped in hummocks. As usual in wooded habitats, the white-footed mouse was the most abundant species taken, totalling 137 in 27 plots (Whitaker et al. 1994).

Mammal species likely to be found in lowland forest in the Grand Calumet River basin

White-footed mouse, *Peromyscus leucopus*
Raccoon, *Procyon lotor*
Northern short-tailed shrew, *Blarina brevicauda*
Opossum, *Didelphis virginiana*
Gray fox, *Urocyon cinereoargenteus*
Eastern chipmunk, *Tamias striatus*

Wetlands

There is a variety of types of wetlands at the Indiana Dunes National Lakeshore, and also in the Grand Calumet River basin, marsh, aquatic shrubland, swamp, panne (depressions among the dunes), and open water.

Marsh

Marsh is prominent in the Indiana Dunes National Lakeshore and in the Grand Calumet River basin. The herb layer was most often of cattail,*Typha*; sedges, *Carex*; bullrushes, *Scirpus*; or blue joint grass, *Calamagrostis canadensis.* The canopy was absent in most areas but consisted of scattered willows or elms in a few. The shrub layer consisted of thick buttonbush, *Cephalanthus occidentalis* in most of the plots, whereas willow was present in one. Muskrats are abundant in many of the marshes, lakes, and ditches of INDU, and would be in the basin if the water was clean.

Mammal species likely to be found in Marshes in the Grand Calumet River basin

Muskrat, *Ondatra zibethicus*
White-footed mouse, *Peromyscus leucopus*

Masked shrew, *Sorex cinereus*

Mammal species likely to be found in Pannes in the Grand Calumet River basin

Meadow vole, *Microtus pennsylvanicus*
White-footed mouse, *Peromyscus leucopus*
Prairie vole, *Microtus pennsylvanicus*

Artificial habitats

Various artificial habitats or developed lands including croplands, residential and industrial areas, right-of-ways, and excavated areas are found at INDU and also in the Grand Calumet River basin.

Right-of-ways

Right-of-ways occur along roads, railroads or trails. However, they do not form a distinct habitat. Instead, they pass through and consist of some other habitat such as mature woods, dry prairie, etc. Therefore many different plants were present, including several grasses and rushes, and several other plant species (*Melilotus, Saponaria, Solidago, Clematis, Dryopteris thelypterus*, *Carex*, *Typha*) occurred as dominants in at least one plot. Because of the high variation in these plots, no list of expected species of mammals is given.

Excavated areas

Excavated areas are places where open sand exists because of human activities. There were three plots in this habitat at INDU (Whitaker et al. 1994), and there are areas with this habitat due to sand-mining in the Grand Calumet River Watershed. One of the Indiana dunes plots had been on the site of a former fly ash seepage area and another was on the site of a steel company acid spill. All three plots completely lacked canopy, and two had poor ground cover, with scattered grasses, including little bluestem, sand reed grass, and nodding wild rye *Elymus canadensis*. The third plot had excellent cover of Joe-pye weed, *Eupatorium serotinum*; bullrush, *Scirpus cyperinus*; and spikerush, *Eleocharis sp.* Only two mammals were trapped in plots in this habitat, a meadow vole and a raccoon. As would be expected, it was not a good habitat for mammals, although mammals do pass through these areas.

R

means in terms of the amount of food that can be raised on agricultural land already present, and to limit development to land already zoned as developed. “Developed” land could be returned to agricultural or natural status.

SPECIFIC RECOMMENDATIONS

1. Acquisition of land

The Grand Calumet watershed, like the Indiana Dunes before it, was made into an

reintroduced into the Grand Calumet watershed in the near future, except perhaps in the Miller Woods area.

b. River otter. The river otter can live and do surprisingly well alongside humans, and it is currently being reintroduced into Indiana. It requires extensive, relatively unpolluted aquatic habitat (ponds, lakes or rivers). If the Grand Calumet River could be cleaned up and protected, the river otter could be considered for reintroduction.

c. Bobcat. The bobcat was thought to be nearly extirpated in Indiana, but it is showing up in various counties. There is some evidence of its occurrence at the Indiana Dunes National Lakeshore. This species can live in fairly close proxity to humans, but it does need rather extensive natural woodland habitat because it moves about considerably. The recommendation that woodland habitat be preserved or created would favor this species. However, I would not recommend that reintroduction of this species be attempted in the Grand Calumet basin at this time because of the lack of woodland.

Management recommendations: Once the Grand Calumet River is cleaned up, some consideration could be given to reintroducing the river otter there, provided that adequate protected habitat is available for them. Consideration should be given to introducing the porcupine at the Indiana Dunes National Lakeshore but not in the Grand Calumet River basin until such time as adequate woodland might exist there.

6. Areas of special interest

There are several areas of ecological interest in the Grand Calumet River Basin such as Roxanna marsh, DuPont wetlands, and the Grand Calumet Lakeshore Lagoons because of their high quality habitat for semi-aquatic mammals. Additional areas of like quality, and also to reduce fragmentation, as many intervening areas as possible should be obtained and purchased or otherwise protected.

Management recommendations: In line with recommendation #3, attempts should be made to obtain such properties, or to reach management agreements for them.

LITERATURE CITED

Baker, R. H. 1983. Michigan mammals. Michigan State University Press, Lansing, Michigan, USA.

Brennan, G. A. 1923. The wonders of the dunes. The Bobbs-Merrill Co., Indianapolis, Indiana, USA.

Brooks, D. M. 1959. Fur animals of Indiana. Pittman-Robertson Bulletin No. 4., Indiana Department of Conservation, Division of Fish and Game. Indianapolis, Indiana, USA.

Cowles, H. C. 1899. The ecological relations of the vegetation on the sand dunes of Lake Michigan. Botanical Gazette **27**:95-117, 167-202, 281-308, 361-391.

Dice, L. R. 1928. The least weasel in Indiana. Journal of Mammalogy. **9**:63.

Evermann, B. W. and A. W. Butler. 1894. Preliminary list of Indiana mammals. Proceedings of the Indiana Academy of Science **3**:124-139.

Ford, S. D. 1977. Range, distribution and habitat of the western harvest mouse, *Reithrodontomys megalotis*, in Indiana. American Midland Naturalist 98:422-432.

Hahn, W. L. 1909. The mammals of Indiana. Pages 417-663. A descriptive catalogue of the mammals occurring in Indiana in recent times. 33rd Annual Report Department of Geology and Natural Resources of Indiana. Indiana, U.S.A.

Hoffmeister, D. F. 1989. Mammals of Illinois. University of Illinois Press, Urbana, Illinois, USA.

Krekeler, C .H. 1981. Ecosystem study of the Indiana Dunes National Lakeshore. The Biota of Indiana Dunes National Lakeshore.Vol 2. Mimeograph **1-346**.

Kurta, A. 1982. A Review of Michigan bats: Seasonal and geographic distribution. Michigan Academician **14**:295-312.

Kurta, A. D. King, J. A. Teramino, J. M. Stribley, and K. J. Williams. 1993. Summer roosts of the endangered Indiana bat (*Myotis sodalis*) on the northern edge of its range. American Midland Naturalist **129**:132-138.

Lindsey, A. A., D. V. Schmeltz and S. A. Nichols. 1969. Natural areas in Indiana and their preservation. Department of Biological Sciences, Purdue University, Lafayette, Indiana, USA.

Long, C. A. 1974. Environmental status of the Lake Michigan region. Vol. 15. Mammals of the Lake Michigan drainage basin. Volume 15. Argonne National Laboratory, Argonne, Illinois, USA.

Lyon, M. W. Jr. 1923. Notes on the mammals of the dune region of Porter County, Indiana. Proceedings of the Indiana Academy of Science **31**:209-221.

Lyon, M. W. Jr. 1936. Mammals of Indiana. American Midland Naturalist **17**:1-384.

Mierzwa, K. S., S. D. Culverson, K. S. King, and C. Ross. 1991. Illinois-Indiana regional airport study: Biotic communities Technical Paper No.7, Appendix E, Volume II. TAMS Consultants, Inc. Chicago, Illinois, USA.

Mumford, R. E., and J. O. Whitaker, Jr. 1982. Mammals of Indiana. Indiana University Press, Bloomington, Indiana, USA. 537 p.

Rand, A. L. and A. S. Rand. 1951. Mammal bones from dunes south of Lake Michigan. American Midland Naturalist **46**:649-659.

Sanborn, C. C. 1925. Mammals of the Chicago area. Field Museum of Natural History. Zoology Leaflet 8. 23 p.

Shelford, V. E. 1912a. Ecological Succession. IV, Vegetation and the control of land animal communities. Biological Bulletin **23**:59-99.

Shelford, V. E. 1912b. Ecological succession. V, Aspects of physiological classification. Biological Bulletin **23**:331-370.

Texas Instruments Inc. 1975-1980. Six annual reports, for the years 1975 through 1980. Bailly Nuclear-1 Site. Prepared for Northern Indiana Public Service Co.

Van Zyll de Jong, C. G. 1979. Distribution and systematic relationships of long-eared *Myotis* in western Canada. Canadian Journal of Zoology **57**:987-994.

Waldren, L. 1983. The Indiana Dunes. The Eastern Acorn Press. New York, New York. USA.

Whitaker, J. O. Jr. 1967. Habitat and reproduction of some of the small mammals of Vigo County, Indiana, with a list of mammals known to occur there. Occasional Papers C.C. Adams Center for Ecological Studies 16. Western Michigan University. Kalamazoo, Michigan, USA. 24p.

Whitaker, J. O. Jr., J. Gibble and E. Kjellmark. 1994. Mammals of Indiana Dunes National Lakeshore. Scientific Monograph. NPS/NRINDU/NRSM-94-24. United States Department of the Interior. National Park Service.

Whitaker, J. O. Jr., and G. R. Sly. 1970. First record of *Reithrodontomys megalotis* in Indiana. Journal of Mammalogy **51**:381.

Attachment 1. Likely mammal species of Calumet basin, presettlement, present, and likely origin (native, reintroduced native, introduced exotic, extirpated)

Native mammals, probably present in presettlement times, but gone now

	PRESETTLEMENT	NOW	STATUS
RODENTIA			
Erethizon dorsatum, American porcupine	yes	no	native
CARNIVORA			
Canis lupus, gray wolf	yes	no	native
Ursus americanus, black bear	yes	no	native
Martes pennanti, fisher	yes	no	native
Lutra canadensis, river otter	yes	no	native
Felis concolor, mountain lion	yes	no	native
Felis lynx			

GRAND CALUMET LAGOONS

Paul M. Stewart and Jason T. Butcher

Lake Michigan Ecological Research Station
Biological Resources Division

Despite gains in control and remediation of conventional pollutants (e.g. nutrients, organic wastes), we still lack knowledge of the eff

The East and Middle Lagoons are connected by a wide channel under the Lake Street bridge. The shoreline of the Middle Lagoon (9.9 hectares) is shared by the city of Gary, homeowners, and the National Park Service. This includes the Miller Woods Unit of INDU with dunes and scattered vegetation to the north and black oak savanna to the south of the Lagoon.

The West Lagoon (14.8 hectares) is connected to the Middle Lagoon by a shallow stream that normally flows to the west. The creek was flowing to the east once during the summer of 1994 (P.M. Stewart, personal observation). This occurred after a storm event with subsequent elevated water levels. A fence along the shore separates the eastern section (approximately one

Algae and periphyton were also collected for identification by site. Data from these samples will be presented and discussed in the final report for this project to the National Park Service.

Water quality

Water quality data at twelve sites were collected four times during the summer of 1994 and three times during the summer of 1996. Only the 1994 data will be discussed here. Variables measured include dissolved oxygen, water temperature, pH, specific conductance, total hardness, total alkalinity, and concentrations of sulfate, chloride, ammonia, nitrite, nitrate, reactive phosphate, and total phosphate.

Dissolved oxygen, pH, specific conductance, and temperature were measured in the field with calibrated meters. Water samples were collected at each site in acid-washed polyethylene bottles. The samples were placed on ice for transport to the laboratory where the remaining variables were measured. All variables were analyzed within a suitable time frame following standard methods (APHA 1992).

According to an independent t-test, alkalinity, total hardness, sulfate, ammonia, and nitrate were significantly higher in the West Lagoon than in the Middle Lagoon. The same variables were significantly higher in the West Pond than in the East Pond ($p<0.001$). Total phosphate was not significantly different between the water bodies. Chloride was significantly higher in the Middle Lagoon and the East Pond than in the West Lagoon and the West Pond ($p<0.001$). Ammonia concentrations were correlated with location in the Lagoons ($r=0.860$, $p<0.001$) and were highest at the western sites and lowest in the eastern sites.

High concentrations of ammonia were found throughout the Lagoons and levels were especially high in the West Lagoon, where mean concentration for site WL5 was 2.07 mg/l in 1994. The high levels of ammonia in the West Lagoon and the West Pond, combined with the high pH of these systems, may contribute to the production of high concentrations of undissociated ammonia hydroxide. This form of ammonia [$NH_4OH + NH_3$(dissolved)] is toxic to many aquatic organisms. According to Trussell (1972), the toxic fraction of ammonia is 15% at a pH of 8.5 and a temperature of 25 ^{0}C. Ammonia is removed from the water by plants thus decreasing CO_2 and increasing pH. If future conditions cause an increase in pH to 9.0, the percentage of the more toxic form of ammonia would exceed 35%, leading to elevated toxicity.

In summary, many of the water quality parameters tested in this study, especially ammonia, total hardness, and sulfate, were significantly higher in the West Lagoon and West Pond. These water bodies were most closely associated with the industrial area. Overall, the variables indicated poorer water quality in the impacted areas, and in some areas the water may be toxic to aquatic life.

Plant communities

Aquatic plant species in the Grand Calumet Lagoons were enumerated with an abundance scale (observed=1, rare=2, rare/common=3, common=4, very common=5, abundant=6) (Table 1). Aquatic plant species were also assigned a growth habit for further

evaluation (floating, emergent, submergent). Using inductively coupled plasma techniques (ICP), aquatic vascular tissues (roots and shoots) were analyzed for heavy metal concentrations.

The aquatic vascular plant taxa found at each site in the Grand Calumet Lagoons are presented along with their abundance scale values (Table 1). Site ML3 had the greatest number of aquatic plant species with all growth habits represented. Site ML2 had only 11 species with no emergent representatives. The number of aquatic plant taxa was significantly reduced at the WL3-5 sites (Figure 2). The Middle Lagoon had the greatest number of aquatic plant taxa, but this number was only significantly different from the WL 3-5 sites. Floating plant species were absent from the West Lagoon at the time of our sampling (Table 2), yet these had been observed

Figure 2. The mean number of aquatic vascular plant taxa per site in the Middle Lagoon (ML1-3), West Lagoon (WL1-2) and (WL3-5), East Pond (EP), and West Pond (WP).

Table 1. Aquatic plant species collected from the Grand Calumet Lagoons and ponds. Numbers refer to abundance scale (observed=1, rare=2, rare/common=3, common=4, very common=5, abundant=6).

on previous trips to the Lagoons. An exotic plant species, *Myrophyllum spicatum*, was found throughout the Middle Lagoon and at several sites in the West Lagoon earlier in the season. Submergent plant species were reduced at the We

Table 2. Number of aquatic plant taxa that occupy the growth habits of submergent, floating, and emergent at each site.

Site	Submergent	Floating	Emergent	Total
WL5			7	7
WL4			10	10
WL3			7	7
WL2	4		10	14
WL1	5		5	10
ML3	9	2	9	20
ML2	8	1		9
ML1	7	2	4	13
WP	6	1	5	12
EP	6	2	6	14

Cluster analysis (an assessment of similarity) based on the coefficient of similarity was done to examine relationships among sites based on aquatic plant distribution (Pearson and Pinkham 1992, Gonzales et al. 1993). Results showed two main groupings (Figure 3). The Middle Lagoon sites and the East and West Ponds formed one such cluster, and the five West Lagoon sites formed another cluster. This suggests that there are differences in aquatic plant distribution that can be attributed to landfill proximity.

Representative aquatic vascular plants were collected from several sites and analyzed for heavy metal content (Table 3). Several heavy metals, including zinc, iron, and aluminum, were at elevated concentrations in some of the tissue samples analyzed. *Scirpus americanus* collected from the pond sites had higher tissue aluminum concentrations in the plant roots than in the shoots (Figure 4). The same species collected from the Lagoons showed the opposite trend. Compared to values found in the literature, aquatic plant tissue metal concentrations at several

Table 3. Selected plant species and heavy metal root and shoot concentrations (mg/kg) for

Figure 4.

Aluminum concentrations in the tissue (shoot and root) of *Scirpus americanus* collected from sites WL4, ML3, WP, and EP.

Macroinvertebrate communities were sampled in the Grand Calumet Lagoons using both Hester-Dendy multiplate samplers (Table 5) and sweep nets of the vegetation (Table 6). The Hester-Dendy plates were left in the field for a month (less than the recommended time of six weeks), and the results were inconclusive. The macroinvertebrate data collected by sweep nets of the vegetation showed more definite results.

Both Shannon-Weaver and Simpson Diversity indices of the sweep net macroinvertebrate data showed significantly ($p<0.05$) lower species diversity at the WL3-5 sites than at the other sites (Figure 5). The Hester-Dendy plate data from the Grand Calumet Lagoons show little discrimination between sites. This lack of difference must be viewed with caution because organisms were only identified to the family level. Genus and species level taxonomic identifications allow for finer discrimination. In addition, Hester-Dendy plates are not normally used in lentic systems (Merritt and Cummins 1984). The sweep net samples show clear differences between community dominants found at ML1-3, WL1-2, and WL3-5 sites (Figure 6). Talitrids were dominant at WL3-5 (more than 95% of the organisms counted at WL5), were fewest at WL1-2, and were present in moderate numbers at ML1-3. Chironomids were most abundant in the center of the Lagoons at sites ML2 and WL2. They comprised a small percentage of the community at ML1 and WL3-5. Coenagrionids were present at all of the Middle Lagoon sites and at WL2, but they were absent from all other sites. Overall, the macroinvertebrate communities at WL3-5 show a negative response to environmental impact.

Figure 5. Macroinvertebrate diversity (Simpson and Shannon-Weaver) from sweep net samples collected from the Grand Calumet Lagoons.

Table 5. List of macroinvertebrate species collected with Hester-Dendy multiplate samplers.

Table 6. List of macroinvertebrate species collected by sweep netting of vegetation.

abundant in the East Pond vegetation than in the West Pond. There were no Asselids found in the West Pond, and no Coengrionids found in the East Pond. These differences in the macroinvertebrate community show a marked response to landfill proximity. The Hester-Dendy data for the ponds suffers from the same weaknesses as the Lagoons.

A cluster analysis of the taxa (families) at each site shows four major groups (Figure 8). WL3-5 formed one cluster, the East Pond sites a second, and the West Pond sites a third. The fourth cluster included samples collected from both the Middle Lagoon and WL1-2. This shows, for the macroinvertebrate community living among the aquatic vegetation, that there were differences between communities found at the far-western locations of the West Lagoon and those of the eastern section of the West Lagoon. The latter were more similar to those communities collected from the Middle Lagoon.

Jokinen (1994) found certain gill-breathing mollusks (prosobranchs) in the East Pond (furthest and over a dune ridge from the slag landfill) but not in the West Pond when she investigated the mollusks of the Indiana Dunes National Lakeshore. These include *Valvata tricarinata* and *Amnicola limosa*. In addition, she found a member of the Lymnaeidae family (*Fossaria exigua*) in the West Pond but not in the East Pond. Three additional species of mollusks, *Pisidium casertanum*, *Physa gyrina*, and *Gyralus parvus*, were found in both ponds.

Macroinvertebrate communities based on family level taxonomic identification clearly demonstrate a difference in community composition relative to their proximity to the industrial landfill. Further taxonomic identification to the genus and species level would undoubtedly strengthen these findings. Sites WL3-5 formed a separate cluster and had fewer taxa and lower species diversity than other sites sampled. These data support the conclusion that the macroinvertebrate community responded to impacts from the industrial landfill and were related to landfill proximity.

Fish communities and metrics

Fish communities were sampled at all twelve sites. Sampling was performed by electroshocking for 100 m in each of the Middle and West Lagoon sites. The ponds were sampled by repeated seines at each of the sites. Fish were counted and identified in the field. Community biometrics were performed for each site. At several sites carp (*Cyprinus carpio*) were collected, placed on ice, and brought to the laboratory for whole tissue analysis of PAH concentrations.

Seventeen species of fish were collected in the sampling effort (Table 7). The East and West Ponds had two species each, and both were heavily dominated by pumpkinseed sunfish (*Lepomis gibbosus*) (more than 320 collected). The second species present was the grass pickerel (*Esox americanus*) in the East Pond and the lake chubsucker (*Erimyzon sucetta*) in the West Pond. Only one lake chubsucker was collected from the West Pond.

Figure 7. Domi


WL3
WL4

Notable in the Grand Calumet Lagoons was the collection of the Iowa darter (*Etheostoma exile*) at several sites, including WL3 and WL4, which had not been previously recorded in northwestern Indiana (Tom Simon, U.S. EPA, personal comm

are formed. The East and West Pond form one cluster and are dominated by pumpkinseed sunfish and one other species. The second cluster was made up of Middle Lagoon sites, and the third cluster included all of the sites from the West Lagoon. A small stream choked
with cattails forms a barrier b8 eGs form

11. % Simple lithophils	**16.6 - 33.9 %**
12. % DELT anomalies	**0.1 - 1.3 %**

Table 10. IBI scores and assessment for fish comm

Table 11. Whole fish tissue PAH concentrations from sites in the Grand Calumet Lagoons.

Analyte (µg/kg)	WL3-A	WL3-B	WL5-A	WL5-B
Napthalene	29.5	29.6	212.1	145.9
Acenaphthene	80.9	34.3	450.9	446.6
Fluorene	14.0	11.1	138.6	105.5
Fluoranthene	11.6	36.5	84.5	18.5
Total PAHs	257.4	282.4	1413.4	807.3

Sediment chemistry

Pyrene	BDL	1,694

that subsequently caused problems with the analytical chemistry. As in our sampling, the sediments between WL4 and WL5 were loose, black, oily, and flocculent, with a strong hydrocarbon odor.

Limited evidence (few samples, but high levels of PAHs) suggests that a major contaminant problem exists in the sediments of the Grand Calumet Lagoons. The PVC pipe used for sediment sampling became covered in a black, tarry coating between sites WL4 and WL5. WL5 has high PAH concentrations, and they are much higher than levels exhibiting major biological effects.

Aquatic toxicity testing

Aquatic toxicity testing was performed on the water and sediments from the West Pond. Despite apparent contamination, neither surface water nor sediments appeared to be toxic. This lack of toxicity could be due to modifying or interfering factors in the tests, such as hardness. It also may have been due to procedural problems which must be addressed.

There is no strong evidence that surface waters in the Lagoons were toxic, so toxicity testing was focussed on the Lagoon sediments. All fathead minnows died within twelve hours when exposed to the sediments from WL5. These sediments (WL5) were not tested further due to their extremely contaminated nature.

Whole sediments from WL4 affected survival and growth rates of amphipods, *Hyalella*, and sediment eluriate was toxic to *Ceriodaphnia*. This suggests that sediments from WL4 may also be toxic to aquatic organisms in the Grand Calumet Lagoons. Sediments at the other sites in the Grand Calumet Lagoons are contaminated with heavy metals. Despite this contamination, neither whole sediments nor eluriate from these sites decreased survival or growth in fathead larvae. Sediments from WL1 and WL3 appeared to affect growth in amphipods, while sediments and eluriate from WL2 and WL3 appeared to affect survival and reproduction in *Ceriodaphnia*. This evidence suggests that sediments from WL2 and WL3 have negative effects on aquatic organisms.

The results of these toxicity assays should be considered preliminary. In some cases, no toxicity was observed from sediments that had previously shown elevated concentrations of contaminants. A resampling effort with further aquatic toxicity testing is necessary to elucidate fully the extent of sediment toxicity in the Grand Calumet Lagoons.

CONCLUSIONS

An obviously impaired aquatic community exists within and surrounding the Grand Calumet Lagoons and ponds. The ponds and surrounding areas include wetlands, dune and swale habitat, and a savanna community. This habitat type is considered to be among the most endangered ecosystems on the planet. Rare habitat and federally endangered and threatened species exist within the area.

This option would allow continued degradation of the park and its biota due to extensive contamination that already exists. Since the Lagoons are the headwaters of the Grand Calumet River, this would allow for continued contamination of the Grand Calumet River west of the culvert. This would leave surface contaminants containing more than 12% PAH in the area, and aquatic toxicity testing has shown the sediments in the far-western regions to be extremely toxic. Further contaminant uptake in the water column would add to downstream discharges. Positive aspects of the option would include avoiding sediment resuspension during dredging and the associated turbidity problems. Existing ecosystems would not be disturbed, so there would be little risk of removing some species from the area. It is the least costly option, at least in the short term.

Option 2 - Remove contaminated material from the area.

This option comes in three related forms. One option is removing hot spots which must first be more clearly identified. The second option is removing a substantial portion of the contamination, and the third option is to remove all contaminants in the area. There is little difference between removing the hot spots and removing a portion of the contamination, and both options would leave a great deal of contamination behind. The more contaminants and sediments are removed, the more this would cost.

Removing all the contaminants in the area would require the removal of a great deal of sediment and fill. This includes the slag piles, hazardous waste dump number 2 , metal burning sites, and the sediments from the Grand Calumet Lagoons. This removal would necessitate the construction and maintenance of a large, lined storage facility. Complete removal of contamination is expensive and extremely difficult. Removal of contaminants past a certain point to the east would destroy the resource that we are trying to protect. Building a landfill on park property would be an extremely contentious issue.

The benefits of this option include the reduced contaminant impact on the park and biota including unique and fragile habitats of oak savanna, dune and swale, and wetlands. The proper disposal of a large volume of material should be addressed including potential locations of a fill site and proper site construction to prevent future recontamination.

Option 3 - Cap the contaminated areas (the landfill), and form a physical containment barrier between more contaminated and less contaminated sediments.

No barrier system will last forever because it assumes complete separation between the upper and lower aquifers. Numerous wells throughout the region need to be studied and perhaps removed because they may have aided in contamination of the lower aquifer. Groundwater monitoring should be instituted. If necessary, pumping and remediation efforts should be initiated to clean up the aquifer and to keep contamination from spreading. Capping the contaminated sediment and then adding clean sediment to the system would potentially benefit the aquatic ecosystem, but further contamination of ground water is also possible. Existing

conditions would be dramatically altered. It may be necessary to establish a groundwater divide that separates moving landfill contaminants in the Lagoons from Lake Michigan.

LITERATURE CITED

APHA. 1992. Standard methods for the examination of water and wastewater, 18th edition. Washington, D.C., USA.

Baumann, P.C., M.J. Mac, S.B. Smith, and J.C. Harshbarger. 1991. Tumor frequencies in Walleye (*Stizostedion vitreum*) and Brown bullhead (*Ictalurus nebulosus*) and sediment contaminants in tributaries of the Laurentian Great Lakes. Canadian Journal of Fisheries and Aquatic Sciences **48**:1804-1810.

Black, J.J., T.F. Hart, Jr. and E. Evans. 1981. HPLC studies of PAH pollution in a Michigan trout stream. Pages 343-355 in M. Cooke and A.J. Dennis, editors. Chemical analysis and biological fate:polynuclear aromatic hydrocarbons. Fifth international symposium. Batelle Press, Columbus, Ohio, USA.

Bosserman, R.W. 1985. Distribution of heavy metals in aquatic macrophytes from Okefenokee Swamp. Heavy metals in water organisms. Pages 31-39 in Symposia Biologica Hungarica Volume 29. Akademiai Kiado, Budapest, Hungary.

Clark, J.U. and A.S. Jarvis. 1990. Regulatory evaluation of petroleum hydrocarbons in dredged material. Miscellaneous paper EL-90-11. U.S. Army Corps of Engineers Waterways Experimental Station, Vicksburg, Mississippi, USA.

Cowgill, U.M. 1973. The determination of all detectable elements in the aquatic plants of Linsley Pond and Cedar Lake (North Bradford, Connecticut) by x-ray emission and optical emission spectroscopy. Applied Spectroscopy **17**:5-9.

Gatz, D.F., V.C. Bowersox, and J. Su. 1989. Lead and cadmium loadings to the Great Lakes from precipitation. Journal of Great Lakes Research **15**:246-264.

Gonzales, D.A., J.G. Pearson, and C.F.A. Pinkham. 1993. User's manual BIOSIM1, Beta version 1.0. A program that applies the coefficient of biotic similarity to complex data matrices. EPA 600-R-93-219. U.S. Environmental Protection Agency, Environmental Monitoring and Support Laboratory, Las Vegas, Nevada, USA.

Hardy, M.A. 1984. Chemical and biological quality of streams at the Indiana Dunes National Lakeshore, Indiana, 1978-1980. U.S. Geological Survey, Water Resources Investigations 83-2408. Indianapolis, Indiana, USA.

Hilsenhoff, W.L. 1988. Rapid field assessment of organic pollution with a family-level biotic index. Journal of the North American Benthological Society **7**:65-68.

Indiana State Department of Health. 1997. Indiana Fish Consumption Advisory. Environmental Epidemiology Section, Indiana State Department of Health, Indianapolis, Indiana, USA.

Jokinen, E. 1994. The freshwater mollusks of the inland waters of the Indiana Dunes

Simon, T.P. 1991. Development of an index of biotic integrity expectations for the ecoregions of Indiana, I. Central Corn Belt. U.S. Environmental Protection Agency, Region 5, Environmental Sciences Division, Monitoring and Quality Assurance Branch: Ambient Monitoring Section, EPA 905-9-91-025. Chicago, Illinois, USA.

Simon, T.P. and P.M. Stewart. In review. Development of fish community reference conditions for diurnal, palustrine wetlands: an emphasis on assessment of non-point source landfill effects. Aquatic Ecosystem Health and Restoration.

Smith, R.A., R.B. Alexander, and M.G. Wolman. 1987. Water-quality trends in the nation's rivers. Science **235**:1607-1615.

St. Cyr, L. and P.G.C. Campbell. 1993. Trace metals in submerged plants of the St. Lawrence River. Canadian Journal of Botany **72**:429-439.

Steffeck, D.W. date unknown. A survey for contaminants near the Midco I, Midco II, and Ninth Avenue Dump hazardous waste sites in Gary, Lake County, Indiana. U.S. Fish and Wildlife Service, Bloomington, Indiana, USA.

Stewart, P.M. and D.J. Robertson. 1992. Aquatic organisms as indicators of water quality in suburban streams of the lower Delaware River Region, USA. Journal of the Pennsylvania Academy of Science **66**:135-141.

Trussell, R.P. 190 llsnae Ppece t of nnitnitzd pamuoia muoia

Shira Hammann, Meredith E. Becker, and Richard L. Whitman

Lake Michigan Ecological Research Station
Biological Resources Division
U.S. Geological Survey
1100 N. Mineral Springs Rd.
Porter, Indiana 46304

I

large monocultures that have choked out many other, more niche-specific species. Processes required for the creation and maintenance of local ecosystems, such as periodic fires and migration, have been suppressed. This interruption has allowed all communities to become overgrown, and some community types have been lost to most of the region. Due to habitat destruction and fragmentation, the gene pools of many niche-specific native species have become dangerously small.

Despite all of the degradation that has taken place, a few high quality natural areas remain. These are small fragments of the former landscape, often isolated both from each other and from the natural processes that formed them, but they have managed to persist largely intact.

The Nature Conservancy, using Natural Heritage Network data, has identified the Greater Calumet Wetlands Site as an area that supports globally and state significant biodiversity. There are 18 natural community types extant within the site. Within these, are found more than 700 species of native plants, of which 85 are globally or state significant; more than 200 species of birds, including 18 species confirmed to nest in the area that are globally or state significant; and 21 species of reptiles and amphibians, eight of which are state significant.

Several federally listed species are known to occur in the study area: Pitcher's thistle (*Cirsium pitcheri*), the Karner blue butterfly (*Lycaeides melissa samuelis*), and the Peregrine falcon (*Falco peregrinus*). Habitat appropriate for Indiana bat (*Myotis sodalis*) is also present, although it has not been demonstrated that the species is there. Numerous state listed species are found throughout the high quality habitats of the study area.

The highest quality natural areas of the study site are Miller Woods, the East and Middle Lagoons of the Grand Calumet Lagoons, Clark and Pine er Wsss, thsruPo wiD(

disturbance. There has been some improvement, with fish communities rebounding during the last decade. Abundance and diversity have improved, and these communities are primarily composed of invasive, tolerant species

RESTORATION PHILOSOPHY AND S

if these are sufficiently cleaned and protected to support it. American porcupine (*Erethizon dorsatum*), and perhaps smooth green snake (*Opheodrys vernalis*) could be reintroduced into the Miller Woods area if enough good habitat can be created.

Sediment clean-up has great potential to benefit the Grand Calumet ecosystem by

be occur along with sediment clean-up, but they will be ongoing projects and cannot be performed in any meaningful and lasting way if they are considered only as adjunct to sediment clean-up.

Continuous monitoring of all aspects of the system will be essential to chart restoration progress. Management and partnerships among involved groups will be imperative for making restoration an ongoing, rather than a sporadic and poorly planned endeavor. Ecological considerations will need to be incorporated in a meaningful way at the beginning and in all steps of the land use planning process. Many sites are indirectly affected by the river, including terrestrial systems, and management for these areas must be considered as a step in the entire system restoration. Further, any restoration efforts must be supported by pollution abatement in order to prevent re-contamination of newly restored areas. In some areas it may not be possible to return to historic conditions, but by decreasing pollution impacts and actively preserving and managing lands, progress toward an integrated system of natural areas can be made.

REACH-BY-REACH RECOMMENDATIONS

The Army Corps of Engineers has divided the river into ten reaches that will correspond to discrete dredging projects. The division between reaches is arbitrary from a habitat standpoint, and reach-by-reach recommendations cannot stand alone without incorporation of the ideas detailed above for the entire study system. Recommendations for activities at specific locations can, however, be broken into reach-unit groupings. These, along with brief descriptions of the habitat found in each reach, are described below.

Lagoons reach

The Lagoons area contains two large tracts with high value as natural areas: The Grand Calumet Lagoons and Miller Woods, which border the Lagoons to the north and south.

Grand Calumet Lagoons

The Grand Calumet Lagoons are a series of lagoons located at the eastern end of the Grand Calumet River where the mouth of the river once opened to Lake Michigan. The Lagoons are now at the River's east end. The area immediately to the south of the Lagoons consists of mesic sand savanna, with interspersed marsh and pond communities in swales. The area to the north borders Lake Michigan and includes unusual habitats, such as pannes. The easternmost lagoon provides habitat for the most diverse array of macroinvertebrates in the study area and for a wide variety of submerged aquatic vascular plant species. The ponds that are closely associated with the Lagoons are also valuable as rare remnants of the once-common panne-type community.

The Lagoons are divided into three parts: The West Lagoon, the Middle Lagoon, and the East Lagoon. The West and Middle Lagoon are separated by a shallow stream that usually flows west. A wide channel separates the Middle Lagoon from the East Lagoon. The Middle Lagoon is thought to be the least contaminated of the three areas. The West Lagoon is highly contaminated, and it is currently under enforcement action under the Resource Conservation and

could include the reintroduction of American porcupine (*Erethizon dorsatum*), and perhaps eventually the river otter (*Lutra canadensis*), into the Miller Woods area. Restoration and management should be carried out in accordance with the advice of local experts and land managers.

In the South Shore portion of the Miller Woods tract, there is extensive illegal dumping of tires and other items. It is recommended that Miller Woods be patrolled and secured from intruders at its southern end.

USX reach

The USX reach of the Grand Calumet River is highly degraded for most of its length. The highest quality natural areas in the reach, both located at the far western end, are the Bongi site and a small portion of the Clark and Pine site that is located mostly in the Gary Sanitary District reach. The Bongi site contains two borrow pits located to the north of the river. The north pit, further from the river, is more contaminated than the south pit perhaps due to fly ash runoff.

Plans for restoration of the USX reach of the River are currently being formulated as part of a RCRA enforcement action against USX Corporation for this area. The plans are currently on hold while the Indiana Department of Environmental Management decides whether or not to issue a water discharge permit for the river dewatering and discharge portion of the USX compliance plan. The plans, which will be paid for by USX Corporation, include dredging within that reach to remove contaminants. Other plans may include the purchase and restoration of natural areas and the control of exotic species.

In addition to these efforts, the U.S. Army Corps of Engineers plans to add deflector logs and submerged gravel weirs in this reach. Deflector logs will help to stabilize the shoreline before and after dredging. Stable shoreline and gravel weirs will provide fish habitat after dredging.

Several concerns remain. Dredging around the sensitive Bongi property at the western end of this reach may cause the banks to erode in this area, thereby destroying portions of this valuable property. Bank sloughing may also cause contamination of the Bongi ponds with sediment-laden waters from the River. The banks here should be stabilized with native vegetation to minimize this impact. The Georgia Pacific Lagoon, located to the south of the river, opposite the Bongi property, is also in danger of contamination due to sediment migration. It has been suggested that a levy be built to protect this area, with tie gates to allow for fish movement in and out of the Lagoon, but these provisions are not included in the current plans for restoration.

Recommendations for the Clark and Pine property are discussed below in the section on the Gary Sanitary District reach.

Possibilities for restoration in this reach of the River have been more extensively studied during the RCRA process than by these authors, so recommendations for this reach are limited. It is highly recommended that any activities in this reach be carefully coordinated between different management agencies. Managers of nearby lands, such as INDU, should be notified in advance of any release of biocontrol organisms, or other activities likely to impact lands beyond the immediate study area.

establish sedge meadow hydrology and a conservative flora and fauna after the channel has closed. The drastic alteration of the local landscape suggested here would carry with it a high risk of invasion by exotics. Careful planning, and long-term continuous monitoring and exotic species elimination would be essential to achieve the desired results.

Alternatively, areas of the bank could be dredged at a shallower level to create backwaters in the river channel proper. These, like the dredged side channels, could potentially serve as turtle habitat, but they also would need to be actively managed to prevent exotic infestation.

In-stream restoration alternatives include the creation of fish habitat in the river. Lunker boxes are structures built by embedding the ends of log platforms in the river banks and allowing them to backfill with rock. These could provide cover and shallow habitat by creating an artificial undercut bank. This should result in an increase in salmonids, and, because the boxes would be built in the open stream rather than in isolated backwaters and ponds, amphibian species should not be affected by an increase in salmonids. Altering river flow to establish large pool/short riffle sequences would also create new habitat within the river channel. These would help fish and macroinvertebrate communities by encouraging natural aeration. Suckers could spawn in the riffles.

The top priorities for restoration in this reach are to pollution-abate, protect, and manage the Clark and Pine site, and to create buffer areas and migration corridors to protect and connect the Clark and Pine and DuPont natural areas. Possible staging areas include the access road at the end of the Gary Regional airport and either side of the River at the landfill.

DuPont reach

quality habitats, but functional limitations due to cultural and land use would limit how successful the corridor could be.

Unfortunately, contaminants are a major problem in the DuPont reach of the Grand Calumet River because concentrations tend to increase further downstream. Fish are greatly affected, and oil-soaked birds are often observed in the area. Contamination affects shoreline plant habitat as well, but it is unknown how deep the contamination reaches into the sediment or how much contamination is present in areas that are not immediately adjacent to the channel. A large stand of cottonwood trees most likely indicates the disposal site of previous dredge spoil. Another major problem in this area is nutrient loading. Nuisance algal blooms have resulted due to deterioration in water quality. Nutrients, including nitrogen and phosphorus, likely enter from inputs upstream, and the resulting blooms do not benefit the ecosystem.

U.S.S. Lead currently has a cleanup program for this area out for public comment. One portion of the cleanup is likely to focus on the channel leading up to the U.S.S. Lead property from the River. Planners of this cleanup should take care to stabilize banks and prevent migration of contaminants into adjacent, highly restorable savanna lands. The U.S.S. Lead property contains extensive marshy habitat which is overgrown with submergent plant species.

deposited in shoreline soils. For this reason, water quality should be monitored intensively and deemed safe before any decisive plans for riverbank flooding are made or implemented.

Removal of exotics will be important not only along the dredge channel, but also within the high quality natural areas. In many locations of the DuPont Property, micro-habitat areas exist that are small enough, and exotic populations are limited enough, that manual control could realistically clear exotics from the sites. Ongoing plans for such removal activities are highly recommended.

A highly aggressive approach should be taken in dredging the channel here, in order to clean this particularly polluted reach, but much care should be taken to protect sensitive habitats along the banks. Further study should determine the extent of contamination in areas associated with the river by hydrology, but too far away from the channel to be included in the dredge process.

East Chicago Sanitary District reach

This section of the River contains several small remnant areas of dune and swale, as well as several very polluted areas. An area of upland meadow is present to the south of the river, between the Indiana Harbor Canal and the Roxanna Substation, which is located to the east of Indianapolis Boulevard on the north side of the river.

The upland meadow at the Roxanna Substation shows great promise as a site for restoration activities because of its connection with the high quality DuPont reach; restoration efforts could upgrade the site from a degraded remnant tract into a productive habitat.

Roxanna Marsh reach

The Roxanna Marsh reach of the River contains an open, shallow pond that was once a stopover area for shorebirds migrating between the Arctic Tundra and South America. The water level has risen during the past ten years, making the area unsuitable as a staging area for these birds. The area is now a large mudflat with very little vegetation growing on it. The area also contains several sites, dominated by a flora of large cottonwood trees, that were previously filled with dredged materials. These are each approximately 30-40 acres in size. One is located neat the Harbison-Walker property and the other is near the U.S.S. Lead. These may support bat populations, which are rare in the study area. Dredging should increase the macroinvertebrate community which will in turn provide an insect food source to bats inhabiting the area.

One option for restoration of Roxanna Pond is to manage water and vegetation levels in the spring and fall so that the area can again be used as a landing and foraging area for migrating shorebirds. There is currently no such area in the entire Midwest, and such an area is needed by the many species of birds that make long-ry poloyttayio cott. These T/5p These d by arequuch 0079 Tw T*.164

suggested plan; a control structure around Roxanna Pond probably would not provide any impediment to migration within the river chann18.l itself,ny d ifhe r arwr ary pedt Bs,

Far West reach

The Illinois section of the Grand Calumet River is not part of the study area, but, as a connected river reach, it merits mention in any paper concerned with the River's ecologic functioning. Burnham Prairie is a site located approximately 2 km to the west of the Illinois-

of when river fish are safe under Lake Michigan guidelines. There is no appreciable hunting in the area.

ii) Tainting of fish and wildlife flavor

Assuming that dredging removes contamination, tainting should be reduced.
See *I) Restriction on fish and wildlife consumption*

iii) Degradation of fish and wildlife populations

Fish and wildlife populations will be improved only with habitat improvement following contamination removal. Dredging removes a great deal of allochthonous material and destroys areas of shallow water habitat. Fish and wildlife habitat, including shallow water areas and riffles, must be artificially created in order to encourage re-colonization. Fragmentation of habitat must also be minimized by the creation of corridors between natural areas, to prevent degradation of populations due to shrinkage of the gene pool for rare species.

iv) Fish tumors and other deformities

Fish deformities should be greatly reduced, if not eliminated, with contamination removal. See *I) Restriction on fish and wildlife consumption*

v) Bird or animal deformities or reproduction problems

Removing contaminants from the river will improve the quality of the fish, over several generations, that the birds are consuming. Therefore, there should be long-term improvement, but immediate results are not likely. There is a risk of harming bird and animal populations far from the River because many organisms that feed in the area are migratory, and this might be eliminated with contamination removal. Pollution abatement is essential to prevent re-contamination. Another potential threat is the area where dredge spoil is stored. Birds and other animals that land, live, or feed in this area will be at risk of contamination, so measures should be taken for the construction of suitable storage facilities.

vi) Degradation of benthos

Dredging should eliminate a majority of the contaminated sediment that is limiting species diversity in the benthos. Pollution abatement is imperative since most contamination accumulates in the sediment. Also, re-contamination from bank sloughing is a possibility if all of the contamination is not removed. Addition of clean sediment should promote colonization by tolerant species, provided that the appropriate sediment type is used and that water quality is improved.

vii) Restrictions on dredging activities

Contaminated sediment brought up from dredging should not pose a threat if a proper disposal plan is in place. Because of the nature of the contaminants, phototoxicity is a possibility, but if proper precautions are taken, risks can be avoided. Disposal areas should be lined and capped so that neither groundwater nor air becomes contaminated.

viii) Eutrophication or undesirable algae

Eutrophication is a result of poor water quality. To address this problem, nutrient inputs must be eliminated. Dredging will not impact the persistence of undesirable algae. Diversity will probably increase when algal species re-colonize, but blue-green algae will dominate regardless of sediment condition.

ix) Restrictions on drinking water consumption or taste and odor problems

Long-term effects on drinking water are not at issue since the public drinking supply comes from Lake Michigan. However, dredging may affect the quality of drinking water tnot i5fEsTc 28nce the public drink

The ecosystem for phytoplankton and zooplankton is toxic now, and removing contamination can only improve conditions. Phytoplankton community structure depends on nutrients and flow rate. A decrease in nutrient loading is essential for improving the phytoplankton populations. Zooplankton are greatly affected by the toxic sediment, so dredging should be an improvement. However, more information on resident planktonic communities is needed for a thorough estimation.

xiv) Loss of fish and wildlife habitat

Fish habitat availability will decrease because of the smooth, U-shaped channel that results from dredging. Contamination should decrease and dissolved oxygen should increase, but fish populations will not be attracted to the area unless artificial habitats are created. For this reason, habitat enhancement in the river channel is an essential part of post-dredging restoration. Other animals should benefit from less contamination, but again, habitat must be created. Dredging along the banks to create shallow backwaters or to eliminate exotic seedbeds could, combined with active ongoing management, create some of this habitat. See *iii) Degradation of fish and wildlife populations.*

CONCLUSIONS

Preservation and restoration in the Calumet region are necessary to re-create viable natural systems in the wake of contamination and fragmentation from human settlement and industry. The historic habitats of the region were diverse and formed from floras of several different regions of the country. Despite extensive degradation, several rare and highly valuable natural areas remain, as fragments of the native landscape. For these to persist as viable, functioning ecosystems, extensive restoration measures must be taken. Dredging and replacement of sediments in the Grand Calumet River are first steps in the process, but they will need to be accompanied by preservation and management activities to be successful.

Dredging will remove many contaminants from the River, but it probably will not fully clean the river system and may have some negative effects. To minimize the negative effects of dredging, banks should be stabilized and fish micro-habitat should be artificially re-constructed. Pollutant concentrations in the waters and sediments of the River should be monitored after dredging to assure that they stabilize at low levels. For dredging to be effective in the long term, re-contamination of the River from point and non-point sources will have to be stopped.

Habitats of the Grand Calumet River system have been degraded not only by contamination, but by fragmentation, invasion of aggressive species, and suppression of natural processes. These problems will also need to be addressed. Buffer areas are needed to prevent further contamination of high quality areas. Corridors of semi-native habitat should be created between high quality areas, to allow for migration of species between them. Populations of invasive species should be eliminated to the greatest extent possible, and replaced with assemblages of native species. Habitats that require fire to persist should be burned on an ongoing basis. Extirpated species should, when appropriate, be re-introduced to the region. Areas should be managed to create a variety of different habitat types, and habitat for rare species should be given priority in restoration. Restoration plans will need to take current conditions, such as the unprecedented rapid flow of the present-day River, into account. It may

not be possible to return to presettlement conditions, but efforts should be made to create a diverse set of ecologic conditions and habitats in today's landscape that have the potential to persist over time and, to some extent, to mimic presettlement conditions.

To accomplish the goal of an integrated, viable landscape, a new land-use ethic will have to be established. Diverse groups will need to cooperate to make restoration a consistent, long-term project. All aspects of the system will require ongoing monitoring, and plans for preservation and restoration will need to be integrated at every stage of the land-use planning process. By acknowledging that the ecologic systems of the region must be understood and maintained, we will move toward the goal of an integrated, functioning landscape where human inhabitants and native communities can successfully coexist.