

The Effect of Seagulls on Recreational Water Quality at Lake Calumet, Illinois**Brie Betz**

Abstract The Lake Calumet region, located 12 miles south of Chicago's downtown financial district, has experienced great industrial growth and decline over the last century and a half, and it has become an isolated sanctuary to ring-billed and herring gulls in recent years. With the slow departure of manufacturing facilities and the degeneration of the surrounding factory supported towns, city officials and locals have suggested turning the lake into a recreational attraction in hopes of community renewal. The lake is currently a gateway for many inland boaters to reach Lake Michigan, and so city planners hope that creating a public park on the lake will attract more nature and boating enthusiasts and in turn more development in the community. Due to the presence of Illinois' largest gull rookery and the correlation of seagulls with heightened bacterial loads, this level of recreation may, however, be unsafe for people to use. After testing various water quality parameters on the lake, I concluded that *Escherichia coli* levels were highest at sites nearest to the gull rookery and with the lowest dissolved oxygen levels. The average *E. coli* load sampled surpasses the US EPA's fresh recreational water standard of 126 cfu, and my results suggest that the presence of the state's largest seagull population is currently preventing recreational use in the lake. My policy recommendation would be to prevent further recreational development on the lake at this time.

can affect *E. coli* levels. One particular study in Corpus Christi, Texas, found that high water temperatures lead to decreased bacterial levels (Crysup and Mott 2001), and another study found that decreased amounts of dissolved oxygen can inhibit *E. coli* reproduction (Sandoval-Basurto *et al.* 2005).

With gulls being a proven source of *E. coli* and *E. coli* being an indicator of potentially harmful water-borne illnesses, lakes with large gull populations should monitor water quality for unsafe bacterial levels. Twelve miles south of the Chicago Loop (the financial district), in a remote part of the southeastern side, is one such lake. This lake, Lake Calumet, is home to Illinois' largest ring-billed and herring gull nesting site, and in the early summer months (the onset of the mating season), the rookery (nesting site) is crowded with gull activity (Calumet Environmental Education Program n.d.). The rookery is only one of the distinguishing features of Lake Calumet; it is also enclosed by a golf course, landfill, and several rare and threatened ecosystems, including oak savannahs, prairie wetlands, and oak woodlands. In addition, it has a 130-year record of intense industrial use and barge traffic being a gateway from the interior of the Chicago region out to the Great Lakes.

The Illinois International Port District currently owns Lake Calumet's shoreline, and there have been heated debates between various environmental groups, the city of Chicago, and potential developers regarding the future of the lake. Presently, Lake Calumet has limited

Methods

I chose fifteen sample sites, spaced evenly in the upper, middle, lower, and peninsular regions of Lake Calumet (Fig. 1). Over the course of the summer, I sampled four times (July 1,




Figure 1: Aerial View of Lake Calumet Sampling Sites (Upper, Middle, Lower, and Peninsular) with Measured Through-Water Distances to the Gull Rookery

Results

After averaging all of the *E. coli* samples (Fig. 2), I found that the mean *E. coli* level is 301.7 cfu (with a standard error of 77.6) which well exceeds the USEPA's fresh recreational water quality standard of 126 cfu (average weekly sample over a single month) (USEPA 2004).




Figure 2: *E. Coli* Levels For Each Sampling Event With an Overall Average of 301.7 cfu

Because of the established correlation of seagulls and elevated bacterial levels, I was attempting to determine whether the gulls are responsible. My hypothesis claimed that if the gulls were causing the elevated levels of *E. coli*, then sites nearest to their rookery should show a correlated increase in bacteria. Due to behavioral observation of the gulls, I predicted that through-water distance would be a more accurate measure of how the bacteria are being transported around the lake. The through-water distances (Fig. 3) did show a stronger correlation to *E. coli* levels than the aerial distances (Fig. 4), and when I plotted a linear fit to these through-water distance and *E. coli*, through-water distance from the rookery explains 28 percent of the variability in *E. coli* levels. The relationship is $\ln(e. coli) = 3.995 - 0.803 \ln(\text{water distance})$, with an R-Square value of 0.284 and the regression had a P-value of $<.0001$. In general, distance from the gull rookery is in fact affecting *E. coli* loads.

In an attempt to verify this correlation, I needed to eliminate any confounding variables, which would be parameters other than distance that are affecting *E. coli* loads, to be certain that distance alone is affecting *E. coli*. To determine whether other water quality parameters were influencing *E. coli* levels, I ran a multivariate analysis on all of my parameters to look for

I then sorted the regression by sampling day and found that although the model explained 65 and 80 percent of the variability in *E. coli* for outings one and two, respectively, it only explained 34 and 28 percent for sampling days three and four.

Discussion

My research shows that there is in fact a correlation between distance from the rookery and *E. coli* levels, and this suggests that the gulls are indeed responsible for the elevated levels of bacteria in the lake. The connection to dissolved oxygen is a bit more difficult to explain, however, because dissolved oxygen fluctuates naturally with time of day, water temperature, depth, and amount of wind (Water On The Web n.d.). This suggests that although my results showed no overall correlation between time of day and water temperature, there may be other possible natural conditions affecting this relationship. For instance, although I measured the total depth of the lake at each sampling site, I took all measurements at a depth of 1 meter. Because dissolved oxygen varies by depth (with oxygen production occurring nearest to the surface), I may be seeing a false correlation without the average dissolved oxygen level for each sampling site. Secondly, I was not able to measure or quantify the amount of wind incident on the lake on any of the sampling days. I made observational notes about the direction of the wind, but the amount of wind is more crucial because it can induce mixing of the surface layers of the lake, thereby increasing the amount of dissolved oxygen at a given site (Water On The Web n.d.). Although there is no noticeable correlation between sampling day and dissolved oxygen level, the amount of wind or some other weather condition may explain the variance in R-square values for each of the four sampling days.

It is unfortunate that the dissolved oxygen data are not conclusive because my research disagrees with a previous study on the inverse relationship between dissolved oxygen and *E. coli* levels. My results are countered by Sandoval-Basurto's study that suggested a directly proportional relationship between dissolved oxygen and *E. coli* levels (2005). This study suggested that *E. coli*

that increased water temperature decreased bacteria levels. Reasons for uncertainty in my conclusion could include the narrow water temperature range I measured as well as a possible fatal temperature at which the bacteria cannot survive. Warm environments can be good breeding grounds for certain pathogens so it is difficult to conclude whether this should be a positive or negative feedback or have no significant interaction whatsoever.

Overall, my research concludes that the water in Lake Calumet is unsuitable for recreational activity; however, my inference that the gull population was responsible for the heightened levels of bacteria is not complete. I was biased in assuming that bacteria from the fecal contamination could only move through the water from the rookery, and in fact, my method of measuring the through-water distances would be difficult to replicate. I think that the interactions between the water quality parameters and the weather conditions might not be considering or observing all interactions, and therefore, other conclusions could be reached about the correlation of distance to the gull rookery and bacteria. In general, my multiple-linear regression fit does not seem ultimately to address the question of whether the gulls are responsible, but I have concluded that the water is un-swimmable by current US EPA fresh recreational water standards. Future recreational development on the lake should be discouraged.

Acknowledgements

I would like to thank my mentor Rob Sulski (Illinois EPA engineer) and my fellow intern Sean Babington (University of Colorado Boulder undergraduate) for assisting me in data collection, the Illinois EPA for funding the bacterial analysis conducted at Stat Labs (Chicago, IL), and to Cristina Castanha and Mike Dwyer for their support and assistance.

References

- Calumet Environmental Education Program. (n.d.)
http://www.fieldmuseum.org/ceep/infoSci_thrive.html.
- Crysup, K. A. and J. B. Mott. "Fecal coliforms, enterococci and E. coli as indicators of water quality in the Oso Creek/Oso Bay Watershed, Corpus Christi, Texas"(2001). 101st General Meeting of the American Society for Microbiology, Orlando, FL, USA; May 20-24, 2001.
- Fitzpatrick, W. P. and N. G. Bhowmik. "Pollutant transport to Lake Calumet and adjacent wetlands and an overview of regional hydrology"(1990). *Illinois State Water Survey: Report no. HWRIC-RR-050*, 88 pp.

Fogarty, L. R.; S. K. Haack; M. J. Wolcott; and R. L. Whitman. "Abundance and characteristics

