THE STATE OF LAKE MICHIGAN IN 2000

Prepared for the Lake Michigan Committee



SPECIAL PUBLICATION 05-01

The Great Lakes Fishery Commission was established by the Convention on Great Lakes Fisheries between Canada and the United States, which was ratified on October 11, 1955. It was organized in April 1956 and assumed its duties as set forth in the Convention on July 1, 1956. The Commission has two major responsibilities: first, develop coordinated programs of research in the Great Lakes, and, on the basis of the findings, recommend measures which will permit the maximum sustained productivity of stocks of fish of common concern; second, formulate and implement a program to eradicate or minimize sea lamprey populations in the Great Lakes.

The Commission is also required to publish or authorize the publication of scientific or other information obtained in the performance of its duties. In fulfillment of this requirement the Commission publishes the Technical Report Series, intended for peer-reviewed scientific literature; Special Publications, designed primarily for dissemination of reports produced by working committees of the Commission; and other (non-serial) publications. Technical Reports are most suitable for either interdisciplinary review and synthesis papers of general interest to Great Lakes fisheries researchers, managers, and administrators, or more narrowly focused material with special relevance to a single but important aspect of the Commission's program. Special Publications, being working documents, may evolve with the findings of and charges to a particular committee. Both publications follow the style of the *Canadian Journal of Fisheries and Aquatic Sciences.* Sponsorship of T

THE STATE OF LAKE MICHIGAN IN 2000

Prepared for the Lake Michigan Committee

Edited by

Mark E. Holey
United States Fish and Wildlife Service (USFWS)
Green Bay Fishery Resources Office
2661 Scott Tower Drive
New Franken, Wisconsin, U.S.A. 54229-9565

Thomas N. Trudeau
Illinois Department of Natural Resources
9511 Harrison St.
Des Plaines, Illinois, U.S.A. 60016

TABLE OF CONTENTS

Executive Summary 1

Goal and Objective Setting, Mark E. Holey 7

Nutrients, Glenn Warren 15

Phytoplanton and Zooplankton, Richard P. Barbiero and Marc L. Tuchman 21

Planktivores, Guy Fleischer, Charles M. Madenjian, Robert F. Elliott, and Michael L. Toneys 25

Salmonine Community, Jory L. Jonas, David F. Clapp, James R. Bence, and Mark E. Holey 33

Inshore Fish Community, David F. Clapp, Philip J. Schneeberger, Richard P. O'Neal, Terry J. Lychwick, Brian Belonger, and Steven M. Shroyer 49

Benthivores, Philip J. Schneeberger, Robert F. Elliott, Jory L. Jonas, and Stephanie Hart 59

Sea Lamprey, Dennis S. Lavis 75

Fish Health, Greg Wright, Dale C. Honeyfield, Scott B. Brown 81

Physical and Chemical Habitat Remediation, Mark E. Holey 87

Open Forum, Mark E. Holey and Thomas N. Trudeau 91

Literature Cited 95

shifted from diatoms to cyanophytes and chlorophytes in the late 1960s and 1970s and then to phytoflagellates in the 1980s and 1990s. The species composition of the zooplankton community has been relatively consistent during the past 15 years. Efforts to restore wetland habitat are under way and some major strides have been made in restoration of historical riverine spawning and nursery habitat, including installation of fish-passage facilities and requiring run-of-the-river flow regimes for dams.

Pursue the reduction and elimination of toxic chemicals, where possible, to enhance fish survival rates and allow for the promotion of human consumption of safe fish.

Goal and Objective Setting

Mark E. Holey
United States Fish and Wildlife Service
Green Bay Fishery Resources Office
2661 Scott Tower Drive

of the fish community. An alphabetical list of the common fish names and their corresponding scientific names is given in Table 1.

Table 1. A list of common and scientific fish names used in this publication.

Common name	Scientific name
alewife	*Alosa pseudoharengus*
Atlantic salmon	*Salmo salar*
bloater	*Coregonus hoyi*
brook trout	*Salvelinus fontinalis*
brown trout	*Salmo trutta*
burbot	*Lota lota*
catfish(s)	*Ictalurus* spp.
chinook salmon	*Oncorhynchus tshawytscha*
coho salmon	*Oncorhynchus kisutch*
deepwater cisco	*Coregonus johannae*
deepwater sculpin	*Myoxocephalus thompsoni*
emerald shiner	*Notropis atherinoides*
johnny darter	*Etheostoma nigrum*
lake herring	*Coregonus artedi*
lake sturgeon	*Acipenser fulvescens*
lake trout	*Salvelinus namaycush*
lake whitefish	*Coregonus clupeaformis*
mottled sculpin	*Cottus bairdi*
muskellunge	

Fig. 1. Lake Michigan depicting inshore (<45 m) waters (shaded), southern and

Beeton et al. (1999) classified Lake Michigan as generally oligotrophic because of its low concentrations of phosphorus, but they reported that the level of total dissolved solids (150 mg/L) is normally associated with eutrophic conditions indicating a possible trend toward mesotrophy. Total alkalinity is 113 ppm; the concentrations of calcium, magnesium, and sodium are 31.5, 10.4, and 3.4 ppm, respectively; the phosphorus concentration is 0.9 ppb; and dissolved oxygen concentrations are near saturation at all depths (Beeton and Chandler 1963).

Goals and Guiding Principles

In recognition of the GLWQA's call for restoration of system integrity and the Joint Plan's focus on sustainable benefits, the LMC's goal statement emphasizes biological integrity, production of desirable fish, and ecological efficiency.

Restore and maintain the biological integrity of the fish community so that production of desirable fish is sustainable and ecologically efficient.

Although the GLWQA divides the ecosystem into three components (chemical, physical, and biological), it implies that integrity is an attribute of the ecosystem as a whole. Fish-community goals, however, address issues related primarily to biological integrity, and achieving them will be limited inevitably by the physical and chemical integrity of the Lake Michigan ecosystem. The term biological integrity was at best abstract and somewhat ambiguous when introduced in water-quality legislation in the 1970s (Karr et al. 1986). Since that time, it has been reworked and defined as, "…the ability of an ecosystem to maintain its structure when confronted with environmental change. Systems that cannot maintain their structure following perturbation are said to have lost their integrity" (Karr and Dudley 1981).

The fluctuations over time in the Lake Michigan fish community can provide examples of the concepts of integrity and ecological efficiency. Before the arrival of European settlers, the fish community was considered to be stable and organized. Lake herring and deepwater ciscoes occupied the offshore zones, and lake whitefish and emerald shiners were important in the inshore areas (Wells and McLain 1972). Burbot and lake trout were the main predators. The low diversity of top

predators was compensated for by differentiation of lake trout into specialized shallow-water and deep-water forms enabling them to utilize prey resources throughout the entire water column (Brown et al. 1981; Eshenroder et al. 1995). Attempts to introduce Pacific salmon in the 1900s failed (Parsons 1973) possibly due to the integrity or stability of the fish community at that time. This system remained stable until the invasion of the sea lamprey, which, coupled with extensive fishing pressure, led to the extirpation of the

only a few options exist for altering community structure in a Great Lake. Habitat manipulation is usually limited to remedial action in nearshore environments and tributary streams. Beyond remediation of habitat, managers exert an influence through the regulation of fisheries, stocking, and sea lamprey control. A third reality is that management actions are inexact—their effects cascade through the trophic pyramid to species well beyond those targeted, and those effects can have different time scales for different species. Short-term responses can be deceptive and long-range prediction can prove difficult. FCOs for an entire lake cannot be taken to a high level of exactness—they are reasoned approximations of likelihoods. Management initiatives aimed at achieving objectives will continue to have a large experimental component, and the time frame needed in meeting some objectives will be measured in decades. Last, humans, more than any other organism on earth, have the capacity to alter their environment to their own benefit and desire. Society places a myriad of demands on the Lake Michigan fish community. The Lake Michigan FCOs are designed to produce the fish desired by society within the framework of the capacity of the lake and its biological integrity. Specific objectives will be identified in appropriate chapters.

Nutrients

Glenn Warren
United States Environmental Protection Agency
Great Lakes National Program Office
77 W. Jackson Boulevard
Chicago, Illinois, U.S.A. 60604

Chemical sampling of Lake Michigan, begun in 1983, was done annually at 11 stations in the spring at a time when the lake is homothermal and behaving similar to a modeler's assumption of "mixed reactors." The spring data will be the focus of this report. The main chemicals of interest are total phosphorus, nitrate/nitrite nitrogen, and dissolved silica, which have major effects on algal production (United States Environmental Protection Agency 2003).

Total Phosphorus

The water-quality guideline for concentration of total phosphorus in the open waters of the lake was established at 7 µg/L (International Joint Commission 1980) to maintain the lake in an oligotrophic to mesotrophic state. Total phosphorus concentrations were below the guideline and ranged from a high of 6.4 µg/L in 1983 to a low of 3.8 µg/L in 1992 (Fig. 2). This downward trend in concentration was reversed after 1992. Since 1994, concentrations have ranged from 5.5 µg/L-6.3 µg/L, very similar to levels that occurred prior to 1990. This increase from the low levels of 1992 is significant ($p < 0.05$) for all years after 1994 and indicates that the decreases seen between the late 1970s and 1992 have been reversed. The concentrations of total phosphorus in the northern and southern basins were not significantly different.



Fig. 2. Total phosphorous concentrations (μg/L) in Lake Michigan (northern, southern, lakewide), 1983-1998. Error bars (±2 standard errors) are included for lakewide data.

The most recent load estimates for total phosphorus are from the Lake Michigan Mass Balance Study and, at approximately 2.5 x 10^6 kg/yr for 1994 and 1995, are close to the average load (3.1 x 10^6 kg/yr) from 1980-1991 based on International Joint Commission (IJC) information (Robertson 1997). The estimates from the study are easily within the

Nitrate/Nitrite-Nitrogen

Nitrogen concentration, particularly nitrate/nitrite nitrogen, has been increasing in Lake Michigan through the monitoring period (Fig. 3). The lakewide average increased from 0.262 mg/L in 1983 to 0.311 mg/L in 1999, with a low of 0.257 mg/L in 1984 and a high of 0.342 mg/L in 1993. Neither basin was consistently higher in nitrate/nitrite nitrogen concentration. Increasing nitrate/nitrite concentrations in the lake,

Phytoplankton and Zooplankton

Richard P. Barbiero
Computer Services Corporation
1359 W. Elmdale Ave Suite 2
Chicago, Illinois, U.S.A. 60660

Marc L. Tuchman
United States Environmental Protection Agency
Great Lakes National Program Office
77 W. Jackson Boulevard
Chicago, Illinois, U.S.A. 60604

The Great Lakes National Program Office (GLNPO) of the United States Environmental Protection Agency has primary responsibility for conducting surveillance monitoring of the offshore waters of the Great Lakes. In this report, we provide a brief summary of the results of the planktonic component of GLNPO's spring (March-May) and summer (August-September) biological surveillance sampling at 11 open-water stations on Lake Michigan inTT0 5,and sompo-2(ale whe se data both)6(twith)6TJ0.0006 Tc 0.108. T

Lake Michigan underwent a dramatic decrease in dissolved silica concentrations from the mid-1950s through 1970 (Schelske 1988), resulting from increased phosphorus loading and a consequent increased deposition of silica to the sediments through increased diatom production (Schelske and Stoermer 1971). These changed nutrient conditions are thought to have promoted a shift in the dominance of the summer community from diatoms to chlorophytes and cyanobacteria during the late 1960s and 1970s (Fahnenstiel and Scavia 1987).

In the 1980s, a further shift towards summer dinoflagellate dominance occurred (Fahnenstiel and Scavia 1987), which our data indicate has continued through the 1990s (Fig. 6). More recently, reductions in phosphorus loadings have reversed the trend of decreasing silica and resulted in increases in summer diatom populations (Barbiero et al. 2002). This trend will likely continue as the silica content of the lake continues to increase, resulting in phytoplankton communities that are closer to the historical condition of year-round diatom dominance.



Fig. 6. Phytoplankton biomass by major taxonomic group in Lake Michigan, summer 1983-1999.

Zooplankton

The median lakewide abundance of crustaceans (excluding nauplii) during spring 1999 was 2,690 animals/m^3, with densities in the southern basin greater than those in the northern basin. Only 11 crustacean taxa were found in the lake during spring. The cyclopoid *Diacyclops thomasi* and the calanoids *Limnocalanus macrurus, Leptodiaptomus ashlandi,* and *L. minutus,* along with immatures of these genera, accounted for most of the spring zooplankton. Lakewide abundances increased in August to 42,655 animals/m^3, but no clear spatial differences in abundances were noted. *D. thomasi* remained one of the dominant species with the smaller cladoceran *Bosmina longirostris* and the typical summer dominant *Daphnia mendotae* also contributing substantial numbers. Two predatory cladocerans were found, the native *Leptodora kindti* and the exotic *Bythotrephes longimanus,* although abundance of both were less than 5 individuals/m^3.

The crustacean community of Lake Michigan has undergone a number of changes in the past 20 years. Prior to 1987, the cladoceran community included three daphnids: *D. retrocurva, D. mendotae* and *D. pulicaria* (Evans and Jude 1986). Since the invasion of *B. longimanus*, *D. retrocurva* and *D. pulicaria* have virtually disappeared from the offshore waters, leaving *D. mendotae* the sole daphnid. A number of less-dominant crustacean species, including *Holopedium gibberum, Eubosmina coregoni,* and the cyclopoid copepod *Mesocyclops edax*, have also declined dramatically (Barbiero and Tuchman 2004). Except for *M. edax,* abundance of the numerically dominant copepod community has not changed and continues to be dominated by the same species of diaptomids (*L. ashlandi*, *L. minutus,* and *Skistodiaptomus oregonensis*) and the cyclopoid *D. thomasi* (Barbiero et al., 2005).

In general, cladoceran abundances were higher during 1991-1999 than 1983-1990 (Fig. 7), perhaps indicating reduced predation pressure. Abundances were unusually low in 1998 (Barbiero et al. 2001) compared to the rest of the decade. In 1999, the crustacean community exhibited a marked increase in the smaller cladoceran *B. longirostris* and the cyclopoid copepod *D. thomasi*.



Fig. 7. Total zooplankton density and relative abundance (%) of major taxonomic groups in Lake Michigan during summer, 1983–1999.

Planktivores

Guy Fleischer and Charles M. Madenjian
United States Geological Survey
Great Lakes Science Center
1451 Green Road
Ann Arbor, Michigan, U.S.A. 48105

Robert F. Elliott
United States Fish and Wildlife Service
Green Bay Fishery Resources Office
2661 Scott Tower Drive
New Franken, Wisconsin, U.S.A. 54229-9565

Michael L. Toneys
Wisconsin Department of Natural Resources
110 South Neenah Avenue
Sturgeon Bay, Wisconsin, U.S.A. 54235-2718

The segment of Lake Michigan's fish community that we classify as planktivores includes species that, as mature adults, prey extensively on zooplankton. We recognize that juvenile life stages of all fish rely on plankton for sustenance and that invertebrates even contribute to the diet

bloaters, the surviving member of the deepwater cisco complex in Lake Michigan (Fig. 8). More recently, bloaters have peaked and subsequently declined in abundance in an apparent density-dependent response to high abundance (TeWinkel et al. 2002) rather than as a response to interactions with other species.



Fig. 8. Biomass of major planktivores (adult) in Lake Michigan, 1973-1999. Estimates based on bottom-trawl surveys performed by the USGS Great Lakes Science Center, Ann Arbor, Michigan.

Planktivore Abundance

Trends in planktivore populations, as indicated from both bottom-trawl catches and acoustic surveys, show that alewife abundance has fluctuated with no consistent trend in recent years. A strong 1995 year class, observed in bottom trawls as adults (Fleischer et al. 1999) and detected in the acoustic surveys as young of the year (YOY) and yearlings (Argyle et al. 1998), now dominates the alewife population. Rainbow

smelt have declined in abundance lakewide throughout the 1990s to new low levels. Due to poor recruitment during the 1990s, bloaters have also declined in abundance to the lowest levels since the 1980s (TeWinkel et al. 2002). Deepwater sculpin (not a planktivore but in the planktivore food web) populations have been stable in recent years (Fleischer et al. 1999). Ninespine stickleback catches, which had previously not shown any trends, have dramatically increased in the most recent years.

The lakewide biomass estimates presented here do not include Green Bay or Traverse Bay. Brandt et al. (1991) found that Green Bay contributed 5-14% of the total biomass during 1987. During the 1990s, alewife biomass has ranged from 13,000 to 54,000 metric tons for adults and 8,400 to 148,000 metric tons for YOY (the latter value is the estimate for the 1995 year class). The biomass of the 1995 year class of yearling alewives was 38,000 metric tons. Standing stocks ranged from 3 to 13 kg/ha for adults and from 3 to 35 kg/ha for juveniles (YOY and yearling). In total, alewife standing stocks ranged from 9 to 47 kg/ha. Based on trawl surveys, bloater biomass has declined from 326,000 to 45,500 metric tons during the1990s; acoustic estimates suggest a peak biomass of 475,000 metric tons. These values correspond to standing stocks ranging from 93 to 13 kg/ha, and a peak acoustic estimate of 115 kg/ha. Acoustic-based estimates of rainbow smelt biomass declined from 84,000 to 16,000 metric tons, and standing stock declined from 20 kg/ha to 3 kg/ha. The biomass of deepwater sculpin ranged from 21,500 to 52,000 metric tons, and standing stocks ranged from 6 to 15 kg/ha. Ninespine stickleback biomass varied between 210 and 9,000 metric tons (0.1-2.5 kg/ha standing stock). These values compare favorably with estimates of biomass in other large freshwater systems. In Lake Tanganyika, Moreau et al. (1993) modeled the trophic structure and estimated the standing stock of smaller pelagic fishes at 65 kg/ha. Using the same approach in Lake Oneida, Stewart et al. (1981) estimated

tdks 0.0009 Tc -0.0009 Tw [(Lake Ontario, Half)8(a Superior)39(,Tw -14.978 -1.vorably)-795 y[(the)-5(sam)8(e

a demersal species, contributed on average only about 33,000 metric tons. These values represent an overall standing-stock range of 150-65 kg/ha, with an average of 114 kg/ha for pelagic planktivores and 9 kg/ha for deepwater sculpin. By way of comparison, Brown (1972) estimated an alewife standing stock of 191 kg/ha.

Demands on Planktivores as Prey

During the past three decades, the lake trout restoration program and introductions of hatchery-reared trout and Pacific salmon have resulted in successful and highly desired fisheries. Due to this success, attention has been focused on the capacity of the planktivore populations, especially of alewife, to sustain these predators. The dramatic shift in planktivore dominance from alewife to bloater that occurred during the late 1980s did not result in a similar shift to bloaters as the dominant prey for most predators. Several authors have concluded that much of the adult bloater biomass in Lake Michigan is too deep and occupies water too c8tp.8o

Michigan, 48105-1563, unpubl. data) and CONNECT (Rutherford 1997) model estimates of total salmonine biomass during 1978-1988, which included estimates of naturally reproduced and age-4 fish, are substantially higher than the estimates of total salmonine biomass by Stewart and Ibarra (1991). According to the CONNECT and SIMPLE models, salmonines were eating between 70,000 and 95,000 metric tons of alewives per year during 1978-1988, which is two to three times higher than estimated by Stewart and Ibarra (1991).

Demands on Planktivores as Commercial Species

A commercial fishery for alewife was established in the Wisconsin waters of Lake Michigan in the 1960s when a trawl fishery was developed to harvest the then extremely abundant alewife that had become a nuisance and health hazard along the lakeshore (Fig. 9). Although alewife was the target species, the trawls caught bloater and rainbow smelt. Bloaters and rainbow smelt not sorted and sold for human consumption were sold, along with alewives, for fishmeal and pet food. In 1986, a quota on alewife was implemented, which was replaced by a targeted rainbow smelt only trawl fishery in 1991. Because of these rule changes and seasonal and area restrictions, the alewife harvest declined from about 7,600 metric tons in 1985 to an average (now incidental) harvest of 12 metric tons after 1990.



Fig. 9. Commercial catch of alewife, rainbow smelt, and bloater in Lake Michigan, 1985-1998 (Fleischer 1992; Kubisiak 2000).

The commercial harvest of bloater and rainbow smelt has also declined substantially since 1985 due to changes in population abundance and in regulation (Fig. 9). The bloater harvest declined from almost 4,700 metric tons in 1990 to less than 1,300 metric tons in 1998. Very low recruitment since 1991 is at least partly responsible for this decline in harvest, especially in recent years. The commercial harvest of rainbow smelt occurs almost exclusively in Wisconsin waters, and trawls account for most of the harvest. The harvest of rainbow smelt has declined substantially from a high of about 2,400 metric tons in 1986 to less than 350 metric tons in 1998. The decline is attributed to poor recruitment os fc 0.1642 inigh ofs aot0tet

Planktivore Biomass Expectations

The planktivore objective for Lake Michigan is to maintain a diversity of

Pink salmon, an accidental introduction into Lake Superior in the 1950s, naturalized and spread to other Great Lakes, including Lake Michigan where populations are small. Brook trout and the brook trout-lake trout hybrid (splake) are stocked in small numbers at a few inshore locations in the northern part of the lake and Green Bay.



Fig. 10. Harvest of major salmonines from Lake Michigan 1920-1997, target harvest range (shaded area), and near-term yield expectations (dashed line) (Eshenroder et al. 1995).

Salmonines were introduced into the Great Lakes for several reasons. Rainbow trout, brook trout, splake, and brown trout were introduced to provide more-diverse fishing opportunities, whereas chinook and coho salmon were introduced to control the large populations of introduced alewife and diversify fishing opportunities (Tody and Tanner 1966; Keller et al. 1990).

Lake Michigan's salmonine fisheries are currently sustained in large part by stocking. However, as introduced salmonines naturalized, determining

the amount of natural reproduction is increasingly important. Because lake trout is the only major salmonine predator that was native to Lake Michigan and knowledge of impediments to their recruitment is incomplete, rehabilitation remains an important and yet unattained goal for managers and researchers.

Salmonine Objectives and Yields

The FCOs for Lake Michigan salmonines (trout and salmon) are: 1) to establish a diverse salmonine community capable of sustaining an annual harvest of 2.7-6.8 million kg, of which 20-25% is lake trout, and 2) to establish self-sustaining lake trout populations (Eshenroder et al. 1995). These objectives recognize the limits of the system and the need to balance a desire for abundant populations of top predators with the possibility that overstocking could lead to a collapse of planktivore populations and instability of predator populations. The decline in chinook salmon fisheries during the late 1980s provided an example of the limits of the Lake Michigan fish community. The objectives of the current management approach include sustaining a diverse predator community that supports sport and commercial fisheries, and that utilizes alewife and rainbow smelt populations sufficiently to minimize the negative influences of these exotic planktivores on native species, especially on the bloater (Eshenroder et al. 1995). Declines in alewife abundance and increases in bloater abundance during the early 1980s emphasized the advantages of a diverse salmonine community that could take advantage of a changing prey-fish community. The lower bound of the desired salmonine yield, 2.7 million kg, was determined from the historical catch of lake trout before the collapse of the native population. The upper bound, 6.8 million kg, was determined from biomass size-spectrum models (Borgmann 1987; Sprules et al. 1991) assuming 100% ecological efficiency. Model estimates indicated that a mix of salmonine species would more efficiently use the pelagic fish community, and thus could support higher yields than were realized from a fish community with only lake trout as the top predator. The LMC agreed to the following species-specific near-term yield expectations (within the 2.7 to 6.8 million-kg range) chinook salmon—3.1 million kg; lake trout—1.1 million kg; coho salmon—0.7 million kg; rainbow trout—0.3 million kg; and brown trout—0.2 million kg; or an overall lakewide yield of 5.5 million kg (Eshenroder et al. 1995; Fig. 11). The LMC recommended

that these initial expectations be refined by evaluating the relation between yield (sport, commercial, harvest-weir) and the mix of predators in the system and that a determination be made as to how these different mixtures of species meet a variety of needs identified by society.

statistics) declined significantly (Rutherford 1997; Fig. 12), probably as a result of density- and forage-mediated increases in disease (Holey et al. 1998). Estimates of biomass for the mid-1990s to late 1990s were at or above former levels. Yield has not increased proportionately with increases in biomass, in part because fishing effort has not increased to levels seen during the 1980s. Management agencies reduced stocking levels of chinook salmon in recent years in an attempt to improve survival by reducing population density.

The relation between the number of chinook salmon stocked and the number harvested changed following the population collapse in the late 1980s. Sport harvest of chinook salmon was positively related to numbers stocked through the late 1980s, but the relation was no longer apparent after 1990 (Fig. 11; Hansen et al. 1990; Hansen et al. 1991; Hansen and Holey 2001). Hence, it now appears that managers cannot increase the yield of chinook salmon simply by increasing the number stocked. Even though the predicted biomass has rebounded to pre-population crash levels or greater, yield remains below the 3.1 million-kg near-term expectation level (Fig. 13) due to reduced fishing effort. Whether or not the chinook salmon harvest can recover to the high levels achieved in the 1980s is now in question. Managers should reexamine the yield expectation for this species.


60
50
40
30
20
10
0
Percent composition
Chinook
Lake trout
Coho
Steelhead
Brown trout

salmon are short-lived, very fast growing, and recruit quickly to the fishery. Consequently, their biomass estimates at the beginning of the year are often well below their annual harvest. Given the yield history and current stocking rates, the near-term yield expectation for coho salmon is probably not realistic.

Rainbow Trout and Brown Trout

Yield of rainbow and brown trout increased during the late 1960s and 1970s as stocking increased (Fig. 11). The relation between yields and stocking in recent years warrants further analysis. Yields and percentage contribution to total salmonine yield for these two species have remained consistently at or above expectations (Figs. 11, 13), so current near-term expectations are probably realistic.

Lake Trout

Yield of lake trout in most years since rehabilitation efforts began in the 1960s has been below the near-term yield expectation identified in the FCOs (Fig. 11). However, that part of the FCOs that state that lake trout should constitute 20-25% of the total harvest has consistently been met in recent years (Fig. 13).

Natural Reproduction and Self-Sustaining Populations

Naturalized populations of salmonines are believed to play an increasingly important role in Lake Michigan fisheries. The FCOs emphasize the desirability of enhanced natural reproduction and establishment, to the extent possible, of self-sustaining populations. Naturalized populations require less-intensive management, may result in more-stable ecosystem dynamics, may increase fitness of populations through genetic selection, and in general are expected to have better survival and productivity in the wild than hatchery-reared fish (Chilcote et al. 1986; Leider et al. 1990; Berejikian et al. 1996). Regulation of hydropower facilities, habitat improvement, and improvements in the water quality of riverine and Great Lakes environments likely contributed to increased productivity of naturalized fish (Holey 2005). In particular, naturalized chinook salmon now make up a large portion of

the chinook salmon population. Based on identification of oxytetracycline (OTC) marks in 1992 and 1993, about 30% of the sport harvest of chinook salmon was made up of naturalized fish (Hesse 1994). Naturalized rainbow trout also make up a large portion of the rainbow trout sport harvest. The proportion of naturalized fish in the entire harvest has yet to be determined, but data from several highly suitable tributaries in Michigan indicate that the contribution of naturalized fish to spawning runs may be as high as 100% (Seelbach and Whelan 1988; Seelbach 1989; Seelbach 1993; Seelbach et al. 1994). In tributaries that are less suitable for rainbow trout reproduction (e.g., St. Joseph and Grand Rivers in Michigan), wild rainbow trout comprise 5-20% of the run (Seelbach et al. 1994). Coho salmon spawn in tributaries of Lake Michigan, but compared to chinook salmon and rainbow trout, much less is known about the contribution of this reproduction to coho salmon populations and fisheries (Becker 1983). Between 1976 and 1979, Carl (1982) found evidence of natural reproduction by coho salmon in 25 of 60 Michigan streams surveyed. Patriarche (1980) reported that 9% of the sport catch in Michigan waters in 1979 was from natural reproduction. Brown trout stocked into Lake Michigan generally do not migrate up streams to spawn, but they may attempt to spawn on structures in the lake. There is little if any information regarding the success of brown trout spawning (Becker 1983). Whether the recruitment of these species has changed considerably since the early 1990s when the FCOs were written is unknown, in large part because lakewide monitoring of natural reproduction has not been consistent. Currently, efforts are being made to increase knowledge of recruitment mechanisms and their influences on estimates of population biomass.

Lake Trout Rehabilitation

Efforts to restore self-sustaining lake trout populations began with the initiation of sea lamprey control, followed by the stocking of lake trout yearlings in 1965 (Holey et al. 1995). Since 1985, a Lakewide Management Plan for Lake Trout Rehabilitation in Lake Michigan (Lake Michigan Lake Trout Technical Committee 1985) has guided restoration efforts. This plan targeted annual stocking at 5.84 million fish, focused stocking efforts in habitats where rehabilitation was judged to have the best chance for success, increased genetic diversity of the stocked fish, and established total mortality limits in areas targeted for rehabilitation

(Krueger et al. 1983; Eshenroder et al. 1984; Lake Michigan Lake Trout Technical Committee 1985; Holey et al. 1995). The plan also defined areas of the lake as refuges or primary, secondary, and deferred rehabilitation zones (Fig. 14). The plan prioritized stocking efforts. Refuges received the highest priority, secondary zones the lowest priority, and deferred zones were not stocked at all. Annual mortality was not to exceed 40% except in deferred zones. Two refuges, mid-lake and northern, were created in 1984-1985 in offshore areas containing high-quality spawning habitat. The refuges were sized to encompass the home range of the fish stocked in them, and sport and commercial harvest was banned to provide maximum protection from fishing (Fig. 14). The rest of Lake Michigan was classified into zones based on the quantity of high-quality spawning habitat, historical lake trout yield from commercial fishing, and total mortality.

Fig. 14. Lake Michigan lake trout refuge areas, and primary, secondary, and

The number of lake trout stocked annually into Lake Michigan has been consistently about 3 million-fish short of the rehabilitation plan target of 5.84 million fish (Holey et al. 1995). Prior to development of a lake trout rehabilitation plan (1965-1984), 53% of available lake trout were stocked into zones designated as secondary or deferred, and only 9% were stocked into areas now designated as refuges. The majority (90%) of lake trout is currently stocked in either refuges (56%) or primary (34%) zones (Fig. 15). After implementation of the 1985 rehabilitation plan, more than 70% of stocked lake trout were transported by boat and released on offshore spawning reefs compared to only 27% prior to plan implementation (Holey et al. 1995).



Implementation of the 1985 rehabilitation plan increased the number of lake trout strains stocked (Fig. 16) in an attempt to address the concern that genetic diversity may be limiting successful rehabilitation (Krueger et al. 1983; Eshenroder et al. 1984; Lake Michigan Lake Trout Technical Committee 1985; Burnham-Curtis et al. 1995). The Marquette strain, a shallow-water lean strain from Lake Superior, was the predominant strain

Inshore Fish Community

David F. Clapp
Michigan Department of Natural Resources
Charlevoix Great Lakes Station
96 Grant Street
Charlevoix, Michigan, U.S.A. 49720

Philip J. Schneeberger
Michigan Department of Natural Resources
Marquette Fisheries Research Station
484 Cherry Creek Road
Marquette, Michigan, U.S.A. 49855

Richard P. O'Neal
Michigan Department of Natural Resources
Central Lake Michigan Management Unit
Oceana Field Office
1757 East M-20
Shelby, Michigan, U.S.A. 49455

Terry J. Lychwick
Wisconsin Department of Natural Resources
200 N. Jefferson
Green Bay, Wisconsin, U.S.A. 54301

Brian Belonger
Wisconsin Department of Natural Resources
P.O. Box 127
Peshtigo, Wisconsin, U.S.A. 54157

Steven M. Shroyer
Minnesota Department of Natural Resources
P.O. Box 86
Waterville, Minnesota, U.S.A. 56096

The inshore fish community of Lake Michigan makes up an extremely important part of the lake ecosystem. Inshore (≤45-m depth) waters make up 31% of the lake area, and include tributary estuaries and drowned river mouths (Fig. 1). Coastal wetlands, especially, provide an interface between the lake and terrestrial habitats. Inshore waters are important areas for nutrient exchange (Hayes and Petrusso 1998) and provide nursery habitat for a variety of fish species (Chubb and Liston 1986; Hayes and Petrusso 1998). Inshore fishes include recreationally and commercially important species such as yellow perch, walleye, smallmouth bass, northern pike, muskellunge, catfish, and panfish, as well as nongame species, including spottail shiner, slimy sculpin, mottled sculpin, trout perch, and johnny darter.


Annual yield (millions of kg)
2.5
2.0
1.5
1.0
0.5
0.0
1900
1920
1940
1960
1980
2000
Year

1990s. Measurable year classes were observed in Green Bay in 1995 and 1998 and in the southern basin in 1998 (Fig. 18).



Fig. 18. Trawl catch (number per hour) of young-of-the-year yellow perch in Green Bay (A) and of age-1 yellow perch in southern Lake Michigan (B), 1975-2000.

Despite poor year classes in the early to mid-1990s, yellow perch remain self-sustaining in Lake Michigan. Recruitment has improved in recent years and, in the long term, the target annual yield of 0.9-1.8 million kg is probably achievable. Factors causing poor yellow perch recruitment need to be identified (Makauskas and Clapp 2000). In Southern Green Bay, there is evidence of predation on yellow perch by exotic white perch, the numbers of which have increased dramatically since the late 1980s (BB, unpublished WDNR file data). In the southern basin, target yields may only be achievable if average alewife abundance is low (Shroyer and McComish 2000). Additional information on dynamics of yellow perch populations will come through development of lakewide population models (Allen 2000; Makauskas and Clapp 2000).

Walleye

The walleye was mainly a commercial fish through the 1950s, but harvest in recent years has been primarily in the sport fishery (Fig. 19). With the exception of the 1890s and 1950s, the annual yield of walleye has generally been below the FCO target range of 0.1-0.2 million kg. Most of the commercial harvest since the 1940s has been from Green Bay. Average annual yield for recent years (1985-1998) was 71,740 kg (range = 37,909-108,915 kg). Yield was within the FCO target range in only three years (1994-1996).



Fig. 19. Annual yield (kg) of walleye from Lake Michigan commercial and recreational fisheries, 1889-1998. Commercial catch data from the waters of Green Bay are available for later years (Comm.-GB). Data are from Baldwin et al. (1979) and Kubisiak (2000).

Walleye utilize tributaries as well as the lake for reproduction. Impoundment of tributaries, declining water quality, high fishing pressure, and abundant exotic species all affected walleye reproduction (Schneider and Leach 1979), such that stocked fish now play a substantial role in walleye recruitment. The Muskegon River is an area of historical importance for walleye in the southern basin (O'Neal 1997). About 600,000 walleye fingerlings have been stocked annually in the Muskegon River system in recent years, and, based on OTC marking, these stocked fish represent the only current source of recruitment to this system (RPO, unpubl. data).

Although walleye recruitment in the southern basin is dependent on stocking, natural recruitment is occurring in Green Bay. Walleye are still stocked (in alternate years) in northern Green Bay (Schneeberger 2000), but walleye are beginning to produce natural year classes in some of the tributary systems (i.e., Cedar River, Menominee River) (Schneeberger

evidence of recruitment in the nearshore areas of southern Lake Michigan from Hammond, Indiana, to Chicago, Illinois (J.T. Francis, Indiana Department of Natural Resources, Lake Michigan Fisheries Station, 100 W. Water St., Michigan City, IN, 46360, personal communication; R. Hess, Illinois Department of Natural Resources, 9511 Harrison St., Des Plaines, IL, 60016, personal communication). However, long-term estimates of recruitment and year-class strength of smallmouth bass are not available for any area of Lake Michigan. Green Bay is probably the only area with substantial populations of northern pike. Green Bay also produced muskellunge historically, and the WDNR is attempting to rehabilitate this population through stocking (BB, personal communication). Stocked fish have survived to adulthood, but reproduction has not been documented. Long-term estimates of recruitment and year-class strength are not available for either northern pike or muskellunge.

Non-Game and Non-Commercial Species

Although not directly harvested commercially or recreationally (with the exception of some limited baitfish harvest), many other inshore fish species are important to the inshore ecosystem. Long-term data are not available for many of these species, but some important population changes for a few have been documented in recent years often coincident with changes in the abundance of sport and commercial species. For example, abundance of spottail shiners increased dramatically in the 1990s coincident with a decline in yellow perch numbers (Tonello 1997). Emerald shiners, in contrast, were scarce throughout the 1970s and 1980s (Eck and Wells 1987) and have remained so through recent years (DFC, unpubl. data). Other changes in non-game, nonecology Bay Td Td (Lake Illild etyees

O'Neal 1997). On the other hand, wetlands along the shore of Green Bay, critical to northern pike reproduction, are being lost to development. Important work is currently under way to quantify and restore wetland and drowned river-mouth resources of Lake Michigan (T. Simon, United States Fish and Wildlife Service, Bloomington Field Office, 620 S. Walker St., Bloomington, IN, 47403-2121, personal communication).

Acknowledgments

The following individuals provided data, comments, ideas, or review that significantly improved the quality of this document: Rich Hess, John Dettmers (Illinois Department of Natural Resources); John Kubisiak (Indiana Department of Natural Resources); Jim Francis, Ed Rutherford, Mark Tonello (Michigan Department of Natural Resources); Mark Holey (United States Fish and Wildlife Service); Guy Fleischer (United States Geological Survey); Matt Coffaro, Pradeep Hirethota, Bill Horns, Tim Kroeff, and Mike Toneys (Wisconsin Department of Natural Resources).

Benthivores

Philip J. Schneeberger
Michigan Department of Natural Resources
Marquette Fisheries Station
484 Cherry Creek Road
Marquette, Michigan, U.S.A. 49855

Robert F. Elliott
United States Fish and Wildlife Service
Green Bay Fishery Resources Office
2661 Scott Tower Drive
New Franken, Wisconsin, U.S.A. 54229-9565

Jory L. Jonas and Stephanie Hart
Michigan Department of Natural Resources
Charlevoix Great Lakes Station
96 Grant Street
Charlevoix, Michigan, U.S.A. 49720

Benthivorous fish species are commonly known as bottom feeders because they target food items that are associated primarily with aquatic substrates. This chapter will cover four benthivorous species in Lake Michigan: lake whitefish, round whitefish, lake sturgeon, and burbot. The lake whitefish is the most important commercial fish in Lake Michigan in terms of numbers caught and monetary value. During 1990-1999, Lake Michigan provided an average 61% by weight and 63% by value of the total lake whitefish commercial yield in the Great Lakes (National Marine Fisheries, http://www.st.nmfs.gov/pls/webpls webst1.MF_GL_SPECIES_HELP .SPECIES#by_lake). Round whitefish have been exploited by commercial fisheries for more than a century, although annual yield is relatively

Lake Whitefish

The historical lake whitefish harvest fluctuated dramatically over the last century, but relatively high and increasing harvests have been sustained since 1971 (Fig. 20). The average annual harvest during 1879-1970 was .983 x 10^9 kg, and the average during 1971-1998 was 2.314 x 10^9 kg. Prior to 1970, harvests above the 1879-1970 average occurred about every 15-40 years. These harvest peaks were short-lived and were attributed to single, strong year classes moving through the fishery. After 1970, average yields increased each decade or so in a stepwise fashion: 1.536 x 10^9 kg during 1971-1980, 2.562 x 10^9 kg during 1981-1990, and 3.418 x 10^9 kg during 1991-1998. The FCO specified a target harvest range for lake whitefish of 1.8-2.7 x 10^9 kg (Fig. 20). Although the lower bound of the targeted range is nearly twice as large as the 1879-1970 average, harvests have been within or exceeded the targeted range since 1979.



Fig. 20. Commercial harvest of lake whitefish from Lake Michigan, 1879-1998, compared with the FCO targeted harvest range (shaded). Data from Baldwin et al. (1979).

Lake whitefish commercial fisheries are monitored at 13 locations around Lake Michigan by personnel from the Michigan Department of Natural Resources (MDNR), Wisconsin Department of Natural Resources (WDNR), and the Chippewa/Ottawa Resource Authority. Although mixing certainly occurs, lake whitefish at these 13 locations are considered to be distinct stocks for management purposes. Proportions of the total harvest taken by Michigan (~45%), Wisconsin (~22%), and Native American (~33%) fishermen remained relatively stable over the last 18 years. An examination of harvest by gear type between 1981 and 1998 showed that trapnets accounted for the largest



Fig. 21. Coefficients of condition (K) for lake whitefish age groups that contribute most fish to the commercial fishery in northern Lake Michigan, 1985-98.

Declines in lake whitefish growth and condition could lead to decreased fecundity and decreased egg quality, and could potentially affect year-class strength and stock stability in future years. Thus far, however, reproduction and survival have been sufficient to sustain productive fisheries even in northern management zones where total annual mortalities are very high (0.70-0.83). These high mortalities are partly offset by ice cover and spring warming conditions, which are generally favorable for overwinter survival of eggs and early growth and survival of fry (Taylor et al. 1987; Brown et al. 1993).

In general, effects of parasites and diseases on lake whitefish populations have not been quantified. Lake whitefish can be infected by *Renibacterium salmoninarum* (Rs), a bacterium that had been previously documented only in salmonines (Jonas et al. 2002). Rs can lead to BKD and cause mortality in chinook salmon, but such a progression has not

structures of round whitefish populations in Lake Michigan, but they appear to be self-sustaining.

Table 2 continued

Location	Adult Status	Run Size	Reproduction	Data Source
Menominee River	Spawning known	>200	Fry produced Juveniles presumed[1]	Tom Thuemler, WDNR Greg Kornely, WDNR
Escanaba River	Occasionally observed	Not quantified (few if any)	No data	Ed Baker, MDNR Nancy Auer, MTU
Manistique River	Regularly observed	Not quantified (<25)	No data	Ed Baker, MDNR

Table 2 continued

Location	Adult Status	Run Size	Reproduction	Data Source
St. Joseph River	Occasionally observed	Not quantified (small numbers)	1 juvenile observed	Jay Wesley, MDNR Ed Baker, MDNR
Wolf Lake, Ill.	Occasionally observed	None known	No data	Rich Hess, ILDNR
Ludington Shoal	Occasionally observed	<10 (1990-99)	No data	Rob Elliott, USFWS Ed Baker, MDNR

Estimates of adult lake sturgeon abundance in tributaries supporting regular spring spawning runs range from just a few fish to several hundred. At least three tributaries have runs of 25 or more adults and two have runs of 100 or more (Table 2). The largest aggregation of lake sturgeon occurs in the Menominee River, tributary to Green Bay. Abundance during summer in the lower Menominee River was estimated to be 457-1329 fish in 1991 (Thuemler 1997). Visual estimates indicate that over 100 adults have been returning to the Peshtigo River each spring, and 30-75 adults have been returning to the Fox River in recent years. Sex ratios on the spawning grounds in the Fox River have been about 1:5, female to male (RFE, unpubl. data). The length range of spawning adults in the Menominee and Peshtigo rivers was 102-165 cm (G. Kornely, Wisconsin Department of Natural Resource, 101 N. Ogden Rd., Peshtigo, Wisconsin, 54157, unpubl. data), and average lengths of males and females in the Fox River in 2000 was 150 cm and 164 cm, respectively (RFE, unpubl. data).

There are several indications that lake sturgeon are increasing in abundance in Lake Michigan. Sightings and incidental catches of fish in tributaries throughout the lake are increasing, although this could be due in part to increased interest and reporting. In rivers having established spawning runs, numbers observed increased in recent years. In some cases, this increase has been in conjunction with improved flow regimes

instituted. In the Manistee River, several strong year classes have been produced in the last 12 years following establishment of run-of-the-river flows (D.L. Peterson, University of Georgia, Warnell School of Forestry

commitment on the part of all of the Lake Michigan management agencies.

Recommendations

Several rehabilitation plans for lake sturgeon have been developed recently, and they include the Lake Sturgeon Rehabilitation Strategy for the State of Michigan (Hay-Chmielewski and Whelan 1997), Wisconsin's Lake Sturgeon Management Plan (Wisconsin Department of Natural Resources 2000), and the Draft Lake Sturgeon Plan for the Green Bay Basin. These plans identify major impediments to lake sturgeon rehabilitation and lay a framework for future management efforts and research needs. A recent Great Lakes lake sturgeon workshop, sponsored by the Great Lakes Fisheries Trust, identified the research needs for the Great Lakes with emphasis on Lake Michigan. Primary research needs for lake sturgeon identified in each of these plans and at the workshop include status assessments that provide inventories of populations, identification of habitat requirements and habitat availability for various life-history stages, the design of effective passage around artificial barriers (dams), and appropriate artificial-propagation strategies.

Several actions and initiatives are under way that will enhance lake sturgeon rehabilitation:

- Increased requirements for run-of-the-river flows at hydropower facilities
- Development of fish passage technologies that will safely pass lake sturgeon both upstream and downstream around dams and hydroelectric facilities
- Genetic analysis of remnant stocks in Lake Michigan and the Great Lakes
- Continuation of a new protocol to maximize protection of lake sturgeon during lampricide applications in rivers with known populations
- Development and refinement of culture and propagation techniques for lake sturgeon

- New harvest restrictions in Michigan and Wisconsin banning harvest throughout the lake and its tributaries, except in the Menominee River where limited harvest is allowed during an alternate-year, fall hook-and-line recreational fishery
- Ongoing population, spawning, and tagging assessments to gather basic biological information on distribution, status, and behavior

Burbot

Although abundant in the Great Lakes, burbot is not often recreationally or commercially harvested. In the sport fishery, most burbot are captured incidentally by ice anglers, and they rank low in angler preference (Quinn 2000). Large-scale commercial fisheries have not been established, perhaps because burbot are best eaten fresh; the meat becomes tough when frozen and has a rubbery texture when thawed (Becker 1983). Burbot is a freshwater member of the cod family, and many people who have tasted fresh burbot appreciate it for its lobster-like taste (Paragamian 2000). In northern Europe, burbot roe is a delicacy, and its liver is prized because of its high vitamin A and D content (Scott and Crossman 1973).

Burbot populations in Lake Michigan experienced dramatic changes in the last 50 years. Predation by sea lamprey drastically reduced burbot populations during the 1940s and 1950s. Swink and Fredericks (2000) found that mortality of small burbot due to attacks by sea lamprey was greater than that observed for small lake trout, and that larger burbot experienced mortalities similar to those of larger lake trout. However, sea lamprey control programs initiated in the 1960s resulted in the steady recovery of burbot populations. The FCO for burbot in Lake Michigan is to maintain self-sustaining stocks, and this is being achieved lakewide. However, the recent increase in burbot abundance raised concerns regarding the negative influence this may have on the fish community (e.g., increased predation on lake trout and alewife) (Eshenroder et al. 1995).

Burbot densities in Lake Michigan are high relative to other systems in the world and are increasing. The highest published densities for burbot (139 individuals per ha) are from Julian's Reef in the southern basin in 1990 (Edsall et al. 1993; McPhail and Paragamian 2000). The bycatch of

burbot in commercial fisheries in Green Bay almost quintupled during 1980-1985 (Rudstam et al. 1995). Trawl surveys conducted lakewide during 1973-1999 by the United States Geological Survey, Great Lakes Science Center, indicate that the CPE of burbot has been increasing since 1983, and that the highest burbot abundance was in 1998 (Fig. 23). A similar trend of increasing abundance was seen in bottom gillnet surveys conducted during 1984-1997 by the MDNR in the northern, eastern, and Grand Traverse Bay regions of Lake Michigan (JLJ, unpubl. data).



Fig. 23. CPE of burbot in Lake Michigan trawl surveys conducted by the United States Geological Survey, Great Lakes Science Center, 1973-1999.

The age structure and longevity of burbot indicates a population recovery. The majority of burbot collected during 1996 and 1997 in bottom gillnet surveys were between the ages of 9 and 16 and ranged up to age 22 (JLJ, unpubl. data).

Knowledge of burbot foraging dynamics is important for evaluating their interactions in the fish community. Diet studies were conducted in western waters in 1986-1988 by the WDNR and University of Wisconsin-Stevens Point (Fratt et al. 1997) and in eastern waters in 1996-1997 by the MDNR and Central Michigan University (Hart 2001).

Recommendations

1. Determine burbot consumption and its effect on the forage community and on other predators, especially lake trout.

Sea Lamprey

Dennis S. Lavis
United States Fish and Wildlife Service
Ludington Biological Station
229 South Jebavy Drive
Ludington, Michigan, U.S.A. 49431

Control of sea lampreys made possible the socioeconomic and biological revitalization of Lake Michigan and

limited due to a variety of mechanical, physical, and biological problems. With the exception of six streams, electrical barriers were operated for only a few years; only three remained operational after 1960 to measure annual changes in the number of spawning migrants, and all were dismantled by 1966.

Chemical treatment of tributaries with TFM began in 1960, and most sea lamprey producing streams had been treated once by 1966. The resultant destruction of larval populations is reflected in the 80-90% decline in the number of adults captured at index electric weirs by 1966.

Larval Populations and Production Areas

Sea lampreys have been found in 121 of the 511 tributaries of Lake Michigan. Estimates of larval populations in individual streams and production of juvenile (metamorphosed) sea lampreys are used to compare streams. Streams are selected for treatment based on their treatment cost versus their estimated production of parasitic-phase sea lampreys. Most larvae occur in 36 tributaries, and those larvae that survive treatment in these streams are responsible for the majority of sea lampreys in the lake; treatments are 95-98% effective. Larvae escape treatment due to the presence of oxbows, backwaters, and groundwater influx where minimum lethal concentrations of lampricide are difficult to maintain. Some sea lampreys are produced from streams that are too small for cost-effective treatment with lampricide. Minor production also occurs from a few (<5) small lentic areas.

Adult Populations

From 1977 to the present, traps fished in 12-14 index tributaries have been used to measure abundance of spawning-phase sea lampreys; however, since 1986, the number of spawning-phase sea lampreys in the Manistique River has been estimated using a stratified, random mark-and-recapture technique. Prior to 1996, the traps were used only as a measure of relative abundance, but since then, total abundance of spawning-phase sea lampreys entering tributaries has been estimated. The estimate is based on the efficiency of traps in the index tributaries and is expanded to other major sea lamprey producing tributaries based on average stream discharge. The whole-lake estimate has ranged from

57,615 (1996) to 92,430 (1998). There appears to be a tread of increasing population levels since estimates were initiated in 1996 (Fig. 25).

southern jurisdictions remained virtually unchanged over the past 23-25 years.

Sea lamprey induced mortality can be computed from a statistical relation between the number of Type AI wounds per fish and the probability of a lake trout surviving a single sea lamprey attack (Swink and Hanson 1986; Swink 1990). The lakewide, sea lamprey induced mortality averaged less than 7% most years during 1984-1998 but increased steadily in the 1990s, especially among large lake trout in

annually has been reduced from 21 per year during 1970-1979 to 12 per year during 1990-1999. The reduction is due in part to sea lampreys not reestablishing in some streams following an initial lampricide treatment and to changes in the criteria used in the stream selection process. Prior to 1995, streams were selected for treatment on the basis of relative abundance of larval sea lamprey. From 1995 to present, estimates of the production of juveniles (metamorphosed larvae) allowed for selection of streams based on the cost to treat a stream and its estimated production to the lake (Gavin Christie, Great Lakes Fishery Commission, 2100

dwest
Environmental Sciences Center, 2630 Fanta Reed Road, LaCrosse, WI,

in TFM use through the implementation of

. and Canadian Sea Lamprey Barrier
Coordinators were appointed in 1993 and charged with preparing a Great

Sterile-Male-Release Technique (SMRT)

The release of sterilized male sea lampreys in Great Lakes tributaries is still experimental and has not seen wide-scale application to date. The technique has not been applied to any Lake Michigan tributary. However, spawning-phase sea lampreys are captured in Lake Michigan streams and used in the SMRT program elsewhere in the Great Lakes, primarily in the St. Marys River.

Fish Health

Greg Wright
Intertribal Fisheries Assessment Program
Chippewa Ottawa Resources Authority
186 East Three Mile Road
Sault Ste. Marie, Michigan, U.S.A. 49783

Dale C. Honeyfield
United States Geological Survey
Research and Development Laboratory
ent LThree Mile Road Canadian C 57re for Inl 10.W8 0rs Tf0 Tc 0 Tw 12 0

Bacterial Kidney Disease (BKD)

BKD is caused by the bacterium Rs and is a slowly progressive, systemic infection that occurs primarily in salmonids (Lasee 1995). BKD can be found all over the world affecting both cultured and wild-ranging fish (Sanders and Fryer 1980). The pathogen can be transmitted from fish to fish (Mitchum and Sherman 1981) or from adults to eggs (Bullock et al. 1978; Bullock 1980). Outbreaks of BKD are thought to be stress related (Piper et al. 1982; Lasee 1995). No adult chinook salmon in a monitored spawning run in a Wisconsin tributary tested positive for the Rs bacteria prior to 1986; however, by 1988, over 66% of the fish returning to spawn tested positive, and many fish were showing overt clinical signs of disease (Fig. 27). Rates decreased after 1988 and averaged just above 3% since 1993 with very few fish showing overt clinical signs. The slow, progressive nature of this infection is disarming, which makes it difficult to determine when the pathogen will initiate a virulent disease. Understanding the relation between the pathogen and disease should be the primary focus for fish-health experts working in the Great Lakes.



Fig. 27. Prevalence of *Renibacterium salmoninarium* in chinook salmon (number testing positive/number tested using Direct Fluorescent Antibody Technique) returning to the Strawberry Creek, Wisconsin, spawning weir in northwestern Lake Michigan, 1982-2001. Data from Holey et al. (1998) and S. Marcquenski, Wisconsin Department of Natural Resources, 101 S. Webster St. Madison, Wisconsin, 53703, personal communication.

declines in lipids indicate fish trying to reestablish a normal metabolism or overcome stress. Many diseases develop and spread because the fish can no longer manage stress. Lake Michigan will be stressful for those exotic species that evolved in warmer environments. Therefore, as long as exotic species are the primary forage and predator species, managing Lake Michigan fisheries will continue to be challenging.

beneficial uses have been identified as impaired (Lake Michigan Technical Coordinating Committee 2000). Most of the remediation activities completed in Lake Michigan AOCs have been the removal of contaminated sediments, primarily those with PCBs. Extensive sediment removal projects have been completed on the lower Fox River, Sheboygan River, Waukegan Harbor, Grand Calumet River, Kalamazoo River, and Manistique River. A total of 453,600 kg of PCBs was removed from Waukegan Harbor, and sediments containing PCB concentrations in excess of 300 ppm were removed from the Sheboygan and lower Fox River. In addition, RAPs resulted in the removal of the first barrier to the lake on the Milwaukee River (North Avenue Dam) and in the stocking of Great Lakes muskellunge in Green Bay (BB, personal communication 1997). A more-detailed description of completed remediation activities for each AOC can be found at http://www.epa.gov/glnpo/aoc/.

Table 3. Lake Michigan Areas of Concern and the number of beneficial-use impairments per site (http://www.epa.gov/glnpo/aoc/).*

Area of Concern	Location	Impairments per site
Wisconsin	Lower Green Bay and Fox River	10
	Sheboygan River	8
	Milwaukee Estuary	11
Illinois	Waukegan Harbor	5
Indiana	Grand Calumet River	14
Michigan	Kalamazoo River	8
	White Lake	8
	Muskegan Lake	5
	Manistique River	5
	Menominee River	7

Designation of AOCs and the development of RAPs is the process that the GLWQA created to identify impaired beneficial uses and how to remediate the impairment. The Superfund and the Natural Resources Damage Assessment (NRDA) provisions of the Comprehensive Environmental Response, Compensation and Liabilities Act of 1980 are

two tools available to water-quality and fishery agencies to achieve goals identified for a given AOC. The Superfund addresses primarily human health issues and resulted in the rem

continued steady decline in PCB concentrations for brown trout, rainbow trout, coho salmon, and chinook salmon, and possible constant levels for lake trout.

PCBs no longer directly affect the ability of lake trout and, likely, other salmonines to produce viable eggs. When both PCB and thiamine concentrations were measured from the same lake trout eggs collected from Lake Michigan from 1996-1998, only thiamine concentrations were correlated with swim-up mortality (Stratus Consulting Inc. 1999). Fitzsimons (1995) also concluded that early mortality was not solely related to the contaminants measured in eggs or larvae. Lake trout embryos from Lake Michigan still have elevated mortalities associated with low egg-thiamine concentrations, and the mechanism or pathway for the lowered thiamine concentrations remains unknown (Honeyfield et al. 1998a; Stratus Consulting Inc. 1999).

Flow Management through Hydropower Dams

The water-flow regime through dams can significantly affect fishery habitat below dams (Hayes 1999). Flow regimes during peaking operations often result in large daily fluctuations in water flow below dams that dewater stream channels and negatively affect fish populations. Flow management that requires a run-of-the-river (i.e., flows below a dam have the same flow pattern as above the dam) operation often results in more-favorable habitat below the dam.

The benefit of run-of-the-river flow management for salmon spawning and nursery habitat is illustrated by the number of chinook salmon smonTf0dD 4 >>BD98 c0m1(1i-16.792

Open Forum

Mark E. Holey
United States Fish and Wildlife Service
Green Bay Fishery Resources Office
2661 Scott Tower Drive

from the lake, the alternatives to alewives need to be considered, because alewives occupy an important niche in the lake's food web. The native species that previously filled this niche were lake herring and deepwater ciscoes.

"If alewives are evil, and the primary control is chinook salmon, are chinook salmon good?"

Few doubt that the chinook salmon played a major role in restoring a more-healthy balance between predators and prey fishes, and this outcome led to increased integrity of the lake's fish community. The pronounced naturalization of chinook salmon is very positive in terms of system health, but the sustainability of chinook salmon remains a concern, especially in view of the spectacular die-off of the species in the late 1980s. Chinook salmon apparently suffers increased mortality during periods of low alewife abundance. Although the chinook salmon, among all of the species of trout and salmon in the lake, is the most efficient predator of alewife, it also appears to be less capable of switching to an alternative prey species when alewives are scarce. This dilemma poses a tough management question for the LMC: what population levels of alewife and chinook salmon should managers seek to achieve? Answering this question will require considerable input from the public, especially from those who fish the lake, regarding their views on a desired fish community.

"Has the relative importance of the species sought by sport fishermen changed?"

Based on the state-of-the-lake presentations to the LMC, more anglers pursued yellow perch than chinook salmon, suggesting a shift in the socioeconomic values of these two species. The economic value of the offshore trout and salmon fishery, however, still exceeds the value of the yellow perch fishery, which operates inshore, even though yellow perch anglers outnumber those fishing for trout and salmon, and the catch in numbers of yellow perch exceeds that of salmon and trout. Nevertheless, the trend since the 1970s has been a shift abul9frbeen irors roon salmch in

"Should we be reintroducing non-sport fish, especially a species like lake herring, into the lake?"

situation in Lake Michigan will probably not turn around until a larger population of lake trout that can respond to more-favorable environmental conditions is established. If such conditions do occur and lake trout do respond, lake trout will likely take over and drive the system as has occurred in Lake Superior.

"Why is so much attention focused on stocking more fish and not on identifying and addressing the bottlenecks that may be limiting natural reproduction?"

Other potential bottlenecks cannot be addressed until the low-population-level bottleneck is overcome first, because large-scale reproduction cannot be achieved with the paltry number of adult fish now surviving to a reproductive age. Bottlenecks or gaps in assessment data, not only for lake trout but also for yellow perch, lake sturgeon and other species, do need to be identified.

"Would the economics of the hatchery-based fishery in Lake Michigan weaken the political will to focus on sustainability and the flexibility needed to remain committed to achieving lake trout rehabilitation?"

Economic pressures to maintain a particular fishery can be so great that fish managers have little leeway to attempt other strategies.

"Is the information available on phosphorus levels and inputs adequate?"

Both the IJC and the GLFC recommended continuation of the phosphorus-loading trend data, especially for Lake Erie, and the same recommendation could be made for Lake Michigan. In Lake Ontario, trend data for phosphorus and lower trophic levels were crucial in enabling biologists to understand how increasing pressure at the top of the food web and the diminishing productivity at the bottom were creating an unsustainable food web.

Literature Cited

Adams, S.M. 2000. Ecological role of lipids in the health and success of fish populations. *In* Lipids in freshwater ecosystems. *Edited by* M.T. Arts, and B.C. Wainman, Springer, New York, New York. pp. 132-160.

Allen, P.J. 2000. A computer simulation model for the yellow perch population in the Indiana waters of Lake Michigan. Master's Thesis. Ball State University, Muncie, Indiana.

Argyle, R.L., Fleischer, G.W., Curtis, G.L., Adams, J.V., and Stickel, R.G. 1998. An integrated acoustic and trawl based prey fish assessment strategy for Lake Michigan. A report to the Illinois Department of Natural Resources, Indiana Department of Natural Resources, Michigan Department of Natural Resources and Wisconsin Department of Natural Resources. U.S. Geological Survey—Great Lakes Science Center, Ann Arbor, MI.

Baldwin, N.S., Saalfeld, R.W., Ross, M.A., and Buettner, H.J. 1979. Commercial fish production in the Great Lakes 1867-1977. Great Lakes Fish. Comm. Tech. Rep. 3.

Barbiero, R.P., and Tuchman, M.L. 2002. Results from GLNPO's biological open water surveillance program of the Laurentian Great Lakes 1999. U.S. Environmental Protection Agency, Great Lakes National Program Office, Chicago, IL, EPA-905-R-02-001.

Barbiero, R.P., and Tuchman, M.L. 2004. Changes in the crustacean communities of Lakes Michigan, Huron and Erie following the invasion of the predatory cladoceran *Bythotrephes longimanus*. Can. J. Fish. Aquat. Sci.

Barbiero, R.P., Little, R.E., and Tuchman, M.L. 2001. Results from the U.S. EPA's biological open water surveillance program of the Laurentian Great Lakes: III. Crustacean zooplankton. J. Great Lakes Res. **27**:167-184.

Barbiero, R.P., Schacht, L.L., Little, R.E., and Tuchman, M.L. 2005. Crustacean zooplankton communities in Lake Michigan. *In* State of Lake Michigan: ecology, health and management. *Edited by* T. Edsall, and M. Munawar. Ecovision World Monograph Series. SBP Academic Publishing, Amsterdam, The Netherlands.

Barbiero, R.P., Tuchman, M.L., Warren, G.J., and Rockwell, D.C. 2002.

Bills, T.D., and Johnson, D.A. 1992. Effect of pH on the toxicity of TFM

Evans, M.S. 1986. Recent major declines in zooplankton populations in the inshore region of Lake Michigan: probable causes and implications. Can. J. Fish..Aquat. Sci. **43**: 154-159.

Evans, M.S., and Jude, D.J. 1986. Recent shifts in *Daphnia* community in Southeastern Lake Michigan: a comparison of the inshore and offshore regions. Limnol. Oceanogr. **31**: 56-67.

Evans, W.C. 1975. Thiaminases and their effects on animals. Vitam. Horm. **33**: 467-504.

Fahnenstiel, G.L., and Scavia, D. 1987. Dynamics of Lake Michigan phytoplankton: recent changes in surface and deep communities. Can. J. Fish. Aquat. Sci. **44**: 509-514.

Fetterolf, Jr., C.M. 1980. Why a Great Lakes Fishery Commission and why a sea lamprey international symposium. Can. J. Fish. Aquat. Sci. **37**: 1588-1593.

Fisher, J.P., Spitsbergen, J.M., Lamonte, T., Little, E. E., and DeLonay, A. 1995. Pathological and behavioral manifestations of the "Cayuga Syndrome," a thiamine deficiency in larval landlocked Atlantic salmon. J. Aquat. Anim. Health **7**:269-283.

Fitzsimons, J.D. 1995. The effect of B-vitamins on swim-up syndrome in Lake Ontario lake trout. J. Great Lakes Res. **21** (Suppl. 1): 286-289.

Fitzsimons, J.D., Brown, S.B., and Vandenbyllaardt, L. 1998. Thiamine levels in food chains of the Great Lakes. *In* early life stage mortality syndrome in fishes of the Great Lakes and the Baltic Sea. *Edited by* G. McDonald, J.D. Fitzsimons, and D.C. Honeyfield. American Fisheries Society Symposium 21, Bethesda, MD. pp. 90-98

Flath, L.E., and Diana, J.S. 1985. Seasonal energy dynamics of the alewife in southeastern Lake Michigan. Trans. Am. Fish. Soc. **114**: 28-337.

Hansen, M.J., and Holey, M.E. 2001. Ecological factors affecting the sustainability of chinook and coho salmon populations in the Great Lakes, especially Lake Michigan. *In* Sustainable salmon fisheries: binational perspectives. *Edited by* K.D. Lynch, M.L. Jones, and

International Joint Commission. 1980. Phosphorus management for the

Lake Michigan Technical Coordinating Committee. 2000. Lake Michigan Lakewide Management Plan. U.S. Environmental Protection Agency, Chicago. <http://www.epa.gov/glnpo/lakemich/index.html>.

Lamon, III, E.C., Carpenter, S.R., and Stow, C.A. 1998. Forecasting PCB concentrations in Lake Michigan salmonids: a dynamic linear model approach. Ecol. Appl. **8**: 659-668.

Moreau, J., Nyakegeni, B., and Pearce, M. 1993. Trophic relationships in the pelagic zone of Lake Tanganyika (Burendi Sector). *In* Trophic models of ecosystems. *Edited by* V. Christensen and D. Pauly. Int. Cent. Living Aquat. Resourc. Manage. (ICLARM) Conference Proceedings 26. pp. 138-143.

National Research Council. 1993. Nutrient requirements of fish. Committee on Animal Nutrition. National Academy Press, Washington, DC.

O'Gorman, R., and Stewart, T.J. 1999. Ascent, dominance, and decline of the alewife in the Great Lakes: food web interactions and management strategies. *In* Great Lakes fisheries policy and management: a binational perspective. *Edited by* W.W. Taylor and C.P. Ferreri. Michigan State University Press, East Lansing, MI. pp. 489-513

O'Neal, R. P. 1997. Muskegon River watershed assessment. Michigan Department of Natural Resources, Fisheries Division Special Report 19, Ann Arbor, Michigan.

Paragamian, V.L. 2000. Preface. *In* Burbot biology, ecology and management. *Edited by* V.L. Paragamian, and D.W. Willis. Fisheries Management Section of the American Fisheries Society, Spokane, WA. p. 7.

Parsons, J.W. 1973. History of salmon in the Great Lakes, 1850-1970. U. S. Fish Wildl. Serv. Tech. Pap. 68.

Patriache, M. 1980. Movement and harvest of coho salmon in Lake Michigan, 1978-1979. Mich. Dept. Nat. Resour., Fish. Res. Rep. 1889, Ann Arbor, MI.

Peck, J.W. 1979. Utilization of traditional spawning reefs by hatchery lake trout in the upper Great Lakes. Mich. Dept. Nat. Resour., Fish. Rep. 1871, Lansing, MI.

Perkins, D.L., and Krueger, C.C. 1995. Dynamics of production by hatchery-origin lake trout (*Salvelinus namaycush*
Reef, Lake Ontario. J. Great Lakes Res., **21** (Suppl. 1): 400-417.

Rybicki, R.W., and D. F. Clapp. 1996. Diet of chinook salmon in eastern Lake Michigan. Michigan Department of Natural Resources, Fisheries Division, Technical Report 2027, Ann Arbor, Michigan.

Sanders, J.E., and Fryer, J.L. 1980. *Renibacterium salmoninarum* gen. nov., sp. nov., the causative agent of bacterial kidney disease in salmonid fishes. Int. J. Syst. Bacteriol. **30**: 496-502.

Sawyer, A. J. 1980. Prospects for integrated pest management of the sea lamprey (*Petromyzon marinus*). Can. J. Fish. Aquat. Sci. **37**: 2081-2089.

Schelske, C.L. 1988. Historic trends in Lake Michigan silica concentrations. Int. Rev. Gesamten Hydrobiol. **73**: 559-591.

Schelske, C.L., and Stoermer, E.F. 1971. Eutrophication, silica depletion and predicted changes in algal quality in Lake Michigan. Science **173**: 423-424.

Schneeberger, P.J. 2000. Population dynamics of contemporary yellow perch and walleye stocks in Michigan waters of Green Bay, Lake Michigan, 1988-96. Michigan Department of Natural Resources, Fisheries Research Report 2055, Ann Arbor, MI.

Schneider, J.C., and Leach, J.H. 1979. Walleye stocks in the Great Lakes, 1800-1975: fluctuations and possible causes. Great Lakes Fish. Comm. Tech. Rep. 31.

Scott, W.B., and Crossman, E.J. 1973. Freshwater fishes of Canada. Bull. Fish. Res. Board Can. 184.

Seelbach, P.W. 1989. Characteristics of adult steelhead populations, including returns of hatchery yearlings, in the St. Joseph and Grand rivers, Michigan 1979-85. Michigan Department of Natural Resources, Fisheries Technical Report 89-3, Ann Arbor, MI.

Seelbach, P.W. 1993. Population biology of steelhead in a stable-flow, low-gradient tributary of Lake Michigan. Trans. Am. Fish. Soc. **122**: 179-198.

Seelbach, P.W., and Whelan, G.E. 1988. Identification and contribution of wild and hatchery steelhead stocks in Lake Michigan tributaries. Michigan Department of Natural Resources, Fisheries Research Report 1950, Lansing, MI.

Seelbach, P.W., Dexter, J.L., and Ledet, N.D. 1994. Performance of steelhead smolts stocked in southern Michigan warmwater rivers. Michigan Department of Natural Resources, Fisheries Research Report 2003, Lansing, MI.

Selgeby, J.H., Bronte, C.R., Brown, Jr., E.H., Hansen, M.J., Holey, M.E., VanAmberg, J.P., Muth, K.M., Makauskas, D.B., McKee, P.C., Anderson, D.M., Ferreri, C.P., and Schram, S.T. 1995. Lake trout restoration in the Great Lakes: stock size criteria for natural reproduction. J. Great Lakes Res. **21** (Suppl. 1): 498-504.

Sherman, K., and Duda, A.M. 1999. Large marine ecosystems: an emerging paradigm for fishery sustainability. Fisheries (Bethesda) **24**(12): 15-26.

Shroyer, S.M., and McComish, T.S. 2000. Relationship between alewife abundance and yellow perch recruitment in southern Lake Michigan. N. Am. J. Fish. Manage. **20**: 220-225.

Smith, S.H. 1968. Species succession and fishery exploitation in the Great Lakes. J. Fish. Res. Board Can. **25**: 667-693.

Sonzogni, W.C., Richardson, W., Rodgers, P., and Monteith, T.J. 1983. Chloride pollution of the Great Lakes. J. Water Poll. Control Fed. **55**: 513-521.

Sprules, W.G., Brandt, S.B., Steward, D.J., Munawar, M., Jin, E.H., and Love, J. 1991. Biomass size spectrum of the Lake Michigan pelagic food web. Can. J. Fish. Aquat. Sci. **48**: 105-115.

Stewart, D.J., and Ibarra, M. 1991. Predation and production by salmonine fishes in Lake Michigan, 1978-88. Can. J. Fish. Aquat. Sci. **48**: 909-922.

Stratus Consulting Inc. 1999. Reproductive injuries to lake trout. *In* Injuries to fishery resources, lower Fox River/Green Bay natural resources damage assessment: final report. Stratus Consulting Inc, Boulder, CO., November 8, 1999. pp. 74-117.

Swink, W. D. 1990. Effect of lake trout size on survival after a single sea lamprey attack. Trans. Am. Fish. Soc. **119**: 996-1002.

Swink, W.D., and Fredericks, K.T. 2000. Mortality of burbot from sea lamprey attack and initial analyses of burbot blood. *In* Burbot biology, ecology and management. *Edited by* V.L. Paragamian, and D.W. Willis. Fisheries Management Section of the American Fisheries Society, Spokane, Washington. pp. 147-154.

Swink, W.D., and Hanson, L.H. 1986. Survival from sea lamprey (*Petromyzon marinus*) predation by two strains of lake trout (*Salvelinus namaycush*). Can. J. Fish. Aquat. Sci. **43**: 2528-2531.

Taylor, W.W., Smale, M.A., and Freeberg, M.H. 1987. Bio(rey)-7(o 162 h. Aq66 ref/Pl analy)-7(ses of bu

Tonello, M.A. 1997. Population structure and dynamics of alewives and spottail shiners in Indiana waters of Lake Michigan, 1984-1996. Master's thesis. Ball State University, Muncie, IN.

Trudeau, T. 1995. Status of coho management. *In* Minutes Lake Michigan Committee (1995 Annual Meeting), Ann Arbor, Michigan, Great Lakes Fishery Commission, March 29, 1995.

United States Environmental Protection Agency. 2003. Sampling and analytical procedures for GLNPO's open water quality survey of the Great Lakes. EPA Rept. No. R05-R-03-002.

Wagner, W.C. 1981. Reproduction of planted lake trout in Lake Michigan. N. Am. J. Fish. Manage. **1**: 159-164.

Wells, L., and McLain, A.L. 1972. Lake Michigan: effects of exploitation, introductions, and eutrophication of the salmonid community. J. Fish. Res. Board Can. **29**: 889-898.

Wisconsin Department of Natural Resources. 2000. Wisconsin's Lake Sturgeon Management Plan. 12 pages. Bureau of Fisheries Management and Habitat Protection, Madison, WI.

Zajicek, J.L., Tillitt, D.E., Honeyfield, D.C., Brown, S.B., and Fitzsimons, J.D.. 2004. A method for measuring thiaminase I in fish tissues. J. Aquat. Anim. Health.

90-4 An ecosystem approach to the integrity of the Great Lakes in turbulent times (proceedings of a 1988 workshop supported by the Great Lakes Fishery Commission and the Science Advisory Board of the International Joint Commission). 1990. Edited by C. J. Edwards and H. A. Regier. 302 p.

91-1 Status of walleye in the Great Lakes: case studies prepared for the 1989 workshop. 1991. Edited by P. J. Colby, C. A. Lewis, and R. L. Eshenroder. 222 p.

91-2 Lake Michigan: an ecosystem approach for remediation of critical pollutants and management of fish communities (report of a round table sponsored in 1990 by the Great Lakes Fishery Commission, the Science Advisory Board of the International Joint Commission, and the Lake Michigan Federation). 1991. Edited by R. L. Eshenroder, J. H. Hartig, and J. E. Gannon. 58 p.

91-3 The state of the Lake Ontario fish community in 1989. 1991. S. J. Kerr and G. C. LeTendre. 38 p.

93-1 Great Lakes fish disease control policy and model program. 1993. Edited by J. G. Hnath. 38 p.

Protocol to minimize the risk of introducing emergency disease agents with importation of salmonid fishes from enzootic areas. 1993. Edited by R.W. Horner and R. L. Eshenroder. 15 p.

94-1 The state of Lake Superior in 1992. 1994. Edited by M. J. Hansen. 110 p.

94-2 An introduction to economic valuation principles for fisheries management. L. G. Anderson. 98 p.

95-1 Fish-community objectives for Lake Huron. 1995. R. L. DesJardine, T. K. Gorenflo, R. N. Payne, and J. D. Schrouder. 38 p.

95-2 The state of Lake Huron in 1992. Edited by M. P. Ebener. 140 p.

95-3 Fish-community objectives for Lake Michigan. R.L. Eshenroder, M.E. Holey, T.K. Gorenflo, and R.D. Clark, Jr. 56 p.

99-1 Fish-community objectives for Lake Ontario. T.J. Stewart, R.E. Lange, S.D. Orsatti, C.P. Schneider, A. Mathers, M.E. Daniels. 56 p.

03-1 Fish-community objectives for Lake Superior. W.H. Horns, C.R. Bronte, T.R. Busiah, M.P. Ebener, R.L. Eshenroder, T. Groenflo, N. Kmiecik, W. Mattes, J.W. Peck, M. Petzold, D.R. Schreiner. 86 p.

03-2 Fish-community goals and objectives for Lake

Special Publications

79-1 Illustrated field guide for the classification of sea lamprey attack marks on Great Lakes lake trout. 1979. E. L. King and T. A. Edsall. 41 p.

82-1 Recommendations for freshwater fisheries research and management from the Stock Concept Symposium (STOCS). 1982. A. H. Berst and G. R. Spangler. 24 p.

82-2 A review of the adaptive management workshop addressing salmonid/lamprey management in the Great Lakes. 1982. Edited by J. F. Koonce, L. Greig, B. Henderson, D. Jester, K. Minns, and G. Spangler. 58 p.

82-3 nd T. G. Carey. 262 p.

84-1 Recommendations for standardizing the reporting of sea lamprey marking data. 1984. R. L. Eshenroder and J. F. Koonce. 22 p.

84-2 Working papers developed at the August 1983 conference on lake trout research. 1984. Edited by R. L. Eshenroder, T. P. Poe, and C. H. Olver.

84-3 Analysis of the response to the use of "Adaptive Environmental Assessment Methodology" by the Great Lakes Fishery Commission. 1984. C. K. Minns, J. M. Cooley, and J. E. Forney. 22 p.

85-1 Lake Erie fish community workshop (report of the April 4-5, 1979 meeting). 1985. Edited by J. R. Paine and R. B. Kenyon. 58 p.

85-2 A workshop concerning the application of integrated pest management (IPM) to sea lamprey control in the Great Lakes. 1985. Edited by G. R. Spangler and L. D. Jacobson. 98 p.

85-3 Presented papers from the Council of Lake Committees plenary session on Great Lakes predator-prey issues, March 20, 1985. 1985. Edited by R. L. Eshenroder. 134 p.

85-4 Great Lakes fish disease control policy and model program. 1985. Edited by J. G. Hnath. 24 p.

85-5 Great Lakes Law Enforcement/Fisheries Management Workshop (report of the 21, 22 September 1983 meeting). 1985. Edited by W. L. Hartman and M. A. Ross. 26 p.

85-6 TFM vs. the sea lamprey: a generation later. 1985. 18 p.

86-1 The lake trout rehabilitation model: program documentation. 1986. C. J. Walters, L. D. Jacobson, and G. R. Spangler. 34 p.

87-1 Guidelines for fish habitat management and planning in the Great Lakes (report of the Habitat Planning and Management Task Force and Habitat Advisory Board of the Great Lakes Fishery Commission). 1987. 16 p.

87-2 Workshop to evaluate sea lamprey populations "WESLP" (background papers and proceedings of the August 1985 workshop). 1987. Edited by B. G. H. Johnson.

87-3 Temperature relationships of Great Lakes fishes: a data compilation. 1987. D. A. Wismer and A. E. Christie. 196 p.

88-1 Committee of the Whole workshop on implementation of the Joint Strategic Plan for